

The Extension Dimension of Subcategories and Recollements of Abelian Categories

Xin MA

College of Science, Henan University of Engineering, Zhengzhou 451191, P. R. China

E-mail: maxin@haue.edu.cn

Ye Yang PENG Zhao Yong HUANG¹⁾

Department of Mathematics, Nanjing University, Nanjing 210093, P. R. China

E-mail: pengyy@smail.nju.edu.cn huangzy@nju.edu.cn

Abstract We investigate the behavior of the extension dimension of subcategories of abelian categories under recollements. Let $\Lambda', \Lambda, \Lambda''$ be artin algebras such that $(\text{mod } \Lambda', \text{mod } \Lambda, \text{mod } \Lambda'')$ is a recollement, and let \mathcal{D}' and \mathcal{D}'' be subcategories of $\text{mod } \Lambda'$ and $\text{mod } \Lambda''$ respectively. For any $n, m \geq 0$, under some conditions, we get $\dim \Omega^k(\mathcal{D}) \leq \dim \Omega^n(\mathcal{D}') + \dim \Omega^m(\mathcal{D}'') + 1$, where $k = \max\{m, n\}$ and \mathcal{D} is the subcategory of $\text{mod } \Lambda$ glued by \mathcal{D}' and \mathcal{D}'' ; moreover, we give a sufficient condition such that the converse inequality holds true. As applications, some results for Igusa–Todorov subcategories and syzygy finite subcategories are obtained.

Keywords Recollements, extension dimension, Igusa–Todorov subcategories, syzygy finite subcategories

MR(2010) Subject Classification 16E05, 16G10

1 Introduction

Given a triangulated category \mathcal{T} , Rouquier introduced in [25, 26] the dimension of \mathcal{T} under the idea of Bondal and van den Bergh in [7]. This dimension and the infimum of the Orlov spectrum of \mathcal{T} coincide, see [4, 20]. This dimension plays an important role in representation theory. For example, it can be used to compute the representation dimension of Artin algebras ([19, 25]). As an analogue of the dimension of triangulated categories, the extension dimension $\dim_{\mathcal{A}} \mathcal{D}$ of a subcategory \mathcal{D} of an abelian category \mathcal{A} was introduced by Beligiannis in [5], also see [10]. Let Λ be an Artin algebra. Note that the representation dimension of Λ is at most two (that is, Λ is of finite representation type) if and only if $\dim \text{mod } \Lambda (= \dim_{\text{mod } \Lambda} \text{mod } \Lambda) = 0$ ([5]). So, like the representation dimension of Λ , the extension dimension $\dim \text{mod } \Lambda$ is also an invariant that measures how far Λ is from of finite representation type. It was shown that the extension dimension is useful in studying the representation type of algebras and finitistic dimension conjecture ([33]).

Received September 22, 2021, accepted October 27, 2022

Supported by NSFC (Grant Nos. 11971225, 12171207, 12001168), Henan University of Engineering (Grant Nos. DKJ2019010, XYR-2021JZ001), the Key Research Project of Education Department of Henan Province (Grant No. 21A110006)

1) Corresponding author

Recollements of triangulated and abelian categories were introduced in [6, 11] in connection with derived categories of sheaves on topological spaces with the idea that one triangulated category may be “glued together” from two others. Recollements provide a useful reduction technique for some homological properties such as the finiteness of global dimension and finitistic dimension [9, 13, 21, 29], the Gorensteinness [1, 12, 17, 24] and the representation type and representation dimension of Artin algebras as well as the extension dimension of abelian categories [21, 33], and so on. Following the above philosophy, we will study the behavior of the extension dimension of certain subcategories of an abelian category under recollements.

For an Artin algebra Λ , we use $\text{mod } \Lambda$ to denote the category of finitely generated left Λ -modules. Let Λ' , Λ and Λ'' be Artin algebras such that there is a recollement of module categories:

$$\text{mod } \Lambda' \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \text{mod } \Lambda \begin{array}{c} \xleftarrow{j_!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \text{mod } \Lambda'' .$$

Our main results are as follows.

Theorem 1.1 (Theorem 3.8) *Let $(\text{mod } \Lambda', \text{mod } \Lambda, \text{mod } \Lambda'')$ be a recollement, and let \mathcal{D}' and \mathcal{D}'' be subcategories of $\text{mod } \Lambda'$ and $\text{mod } \Lambda''$ respectively. Assume that $i^!$ is exact. For any $n, m \geq 0$, if one of the following conditions holds:*

- (1) $m = 0$,
- (2) $m \geq 1$ and $i^!j_!$ preserves projective objects,

then

$$\dim_{\text{mod } \Lambda} \Omega^k(\mathcal{D}) \leq \dim_{\text{mod } \Lambda'} \Omega^n(\mathcal{D}') + \dim_{\text{mod } \Lambda''} \Omega^m(\mathcal{D}'') + 1,$$

where

$$k = \max\{m, n\} \quad \text{and} \quad \mathcal{D} = \{D \in \text{mod } \Lambda \mid i^!(D) \in \mathcal{D}' \text{ and } j^*(D) \in \mathcal{D}''\}.$$

Moreover, we have the following

Theorem 1.2 (Theorem 3.12) *Let $(\text{mod } \Lambda', \text{mod } \Lambda, \text{mod } \Lambda'')$ be a recollement, and let \mathcal{D} be a subcategory of $\text{mod } \Lambda$ with $i_*i^!(\mathcal{D}) \subseteq \mathcal{D}$ and $j_*j^*(\mathcal{D}) \subseteq \mathcal{D}$. If $i^!$ is exact, then*

$$\max\{\dim_{\text{mod } \Lambda'} \Omega^n(i^!(\mathcal{D})), \dim_{\text{mod } \Lambda''} \Omega^n(j^*(\mathcal{D}))\} \leq \dim_{\text{mod } \Lambda} \Omega^n(\mathcal{D})$$

for some $n \geq 0$.

Then we apply these results to Igusa–Todorov subcategories and syzygy finite subcategories. Some known results are obtained as corollaries. Finally, we give some examples to illustrate the obtained results.

Throughout this paper, all abelian categories have enough projective and injective objects and all subcategories are full, additive and closed under isomorphisms. All algebras are Artin algebras. Finally, we recall the notion of upper triangular matrix Artin algebras. Let Λ', Λ'' be Artin algebras and ${}_{\Lambda'}M_{\Lambda''}$ an (Λ', Λ'') -bimodule such that ${}_{\Lambda'}M$ and $M_{\Lambda''}$ are finitely generated, and let $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ be a triangular matrix algebra. Then Λ is an Artin algebra ([3, Proposition III.2.1]). A module in $\text{mod } \Lambda$ can be uniquely written as a triple $\begin{pmatrix} X \\ Y \end{pmatrix}_f$ with $X \in \text{mod } \Lambda'$, $Y \in \text{mod } \Lambda''$ and $f \in \text{Hom}_{\Lambda'}(M \otimes_{\Lambda''} Y, X)$ ([3, p.76]).

2 Preliminaries

Let \mathcal{A} be an abelian category, and let \mathcal{D} be a class of objects in \mathcal{A} . We use $\text{add } \mathcal{D}$ to denote the subcategory of \mathcal{A} consisting of direct summands of finite direct sums of objects in \mathcal{D} .

Let $\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n$ be classes of objects in \mathcal{A} . Define

$$\mathcal{U}_1 \diamond \mathcal{U}_2 := \text{add}\{A \in \mathcal{A} \mid \text{there exists an exact sequence} \\ 0 \longrightarrow U_1 \longrightarrow A \longrightarrow U_2 \longrightarrow 0 \text{ in } \mathcal{A} \text{ with } U_1 \in \mathcal{U}_1 \text{ and } U_2 \in \mathcal{U}_2\}.$$

Inductively, define

$$\mathcal{U}_1 \diamond \mathcal{U}_2 \diamond \dots \diamond \mathcal{U}_n := \text{add}\{A \in \mathcal{A} \mid \text{there exists an exact sequence } 0 \longrightarrow U \longrightarrow A \longrightarrow V \longrightarrow 0 \\ \text{in } \mathcal{A} \text{ with } U \in \mathcal{U}_1 \text{ and } V \in \mathcal{U}_2 \diamond \dots \diamond \mathcal{U}_n\}.$$

For a class \mathcal{U} of \mathcal{A} , set $\langle \mathcal{U} \rangle_0 := 0$, $\langle \mathcal{U} \rangle_1 := \text{add } \mathcal{U}$, $\langle \mathcal{U} \rangle_n := \langle \mathcal{U} \rangle_1 \diamond \langle \mathcal{U} \rangle_{n-1}$ for any $n \geq 2$, and $\langle \mathcal{U} \rangle_\infty := \bigcup_{n \geq 0} \langle \mathcal{U} \rangle_n$ ([5]). For subcategories \mathcal{U}, \mathcal{V} and \mathcal{W} of \mathcal{A} , by [10, Proposition 2.2], we have

$$(\mathcal{U} \diamond \mathcal{V}) \diamond \mathcal{W} = \mathcal{U} \diamond (\mathcal{V} \diamond \mathcal{W}).$$

Definition 2.1 ([5, 10, 33]) *For a subcategory \mathcal{D} of \mathcal{A} , the extension dimension $\text{dim}_{\mathcal{A}} \mathcal{D}$ of \mathcal{D} is defined as*

$$\text{dim}_{\mathcal{A}} \mathcal{D} := \inf\{n \geq 0 \mid \mathcal{D} \subseteq \langle T \rangle_{n+1} \text{ with } T \in \mathcal{A}\}.$$

When there is no ambiguity, we write $\text{dim } \mathcal{D} := \text{dim}_{\mathcal{A}} \mathcal{D}$ for short.

Let \mathcal{A} be an abelian category, and let $M \in \mathcal{A}$ and $m \geq 0$. We use $\Omega_{\mathcal{A}}^m(M)$ to denote the m -th syzygy of M ; in particular, $\Omega_{\mathcal{A}}^0(M) = M$. Let \mathcal{D} be a subcategory of \mathcal{A} . We use $\Omega_{\mathcal{A}}^m(\mathcal{D})$ to denote the full subcategory of \mathcal{A} consisting of those objects in \mathcal{A} that are either projective or direct summands of m -th syzygies of objects in \mathcal{D} . Dually, the m -th cosyzygy $\Omega_{\mathcal{A}}^{-m}(M)$ of M and the subcategory $\Omega_{\mathcal{A}}^{-m}(\mathcal{D})$ are defined.

Lemma 2.2 *Let \mathcal{A} be an abelian category and let $X, T \in \mathcal{A}$. If $X \in \langle T \rangle_n$, then for any $n \geq 1$ and $i \geq 0$, we have*

- (1) $\Omega_{\mathcal{A}}^i(X) \in \langle \Omega_{\mathcal{A}}^i(T) \rangle_n$.
- (2) $\Omega_{\mathcal{A}}^{-i}(X) \in \langle \Omega_{\mathcal{A}}^{-i}(T) \rangle_n$.

Immediately, we get the following result.

Lemma 2.3 *Let \mathcal{A} be an abelian category and \mathcal{D} a subcategory of \mathcal{A} . Then for any $m \geq n \geq 0$, we have*

$$\text{dim } \Omega_{\mathcal{A}}^m(\mathcal{D}) \leq \text{dim } \Omega_{\mathcal{A}}^n(\mathcal{D}).$$

Lemma 2.4 *Let \mathcal{A} be an abelian category and $n \geq 1$, and let*

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

be an exact sequence in \mathcal{A} . If there exist $T_1, T_2 \in \mathcal{A}$ such that $A \in \langle T_1 \rangle_n$ and $B \in \text{add } T_2$, then $C \in \langle T_2 \oplus \Omega_{\mathcal{A}}^{-1}(T_1) \rangle_{n+1}$.

Proof By [33, Lemma 3.2], we have the following exact sequence

$$0 \longrightarrow B \longrightarrow C \oplus I \longrightarrow \Omega_{\mathcal{A}}^{-1}(A) \longrightarrow 0$$

in \mathcal{A} with I injective. Then, by Lemma 2.2 and [33, Proposition 2.2(1) and Corollary 2.3(1)], we have

$$C \in \langle T_2 \rangle_1 \diamond \langle \Omega_{\mathcal{A}}^{-1}(T_1) \rangle_n \subseteq \langle T_2 \oplus \Omega_{\mathcal{A}}^{-1}(T_1) \rangle_{n+1}. \quad \square$$

We need the following easy and useful fact.

Lemma 2.5 *Let \mathcal{A} and \mathcal{B} be abelian categories and $n \geq 0$, and let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an exact functor. If F preserves projective objects, then*

$$\Omega_{\mathcal{B}}^n(F(X)) = F(\Omega_{\mathcal{A}}^n(X))$$

for any $X \in \mathcal{A}$.

Proof For any $X \in \mathcal{A}$, consider the following exact sequence

$$0 \longrightarrow \Omega_{\mathcal{A}}^n(X) \longrightarrow P_{n-1} \longrightarrow P_{n-2} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow X \longrightarrow 0$$

in \mathcal{A} with all P_i projective. Applying the functor F to it yields an exact sequence

$$0 \longrightarrow F(\Omega_{\mathcal{A}}^n(X)) \longrightarrow F(P_{n-1}) \longrightarrow F(P_{n-2}) \longrightarrow \cdots \longrightarrow F(P_0) \longrightarrow F(X) \longrightarrow 0$$

in \mathcal{B} with all $F(P_i)$ projective by assumption. Thus

$$\Omega_{\mathcal{B}}^n(F(X)) = F(\Omega_{\mathcal{A}}^n(X)). \quad \square$$

The following definition is cited from [11].

Definition 2.6 *A recollement, denoted by $(\mathcal{A}, \mathcal{B}, \mathcal{C})$, of abelian categories is a diagram*

$$\begin{array}{ccccc} \longleftarrow i^* \longleftarrow & & \longleftarrow j_! \longleftarrow & & \\ \mathcal{A} \xrightarrow{i_*} & \mathcal{B} & \xrightarrow{j^*} & \mathcal{C} & \\ \longleftarrow i^! \longleftarrow & & \longleftarrow j_* \longleftarrow & & \end{array}$$

- (1) (i^*, i_*) , $(i_*, i^!)$, $(j_!, j^*)$ and (j^*, j_*) are adjoint pairs.
- (2) i_* , $j_!$ and j_* are fully faithful.
- (3) $\text{Im } i_* = \text{Ker } j^*$.

In the rest of this section, we assume that $(\mathcal{A}, \mathcal{B}, \mathcal{C})$ is a recollement of abelian categories as in Definition 2.6. We list some properties of such recollements (see [11, 16, 18–23] and [33]), which will be used in the sequel.

Lemma 2.7 *We have*

- (1) $i^*j_! = 0 = i^!j_*$.
- (2) The functors i_* , j^* are exact, and the functors i^* , $j_!$ are right exact, and the functors $i^!$, j_* are left exact.
- (3) All natural transformations

$$i^*i_* \longrightarrow 1_{\mathcal{A}}, \quad 1_{\mathcal{A}} \longrightarrow i^!i_*, \quad 1_{\mathcal{C}} \longrightarrow j^*j_! \quad \text{and} \quad j^*j_* \longrightarrow 1_{\mathcal{C}}$$

are natural isomorphisms. Moreover, all functors i^* , $i^!$ and j^* are dense.

- (4) For any object $X \in \mathcal{B}$, if i^* is exact, then there is the following exact sequence

$$0 \longrightarrow j_!j^*(X) \longrightarrow X \longrightarrow i_*i^*(X) \longrightarrow 0$$

in \mathcal{B} ; and if $i^!$ is exact, then there is the following exact sequence

$$0 \longrightarrow i_*i^!(X) \longrightarrow X \longrightarrow j_*j^*(X) \longrightarrow 0$$

in \mathcal{B} .

(5) If i^* is exact, then $i^!j_! = 0$ and $j_!$ is exact; and if $i^!$ is exact, then $i^*j_* = 0$ and j_* is exact.

(6) The functors i^* and $j_!$ preserve projective objects. If $i^!$ is exact, then i_* and j^* preserve projective objects.

We get the following observation.

Lemma 2.8 We have the following assertions.

(1) If $i^!$ is exact, then there is an exact sequence of natural transformations

$$0 \longrightarrow i_*i^!j_! \longrightarrow j_! \longrightarrow j_* \longrightarrow 0 .$$

(2) If i^* is exact, then there is an exact sequence of natural transformations

$$0 \longrightarrow j_! \longrightarrow j_* \longrightarrow i_*i^*j_* \longrightarrow 0 .$$

Proof (1) By [11, Proposition 4.4], there is an exact sequence of natural transformations

$$0 \longrightarrow i_*i^!j_! \longrightarrow j_! \longrightarrow j_* \longrightarrow i_*i^*j_* \longrightarrow 0 .$$

Since $i^!$ is exact, we have $i^*j_* = 0$ by Lemma 2.7, and thus the assertion follows.

(2) It is a dual of (1). □

As a consequence of Lemmas 2.7 and 2.8, we get the following

Remark 2.9 Let

$$0 \longrightarrow C'' \longrightarrow C' \longrightarrow C \longrightarrow 0$$

be an exact sequence in \mathcal{C} . Assume that $i^!$ is exact. By Lemma 2.7, we have that j_* is exact. So

$$0 \longrightarrow j_*(C'') \longrightarrow j_*(C') \longrightarrow j_*(C) \longrightarrow 0$$

is exact in \mathcal{B} . By Lemma 2.8, we have the following exact sequence

$$0 \longrightarrow i_*i^!j_!(C') \longrightarrow j_!(C') \longrightarrow j_*(C') \longrightarrow 0$$

in \mathcal{B} . One can get the following pullback diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & i_*i^!j_!(C') = i_*i^!j_!(C') & & & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \dashrightarrow & K & \dashrightarrow & j_!(C') & \dashrightarrow & j_*(C) \dashrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & j_*(C'') & \longrightarrow & j_*(C') & \longrightarrow & j_*(C) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

The following result generalizes [8, Lemma 4.2], which is useful in the sequel.

Lemma 2.10 For any $n \geq 1$, assume that $i^!$ is exact and $i^!j_!$ preserves projective objects. Then, for any $X \in \mathcal{B}$, there exists an exact sequence

$$0 \longrightarrow i_*i^!j_!(P_{n-1}) \longrightarrow \Omega_{\mathcal{B}}^n(j_*j^*(X)) \longrightarrow j_*(\Omega_{\mathcal{C}}^n(j^*(X))) \longrightarrow 0$$

in \mathcal{B} , where P_{n-1} , a projective object in \mathcal{C} , lies in the exact sequence

$$0 \longrightarrow \Omega_{\mathcal{C}}^n(j^*(X)) \longrightarrow P_{n-1} \longrightarrow \Omega_{\mathcal{C}}^{n-1}(j^*(X)) \longrightarrow 0.$$

Proof Notice that $j^*(X) \in \mathcal{C}$. Consider the following exact sequence

$$0 \longrightarrow \Omega_{\mathcal{C}}^1(j^*(X)) \longrightarrow P_0 \longrightarrow j^*(X) \longrightarrow 0$$

in \mathcal{C} with P_0 a projective object. By Remark 2.9, we get the following pullback diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & \downarrow & & \downarrow & & \\ & & i_*i^!j_!(P_0) = & = & i_*i^!j_!(P_0) & & \\ & & \downarrow & & \downarrow & & \\ 0 & \dashrightarrow & K_1 & \dashrightarrow & j_!(P_0) & \dashrightarrow & j_*j^*(X) \dashrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & j_*(\Omega_{\mathcal{C}}^1(j^*(X))) & \longrightarrow & j_*(P_0) & \longrightarrow & j_*j^*(X) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

Since $j_!$ preserves projective objects by Lemma 2.7, $j_!(P_0)$ is a projective object in \mathcal{B} . So $\Omega_{\mathcal{B}}^1(j_*j^*(X)) = K_1$ and the assertion for $n = 1$ follows.

Now applying Remark 2.9 to the exact sequence

$$0 \longrightarrow \Omega_{\mathcal{C}}^2(j^*(X)) \longrightarrow P_1 \longrightarrow \Omega_{\mathcal{C}}^1(j^*(X)) \longrightarrow 0$$

in \mathcal{C} with P_1 projective yields the following pullback diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & \downarrow & & \downarrow & & \\ & & i_*i^!j_!(P_1) = & = & i_*i^!j_!(P_1) & & \\ & & \downarrow & & \downarrow & & \\ 0 & \dashrightarrow & K_2 & \dashrightarrow & j_!(P_1) & \dashrightarrow & j_*(\Omega_{\mathcal{C}}^1(j^*(X))) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & j_*(\Omega_{\mathcal{C}}^2(j^*(X))) & \longrightarrow & j_*(P_1) & \longrightarrow & j_*(\Omega_{\mathcal{C}}^1(j^*(X))) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

We also get the following pullback diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & K_2 & = & K_2 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \twoheadrightarrow & i_* i^! j_!(P_0) & \twoheadrightarrow & i_* i^! j_!(P_0) \oplus j_!(P_1) & \dashrightarrow & j_!(P_1) \dashrightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 & & \parallel & & K_1 & \longrightarrow & j_*(\Omega_{\mathcal{C}}^1(j^*(X))) \longrightarrow 0 \\
 0 & \longrightarrow & i_* i^! j_!(P_0) & \longrightarrow & K_1 & \longrightarrow & j_*(\Omega_{\mathcal{C}}^1(j^*(X))) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0.
 \end{array}$$

Notice that i_* and $i^!j_!$ preserve projective objects by Lemma 2.7 and assumption, so $\Omega_{\mathcal{B}}^2(j_*j^*(X)) = K_2$. Repeating this process, we can get the desired exact sequence

$$0 \longrightarrow i_* i^! j_!(P_{n-1}) \longrightarrow \Omega_{\mathcal{B}}^n(j_*j^*(X)) \longrightarrow j_*(\Omega^n(j^*(X))) \longrightarrow 0$$

in \mathcal{B} , where P_{n-1} , a projective object, lies in the exact sequence

$$0 \longrightarrow \Omega_{\mathcal{C}}^n(j^*(X)) \longrightarrow P_{n-1} \longrightarrow \Omega_{\mathcal{C}}^{n-1}(j^*(X)) \longrightarrow 0. \quad \square$$

3 Main Results

Let Λ be an Artin algebra, and \mathcal{D} a subcategory of $\text{mod } \Lambda$.

(1) \mathcal{D} is said to be of *finite representation type*, if there is some $N \in \text{mod } \Lambda$ such that $\text{add } \mathcal{D} = \text{add } N$; that is, the number of non-isomorphic indecomposable Λ -modules appeared in \mathcal{D} is finite. In particular, if $\mathcal{D} = \text{mod } \Lambda$, it is said that Λ is of *finite representation type* (see [2]).

(2) \mathcal{D} is said to be *m-syzygy finite* if the subcategory $\Omega^m(\mathcal{D}) := \Omega_{\text{mod } \Lambda}^m(\mathcal{D})$ is of finite representation type. In particular, if $\mathcal{D} = \text{mod } \Lambda$, it is said that Λ is an *m-syzygy finite algebra* (see [27]).

Remark 3.1 A subcategory $\mathcal{D} \subseteq \text{mod } \Lambda$ is *n-syzygy finite* if and only if $\dim \Omega^n(\mathcal{D}) = 0$. In particular, Λ is of finite representation type if and only if $\dim \text{mod } \Lambda = 0$ ([33, Corollary 3.8]).

Definition 3.2 ([28, Definition 3.1]) *Let \mathcal{D} be a subcategory of $\text{mod } \Lambda$. Then \mathcal{D} is said to be (n-)Igusa–Todorov provided that there exist $V \in \text{mod } \Lambda$ and $n \geq 0$, such that for any $M \in \Omega^n(\mathcal{D})$, there is an exact sequence*

$$0 \longrightarrow V_1 \longrightarrow V_0 \longrightarrow M \longrightarrow 0$$

in $\text{mod } \Lambda$ with $V_1, V_0 \in \text{add } V$. The module V is then called an (n-)D-Igusa–Todorov module.

In particular, if $\mathcal{D} = \text{mod } \Lambda$, it is said that Λ is an (n-)Igusa–Todorov algebra and the module V is then called an (n-)Igusa–Todorov module (see [27] and [14, Lemma 3.6]).

The following result generalizes [33, Theorem 3.14], which gives an equivalent characterization of (n-)Igusa–Todorov subcategories and means that $\dim \Omega^n(\mathcal{D})$ is an invariant for measuring how far a subcategory of $\text{mod } \Lambda$ is from being Igusa–Todorov.

Proposition 3.3 *Let \mathcal{D} be a subcategory of $\text{mod } \Lambda$. Then for any $n \geq 0$, the following statements are equivalent.*

- (1) \mathcal{D} is n -Igusa–Todorov.
- (2) $\dim \Omega^n(\mathcal{D}) \leq 1$.

Proof (1) \Rightarrow (2) Assume that \mathcal{D} is n -Igusa–Todorov. Let $X \in \Omega^n(\mathcal{D})$. Then there exists $V \in \text{mod } \Lambda$ such that the following sequence

$$0 \longrightarrow V_1 \longrightarrow V_0 \longrightarrow X \longrightarrow 0$$

in $\text{mod } \Lambda$ with $V_1, V_0 \in \text{add } V$ is exact. By Lemma 2.4, we have

$$X \in \langle V \oplus \Omega^{-1}(V) \rangle_2.$$

Thus $\dim \Omega^n(\mathcal{D}) \leq 1$.

(2) \Rightarrow (1) Assume that $\dim \Omega^n(\mathcal{D}) \leq 1$ and $X \in \mathcal{D}$. Then there exists $V \in \text{mod } \Lambda$ such that the following sequence

$$0 \longrightarrow V_1 \longrightarrow \Omega^n(X) \longrightarrow V_2 \longrightarrow 0,$$

in $\text{mod } \Lambda$ with $V_1, V_2 \in \langle V \rangle_1$ is exact. By [33, Lemma 3.2], we obtain the following exact sequence

$$0 \longrightarrow \Omega^1(V_2) \longrightarrow V_1 \oplus P \longrightarrow \Omega^n(X) \longrightarrow 0$$

in $\text{mod } \Lambda$ with P projective. Notice that both $\Omega^1(V_2)$ and $V_1 \oplus P$ are in $\text{add}(\Omega^1(V) \oplus V \oplus \Lambda)$, so \mathcal{D} is n -Igusa–Todorov. □

As an immediate consequence, we get the following

Corollary 3.4 ([28, Proposition 3.4]) *Let \mathcal{D} be a subcategory of $\text{mod } \Lambda$. If \mathcal{D} is n -Igusa–Todorov, then $\Omega^i(\mathcal{D})$ is also n -Igusa–Todorov for any $i \geq 1$. In particular, \mathcal{D} is m -Igusa–Todorov for $m \geq n$ in that case.*

Proof It follows from Lemma 2.3 and Proposition 3.3. □

Remark 3.5 ([27, Remark 2.4]) *If Λ is an n -Igusa–Todorov algebra, then Λ is also an m -Igusa–Todorov algebra for any $m \geq n$.*

The following result generalizes [33, Proposition 3.15].

Proposition 3.6 *Let \mathcal{D} be a subcategory of $\text{mod } \Lambda$ and $m, n \geq 0$. If $\dim \Omega^n(\mathcal{D}) \leq m$, then $\dim \Omega^{n-i}(\mathcal{D}) \leq m + i$ for any $0 \leq i \leq n$. In particular, if \mathcal{D} is n -Igusa–Todorov, then $\dim \mathcal{D} \leq n + 1$.*

Proof Let $X \in \mathcal{D}$. If $n = 0$ (also $i = 0$), then $\dim \mathcal{D} \leq m$. Now suppose $n \geq 1$. Consider the following exact sequence

$$\begin{array}{cccccccccccc}
0 & \longrightarrow & \Omega^n(X) & \longrightarrow & P_{n-1} & \longrightarrow & \cdots & \longrightarrow & P_{n-i} & \longrightarrow & P_{n-i-1} & \cdots & \longrightarrow & P_0 & \longrightarrow & X & \longrightarrow & 0 \\
& & & & & & & & & & & & & & & & & & \\
& & & & & & & & & \searrow & \nearrow & & & & & & & & \\
& & & & & & & & & & \Omega^{n-i}(X) & & & & & & & &
\end{array}$$

in $\text{mod } \Lambda$ with all P_i projective. Using Lemma 2.4 repeatedly, we have $\Omega^{n-i}(X) \in \langle T \rangle_{m+i+1}$ for some $T \in \text{mod } \Lambda$. Thus $\dim \Omega^{n-i}(\mathcal{D}) \leq m + i$. The last assertion follows from Proposition 3.3. □

We have the following

Remark 3.7 Let \mathcal{D} be a subcategory of $\text{mod } \Lambda$.

(1) If $\dim \mathcal{D} = n$, then $\dim \Omega^i(\mathcal{D}) \geq n - i$ for any $0 \leq i \leq n$ by Proposition 3.6. In particular, if $\dim \text{mod } \Lambda = n$, then $\dim \Omega^i(\text{mod } \Lambda) \geq n - i$ for any $0 \leq i \leq n$.

(2) If \mathcal{D} is n -syzygy finite, then $\dim \Omega^{n-1}(\mathcal{D}) \leq 1$, and so \mathcal{D} is $(n - 1)$ -Igusa–Todorov by Proposition 3.3. In addition, we have $\dim \mathcal{D} \leq n$ by Proposition 3.6. In particular, if Λ is n -syzygy finite, then $\dim \Omega^{n-1}(\text{mod } \Lambda) \leq 1$, it follows from Proposition 3.3 that Λ is an $(n - 1)$ -Igusa–Todorov algebra ([27, Proposition 2.5]). In addition, we have $\dim \text{mod } \Lambda \leq n$.

From now on, assume that Λ' , Λ and Λ'' are Artin algebras and

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j_!} & \\ \text{mod } \Lambda' & \xrightarrow{i_*} & \text{mod } \Lambda & \xrightarrow{j^*} & \text{mod } \Lambda'' \\ & \xleftarrow{i^!} & & \xleftarrow{j_*} & \end{array}$$

is a recollement.

Theorem 3.8 Let \mathcal{D}' and \mathcal{D}'' be subcategories of $\text{mod } \Lambda'$ and $\text{mod } \Lambda''$ respectively. Assume that $i^!$ is exact. For any $n, m \geq 0$, if one of the following conditions holds:

- (1) $m = 0$,
- (2) $m \geq 1$ and $i^!j_!$ preserves projective modules,

then

$$\dim \Omega^k(\mathcal{D}) \leq \dim \Omega^n(\mathcal{D}') + \dim \Omega^m(\mathcal{D}'') + 1,$$

where $k = \max\{m, n\}$ and $\mathcal{D} = \{D \in \text{mod } \Lambda \mid i^!(D) \in \mathcal{D}' \text{ and } j^*(D) \in \mathcal{D}''\}$.

Proof Let $X \in \mathcal{D}$. Then $i^!(X) \in \mathcal{D}'$ and $j^*(X) \in \mathcal{D}''$. Suppose $\dim \Omega^n(\mathcal{D}') = p$ and $\dim \Omega^m(\mathcal{D}'') = q$. Then there exist $V' \in \text{mod } \Lambda'$ and $V'' \in \text{mod } \Lambda''$ such that $\Omega^n(i^!(X)) \in \langle V' \rangle_{p+1}$ and $\Omega^m(j^*(X)) \in \langle V'' \rangle_{q+1}$.

Since $i^!$ is exact, there exists the following exact sequence

$$0 \longrightarrow i_*i^!(X) \longrightarrow X \longrightarrow j_*j^*(X) \longrightarrow 0$$

in $\text{mod } \Lambda$ by Lemma 2.7. Set $k := \max\{m, n\}$. By the horseshoe lemma, there exists an exact sequence

$$0 \longrightarrow \Omega^k(i_*i^!(X)) \longrightarrow \Omega^k(X) \longrightarrow \Omega^k(j_*j^*(X)) \longrightarrow 0 \tag{3.1}$$

in $\text{mod } \Lambda$.

- (1) If $m = 0$, then $j^*(X) \in \langle V'' \rangle_{q+1}$. Since j_* is exact by Lemma 2.7, we have

$$j_*j^*(X) \in j_*\langle V'' \rangle_{q+1} \subseteq \langle j_*(V'') \rangle_{q+1}$$

by [33, Lemma 2.4]. It follows from Lemma 2.2 that $\Omega^k(j_*j^*(X)) \in \langle \Omega^k(j_*(V'')) \rangle_{q+1}$.

Note that $\Omega^k(i^!(X)) \in \langle \widetilde{V}' \rangle_{p+1}$ for some $\widetilde{V}' \in \text{mod } \Lambda'$ by Lemma 2.3. Since i_* is exact and preserves projective modules by Lemma 2.7, we have

$$\begin{aligned} \Omega^k(i_*i^!(X)) &= i_*(\Omega^k(i^!(X))) && \text{(by Lemma 2.5)} \\ &\in i_*(\langle \widetilde{V}' \rangle_{p+1}) \subseteq \langle i_*(\widetilde{V}') \rangle_{p+1}. && \text{(by [33, Lemma 2.4])} \end{aligned}$$

Following the exact sequence (3.1), we have

$$\Omega^k(X) \in \langle i_*(\widetilde{V}') \rangle_{p+1} \diamond \langle \Omega^k(j_*(V'')) \rangle_{q+1} \subseteq \langle i_*(\widetilde{V}') \oplus \Omega^k(j_*(V'')) \rangle_{p+q+2}$$

by [33, Proposition 2.2(1) and Corollary 2.3(1)]. Thus $\dim \Omega^k(\mathcal{D}) \leq p + q + 1$.

(2) Now let $m \geq 1$. Suppose $p \geq 1$ and $q \geq 0$. By Lemma 2.3, we have $\dim \Omega^k(\mathcal{D}') \leq p$ and $\dim \Omega^k(\mathcal{D}'') \leq q$. Without loss of generality, we still assume $\Omega^k(i^!(X)) \in \langle V' \rangle_{p+1}$ and $\Omega^k(j^*(X)) \in \langle V'' \rangle_{q+1}$ for some $V' \in \text{mod } \Lambda'$ and $V'' \in \text{mod } \Lambda''$. Consider the following exact sequence

$$0 \longrightarrow V'_1 \longrightarrow \Omega^k(i^!(X)) \longrightarrow V'_2 \longrightarrow 0$$

in $\text{mod } \Lambda'$ with $V'_1 \in \text{add } V'$ and $V'_2 \in \langle V' \rangle_p$. By [31, Lemma 3.3], there is the following exact sequence

$$0 \longrightarrow \Omega^1(V'_2) \longrightarrow P' \oplus V'_1 \longrightarrow \Omega^k(i^!(X)) \longrightarrow 0$$

in $\text{mod } \Lambda'$ with P' projective. Applying the exact functor i_* to it yields the following exact sequence

$$0 \longrightarrow i_*(\Omega^1(V'_2)) \longrightarrow i_*(P' \oplus V'_1) \longrightarrow i_*(\Omega^k(i^!(X))) \longrightarrow 0,$$

where $\Omega^1(V'_2) \in \langle \Omega^1(V') \rangle_p$ by Lemma 2.2. Then by [33, Lemma 2.4], we have

$$i_*(\Omega^1(V'_2)) \in i_*(\langle \Omega^1(V') \rangle_p) \subseteq \langle i_*(\Omega^1(V')) \rangle_p.$$

Since i_* is exact and preserves projective modules by Lemma 2.7, we have that $i_*(P')$ is projective in $\text{mod } \Lambda$ and $\Omega^k(i_*i^!(X)) = i_*(\Omega^k(i^!(X)))$ by Lemma 2.5.

By Lemma 2.10, there exists the following exact sequence

$$0 \longrightarrow i_*i^!j_!(P_{k-1}) \longrightarrow \Omega^k(j_*j^*(X)) \longrightarrow j_*(\Omega^k(j^*(X))) \longrightarrow 0$$

in $\text{mod } \Lambda$ with P_{k-1} projective. By [31, Lemma 3.3], there exists the following exact sequence

$$0 \longrightarrow \Omega^1(j_*(\Omega^k(j^*(X)))) \longrightarrow P \oplus i_*i^!j_!(P_{k-1}) \longrightarrow \Omega^k(j_*j^*(X)) \longrightarrow 0$$

in $\text{mod } \Lambda$ with P projective. Since j_* is exact by Lemma 2.7, we have

$$j_*(\Omega^k(j^*(X))) \in j_*(\langle V'' \rangle_{q+1}) \subseteq \langle j_*(V'') \rangle_{q+1}$$

by [33, Lemma 2.4]. It follows from Lemma 2.2 that

$$\Omega^1(j_*(\Omega^k(j^*(X)))) \in \langle \Omega^1(j_*(V'')) \rangle_{q+1}.$$

Notice that i_* and $i^!j_!$ preserve projective modules by Lemma 2.7 and assumption, so $i_*i^!j_!(P_{k-1})$ is a projective Λ -module. Consider the following commutative diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 & & (3.2) \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & i_*(\Omega^1(V'_2)) & \dashrightarrow & V'_1 & \dashrightarrow & \Omega^1(j_*(\Omega^k(j^*(X)))) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & i_*(P' \oplus V'_1) & \longrightarrow & i_*(P' \oplus V'_1) \oplus P \oplus i_*i^!j_!(P_{k-1}) & \longrightarrow & P \oplus i_*i^!j_!(P_{k-1}) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & \Omega^k(i_*i^!(X)) & \longrightarrow & \Omega^k(X) & \longrightarrow & \Omega^k(j_*j^*(X)) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & &
 \end{array}$$

Then

$$\begin{aligned} V_1 &\in \langle i_*(\Omega(V')) \rangle_p \diamond \langle \Omega(j_*(V'')) \rangle_{q+1} \\ &\subseteq \langle i_*(\Omega(V')) \oplus \Omega(j_*(V'')) \rangle_{p+q+1} \end{aligned}$$

by [33, Proposition 2.2(1) and Corollary 2.3(1)]. Applying Lemma 2.4 to the middle column in the diagram (3.2) yields

$$\Omega^k(X) \in \langle V \rangle_{p+q+2},$$

where $V := i_*(V') \oplus \Lambda \oplus \Omega^{-1}(i_*(\Omega(V')) \oplus \Omega(j_*(V'')))$. Thus

$$\dim \Omega^k(\mathcal{D}) \leq p + q + 1.$$

If $p = 0$, then $V'_2 = 0$, and the desired assertion also follows. □

Combining with Theorem 3.8 with Proposition 3.3, we have

Corollary 3.9 *Let \mathcal{D}' and \mathcal{D}'' be subcategories of $\text{mod } \Lambda'$ and $\text{mod } \Lambda''$ respectively. Assume that $i^!$ is exact and $i^!j_!$ preserves projective modules. If \mathcal{D}' is n -Igusa–Todorov and \mathcal{D}'' is m -Igusa–Todorov, then*

$$\dim \Omega^k(\mathcal{D}) \leq 3,$$

where $k = \max\{m, n\}$ and $\mathcal{D} := \{D \in \text{mod } \Lambda \mid i^!(D) \in \mathcal{D}' \text{ and } j^*(D) \in \mathcal{D}''\}$.

The following result provides a sufficient condition for a subcategory $\mathcal{D} \subseteq \text{mod } \Lambda$ being n -Igusa–Todorov.

Corollary 3.10 *Let \mathcal{D}' and \mathcal{D}'' be subcategories of $\text{mod } \Lambda'$ and $\text{mod } \Lambda''$ respectively. Assume that $i^!$ is exact and $i^!j_!$ preserves projective modules. If \mathcal{D}' is n -syzygy finite and \mathcal{D}'' is m -syzygy finite, then \mathcal{D} is a k -Igusa–Todorov subcategory, where $k = \max\{m, n\}$ and $\mathcal{D} = \{D \in \text{mod } \Lambda \mid i^!(D) \in \mathcal{D}' \text{ and } j^*(D) \in \mathcal{D}''\}$.*

Proof It follows from Theorem 3.8, Remark 3.1 and Proposition 3.3. □

Taking $\mathcal{D}' = \text{mod } \Lambda'$ and $\mathcal{D}'' = \text{mod } \Lambda''$ in Theorem 3.8, it is easy to check that $\mathcal{D} = \{D \in \text{mod } \Lambda \mid i^!(D) \in \mathcal{D}' \text{ and } j^*(D) \in \mathcal{D}''\} = \text{mod } \Lambda$. Then we have

Remark 3.11 Assume that $i^!$ is exact and $n, m \geq 0$. Set $k := \max\{m, n\}$. If one of the following conditions holds:

- (1) $m = 0$,
- (2) $m \geq 1$ and $i^!j_!$ preserves projective objects,

then

$$\dim \Omega^k(\text{mod } \Lambda) \leq \dim \Omega^n(\text{mod } \Lambda') + \dim \Omega^m(\text{mod } \Lambda'') + 1.$$

In particular, we have

- (a) (see [33, Theorem 5.5]) If $n = 0 = m$, then $\dim \text{mod } \Lambda \leq \dim \text{mod } \Lambda' + \dim \text{mod } \Lambda'' + 1$.
- (b) If Λ' is n -Igusa–Todorov and Λ'' is m -Igusa–Todorov, then $\dim \Omega^k(\text{mod } \Lambda) \leq 3$.
- (c) If Λ' is n -syzygy finite and Λ'' is m -syzygy finite, then Λ is k -Igusa–Todorov.

The following result shows that the converse inequality in Theorem 3.8 holds true under certain conditions.

Theorem 3.12 *Let \mathcal{D} be a subcategory of $\text{mod } \Lambda$ with $i_*i^!(\mathcal{D}) \subseteq \mathcal{D}$ and $j_*j^*(\mathcal{D}) \subseteq \mathcal{D}$. If $i^!$ is exact, then*

$$\max\{\dim \Omega^n(i^!(\mathcal{D})), \dim \Omega^n(j^*(\mathcal{D}))\} \leq \dim \Omega^n(\mathcal{D})$$

for some $n \geq 0$.

Proof Suppose $\dim \Omega^n(\mathcal{D}) = p$. Then for any $X \in \mathcal{D}$, there exists $V \in \text{mod } \Lambda$ such that $\Omega^n(X) \in \langle V \rangle_{p+1}$.

Let $X' \in i^!(\mathcal{D})$. Consider the following exact sequence

$$0 \longrightarrow \Omega^n(X') \longrightarrow P'_{n-1} \longrightarrow P'_{n-2} \longrightarrow \cdots \longrightarrow P'_0 \longrightarrow X' \longrightarrow 0$$

in $\text{mod } \Lambda'$ with all P'_i projective. Since i_* is exact and preserves projective modules by Lemma 2.7, we have $\Omega^n(i_*(X')) = i_*(\Omega^n(X'))$ by Lemma 2.5. Notice that $i_*(X') \in \mathcal{D}$, so $i_*(\Omega^n(X')) \in \langle V \rangle_{p+1}$ by assumption. Since $i^!$ is exact again by assumption, we have

$$\Omega^n(X') \cong i^!i_*(\Omega^n(X')) \in i^!(\langle V \rangle_{p+1}) \subseteq \langle i^!(V) \rangle_{p+1}$$

by [33, Lemma 2.4]. Thus $\dim \Omega^n(i^!(\mathcal{D})) \leq p$.

Let $X'' \in j^*(\mathcal{D})$. Notice that $j_*(X'') \in \mathcal{D}$, so $\Omega^n(j_*(X'')) \in \langle V \rangle_{p+1}$ by assumption. Since j^* is exact and preserves projective modules by Lemma 2.7, we have

$$\begin{aligned} \Omega^n(X'') &\cong \Omega^n(j_*j^*(X'')) = j^*(\Omega^n(j_*(X''))) && \text{(by Lemma 2.5)} \\ &\in j^*(\langle V \rangle_{p+1}) \subseteq \langle j^*(V) \rangle_{p+1}. && \text{(by [33, Lemma 2.4])} \end{aligned}$$

Thus $\dim \Omega^n(j^*(\mathcal{D})) \leq p$. □

By Theorem 3.12, Proposition 3.3 and Remark 3.1, we have

Corollary 3.13 *Let \mathcal{D} be a subcategory of $\text{mod } \Lambda$ with $i_*i^!(\mathcal{D}) \subseteq \mathcal{D}$ and $j_*j^*(\mathcal{D}) \subseteq \mathcal{D}$ and $n \geq 0$. Assume that $i^!$ is exact. Then we have*

- (1) *If \mathcal{D} is n -Igusa–Todorov, then both $i^!(\mathcal{D})$ and $j^*(\mathcal{D})$ are n -Igusa–Todorov.*
- (2) *If \mathcal{D} is n -syzygy finite, then both $i^!(\mathcal{D})$ and $j^*(\mathcal{D})$ are n -syzygy finite.*

Taking $\mathcal{D} = \text{mod } \Lambda$ in Theorem 3.12, it is clear that $i_*i^!(\mathcal{D}) \subseteq \mathcal{D}$ and $j_*j^*(\mathcal{D}) \subseteq \mathcal{D}$ and that $i^!(\mathcal{D}) = \text{mod } \Lambda'$ and $j^*(\mathcal{D}) = \text{mod } \Lambda''$. Then we have

Remark 3.14 If $i^!$ is exact, then

$$\max\{\dim \Omega^n(\text{mod } \Lambda'), \dim \Omega^n(\text{mod } \Lambda'')\} \leq \dim \Omega^n(\text{mod } \Lambda)$$

for some $n \geq 0$.

In particular, we have

- (1) (see [33, Theorem 5.5]) If $n = 0$, then $\max\{\dim \text{mod } \Lambda', \dim \text{mod } \Lambda''\} \leq \dim \text{mod } \Lambda$.
- (2) If Λ is n -Igusa–Todorov, then both Λ' and Λ'' are n -Igusa–Todorov.
- (3) If Λ is n -syzygy finite, then both Λ' and Λ'' are n -syzygy finite.

Let $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ be a triangular matrix Artin algebra. By [21, Example 2.12], we know that

$$\text{mod } \Lambda' \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \text{mod } \Lambda \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j_*} \\ \xleftarrow{j^*} \end{array} \text{mod } \Lambda''$$

is a recollement of module categories, where

$$i^* \left(\begin{pmatrix} X \\ Y \end{pmatrix}_f \right) = \text{Coker } f, \quad i_*(X) = \begin{pmatrix} X \\ 0 \end{pmatrix}, \quad i^! \left(\begin{pmatrix} X \\ Y \end{pmatrix}_f \right) = X,$$

$$j_!(Y) = \begin{pmatrix} M \otimes_{\Lambda''} Y \\ Y \end{pmatrix}_1, \quad j^* \left(\begin{pmatrix} X \\ Y \end{pmatrix}_f \right) = Y, \quad j_*(Y) = \begin{pmatrix} 0 \\ Y \end{pmatrix}.$$

By [15, Lemma 3.2], $i^!$ admits a right adjoint, then $i^!$ is exact. If $M_{\Lambda''}$ is projective, then $j_!$ is exact. If ${}_{\Lambda'} M$ is projective, notice that $i^! j_!(Y) = i^! \left(\begin{pmatrix} M \otimes_{\Lambda''} Y \\ Y \end{pmatrix}_1 \right) = M \otimes_{\Lambda''} Y$, then $i^! j_!$ preserves projective objects. By Remark 3.14, we get immediately the following corollary, which generalizes some results in [8].


Corollary 3.15 *Let $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$. Then we have*

- (1) (cf. [8, Theorem 4.5]) *If Λ is n -Igusa–Todorov, then both Λ' and Λ'' are n -Igusa–Todorov.*
- (2) (cf. [8, Theorem 4.3]) *If Λ is n -syzygy finite, then both Λ' and Λ'' are n -syzygy finite.*

4 Examples

In this section, all algebras are finite dimensional algebras over an algebraically closed field. For a quiver Q , we use e_i to denote the idempotent corresponding to the vertex i .

Example 4.1 Let Λ be a finite dimensional algebra given by




By [27, Example 1], Λ is a 1-Igusa–Todorov algebra.

(a) Put $e = e_1 + e_2 + e_3$, $\Lambda' := e\Lambda e$ and $\Lambda'' := (1 - e)\Lambda(1 - e)$. It follows that $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ with $M = (1 - e)\Lambda e$, where Λ' is a finite dimensional algebra given by $\bullet \longleftarrow \bullet \longleftarrow \bullet$ and Λ'' is a finite dimensional algebra given by




By Corollary 3.15, Λ' and Λ'' are 1-Igusa–Todorov algebras. In fact, Λ' (it is of finite representation type) and Λ'' (it is a matrix algebra formed by two representation-finite algebras [27, Corollary 3.3]) are 0-Igusa–Todorov algebras.

(b) Put $e := e_3$, $\Lambda' := e\Lambda e$ and $\Lambda'' := (1 - e)\Lambda(1 - e)$. It follows that $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ with $M = (1 - e)\Lambda e$, where Λ' is a finite dimensional algebra given by \bullet and Λ'' is a finite dimensional algebra given by




By Corollary 3.15, Λ' and Λ'' are 1-Igusa–Todorov algebras. In fact, Λ' (it is of finite representation type) is a 0-Igusa–Todorov algebra.

We claim that Λ'' is a 1-Igusa–Todorov algebra. Let $\widetilde{\Lambda}''$ be a finite dimensional algebra given by




Then $\widetilde{\Lambda}''$ is a trivially twisted extension (see [30] for details) of two representation-finite algebras:



By [27, Corollary 3.3], we have that $\widetilde{\Lambda}''$ is a 0-Igusa Todorov algebra. Let I be the ideal of Λ'' generated by θ . Then $\widetilde{\Lambda}''$ is the quotient algebra of Λ by I . Since $I \text{rad } \Lambda'' = 0$, it follows from [27, Theorem 3.4] that Λ'' is a 1-Igusa–Todorov algebra.

Example 4.2 Let Λ be a finite dimensional algebra given by




By [27, Example 2], Λ is a 2-Igusa–Todorov algebra. Put $e := e_1 + e_2 + e_3, \Lambda' := e\Lambda e$ and $\Lambda'' := (1 - e)\Lambda(1 - e)$. It follows that $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ with $M = (1 - e)\Lambda e$, where Λ' is a finite dimensional algebra given by $\bullet \leftarrow \bullet \leftarrow \bullet$ and Λ'' is a finite dimensional algebra given by




By Corollary 3.15, Λ' and Λ'' are 2-Igusa–Todorov algebras. In fact, Λ' (it is of finite representation type) and Λ'' (it is a matrix algebra formed by two representation-finite algebras) are 0-Igusa–Todorov algebras.

Example 4.3 Let Λ be a finite dimensional algebra given by



Put $e := e_1 + e_3$, $\Lambda' := e\Lambda e$ and $\Lambda'' := (1 - e)\Lambda(1 - e)$. It follows that $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ with $M = (1 - e)\Lambda e$, where Λ' is a finite dimensional algebra given by




and Λ'' is a finite dimensional algebra given by \bullet .

Clearly, Λ' and Λ'' are of finite representation type, and so $\dim \text{mod } \Lambda' = 0 = \dim \text{mod } \Lambda''$ by Remark 3.1. By Theorem 3.8 (or [33, Theorem 5.5]), we have $\dim \text{mod } \Lambda \leq 1$. In fact, $\dim \text{mod } \Lambda = 1$. We know that Λ is a 0-Igusa–Todorov algebra from Proposition 3.3 (or [33, Theorem 3.14]). Note that Λ is viewed as a 2-Igusa–Todorov algebra in [27, Example 3].

Example 4.4 Let Λ be a finite dimensional algebra given by




Put $e := e_3$, $\Lambda' := e\Lambda e$ and $\Lambda'' := (1 - e)\Lambda(1 - e)$. It follows that $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ with $M = (1 - e)\Lambda e$, where Λ' is a finite dimensional algebra given by \bullet and Λ'' is a finite dimensional algebra given by



Clearly, $\dim \text{mod } \Lambda' = 0$ and ${}_{\Lambda'} M$ is projective. Notice that Λ'' is a 0-Igusa–Todorov algebra (it is a matrix algebra formed by two representation-finite algebras) and is of representation infinite type, so $\dim \text{mod } \Lambda'' = 1$. Notice that ${}_{\Lambda'} M$ is projective, by Theorem 3.8 and Remark 3.14 (see [33, Theorem 5.5]), we have $1 \leq \dim \text{mod } \Lambda \leq 2$. In fact, $\dim \text{mod } \Lambda = 2$ by [32, Example 3.4].

Example 4.5 Let Λ be a finite dimensional algebra given by



with $\alpha^3 = 0$, $x_i x_j = x_j x_i$, $x_i \eta = \xi \theta$, $3 \leq i, j \leq 6$.

Put $e = e_1 + e_2$, $\Lambda' := e\Lambda e$ and $\Lambda'' := (1 - e)\Lambda(1 - e)$. It follows that $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ with $M = (1 - e)\Lambda e$, where Λ' is a finite dimensional algebra given by



and Λ'' is a finite dimensional algebra given by

$$\begin{array}{ccccccc}
 & \xrightarrow{x_3} & & \xrightarrow{x_3} & & \xrightarrow{x_3} & \\
 & \xrightarrow{x_4} & & \xrightarrow{x_4} & & \xrightarrow{x_4} & \\
 3\bullet & & 4\bullet & & 5\bullet & & 6\bullet \\
 & \xrightarrow{x_5} & & \xrightarrow{x_5} & & \xrightarrow{x_5} & \\
 & \xrightarrow{x_6} & & \xrightarrow{x_6} & & \xrightarrow{x_6} &
 \end{array}
 \quad \text{with } x_i x_j = x_j x_i, \ 3 \leq i, j \leq 6.$$

Notice that ${}_{\Lambda'} M$ is projective. As in Example 4.2, Λ' is a 0-Igusa–Todorov algebra, so $\dim \text{mod } \Lambda' \leq 1$ (in fact, $\dim \text{mod } \Lambda' = 1$). By [32, Example 3.4], we have $\dim \text{mod } \Lambda'' = 3$. By Remark 3.7, we have $\dim \Omega^1(\text{mod } \Lambda'') \geq 2$. Notice that Λ'' is 3-syzygy finite, so $\dim \Omega^1(\text{mod } \Lambda'') \leq 2$ by Proposition 3.6, and thus $\dim \Omega^1(\text{mod } \Lambda'') = 2$. Similarly, we have $\dim \Omega^1(\text{mod } \Lambda') = 1$. On the other hand, Λ' is 2-syzygy finite (Λ' is a monomial algebra), so $\dim \text{mod } \Lambda' = 0$. Then

(a) By Remarks 3.11 and 3.14 (or [33, Theorem 5.5]), $3 \leq \dim \text{mod } \Lambda \leq 5$. By Lemma 2.3 and Remark 3.7, we have $2 \leq \dim \Omega^1(\text{mod } \Lambda) \leq 5$, and $1 \leq \dim \Omega^2(\text{mod } \Lambda) \leq 5$.

(b) By Remark 3.11 and Remark 3.14, we have $2 \leq \dim \Omega^1(\text{mod } \Lambda) \leq 4$, and $1 \leq \dim \Omega^2(\text{mod } \Lambda) \leq 2$. The upper bound here is better than that in (a).

Example 4.6 Let Λ be a finite dimensional algebra given by

$$\begin{array}{ccccccc}
 & \xrightarrow{x_1} & & \xrightarrow{x_1} & & \xrightarrow{x_1} & \\
 & \xrightarrow{x_2} & & \xrightarrow{x_2} & & \xrightarrow{x_2} & \\
 1\bullet & & 2\bullet & & 3\bullet & & 4\bullet \xrightarrow{\delta} 5\bullet \\
 & \xrightarrow{x_3} & & \xrightarrow{x_3} & & \xrightarrow{x_3} & \\
 & \xrightarrow{x_4} & & \xrightarrow{x_4} & & \xrightarrow{x_4} &
 \end{array}$$

with $x_i x_j = x_j x_i$, $\delta x_i = \delta x_j$, $1 \leq i, j \leq 4$.

Put $e = e_5$, $\Lambda' := e\Lambda e$ and $\Lambda'' := (1-e)\Lambda(1-e)$. It follows that $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ with $M = (1-e)\Lambda e$, where Λ' is a finite dimensional algebra given by \bullet and Λ'' is a finite dimensional algebra given by

$$\begin{array}{ccccccc}
 & \xrightarrow{x_1} & & \xrightarrow{x_1} & & \xrightarrow{x_1} & \\
 & \xrightarrow{x_2} & & \xrightarrow{x_2} & & \xrightarrow{x_2} & \\
 1\bullet & & 2\bullet & & 3\bullet & & 4\bullet \\
 & \xrightarrow{x_3} & & \xrightarrow{x_3} & & \xrightarrow{x_3} & \\
 & \xrightarrow{x_4} & & \xrightarrow{x_4} & & \xrightarrow{x_4} &
 \end{array}
 \quad \text{with } x_i x_j = x_j x_i, \ 1 \leq i, j \leq 4.$$

Notice that ${}_{\Lambda'} M$ is projective and $\dim \text{mod } \Lambda' = 0$. By [32, Example 3.4], we have $\dim \text{mod } \Lambda'' = 3$. By Example 4.5, we have $\dim \Omega^1(\text{mod } \Lambda'') = 2$. Then

(a) By Remarks 3.11 and 3.14 (or [33, Theorem 5.5]), we have $3 \leq \dim \text{mod } \Lambda \leq 4$. Then by Lemma 2.3 and Remark 3.7, we have $2 \leq \dim \Omega^1(\text{mod } \Lambda) \leq 4$.

(b) By Remarks 3.11 and 3.14, we have $2 \leq \dim \Omega^1(\text{mod } \Lambda) \leq 3$. The upper bound here is better than that in (a).

Conflict of Interest The authors declare no conflict of interest.

References

[1] Asadollahi, J., Bahiraei, P., Hafezi, R., et al.: On relative derived categories. *Comm. Algebra*, **44**(12), 5454–5477 (2016)

[2] Auslander, M.: Representation dimension of Artin algebras, Queen Mary College Mathematical Notes, University of London, 1971

[3] Auslander, M., Reiten, I., Smalø, S. O.: Representation Theory of Artin Algebras, Cambridge Studies in Advanced Mathematics, 36, Cambridge University Press, Cambridge, 1995

- [4] Ballard, M., Favero, D., Katzarkov, L.: Orlov spectra: bounds and gaps. *Invent. Math.*, **189**, 359–430 (2012)
- [5] Beligiannis, A.: Some ghost lemmas, survey for “The representation dimension of Artin algebras”, Bielefeld 2008, <http://www.mathematik.uni-bielefeld.de/~sek/2008/ghosts.pdf>
- [6] Beilinson, A. A., Bernstein J., Deligne, P.: Faisceaux pervers, Astérisque, 100, Soc. Math. France, Paris, 1982
- [7] Bondal, A. I., Van den Bergh, M.: Generators and representability of functors in commutative and non-commutative geometry. *Moscow Math. J.*, **3**, 1–36 (2003)
- [8] Bravo, D., Lanzilotta, M., Mendoza, O.: Pullback diagrams, syzygy finite classes and Igusa–Todorov algebras. *J. Pure Appl. Algebra*, **223**, 4494–4508 (2019)
- [9] Chen, H., Xi, C.: Recollements of derived categories III: Finitistic dimensions. *J. Lond. Math. Soc.*, **95**, 633–658 (2017)
- [10] Dao, H., Takahashi, R.: The radius of a subcategory of modules. *Algebra Number Theory*, **8**, 141–172 (2014)
- [11] Franjou, V., Pirashvili, T.: Comparison of abelian categories recollements. *Doc. Math.*, **9**, 41–56 (2004)
- [12] Gao, N.: Gorensteinness, homological invariants and Gorenstein derived categories. *Sci. China Math.*, **60**, 431–438 (2017)
- [13] Happel, D.: Reduction techniques for homological conjectures. *Tsukuba J. Math.*, **17**, 115–130 (1993)
- [14] Huang, Z., Sun, J.: Endomorphism algebras and Igusa–Todorov algebras. *Acta Math. Hungar.*, **140**, 60–70 (2013)
- [15] Lu, M.: Gorenstein defect categories of triangular matrix algebras. *J. Algebra*, **480**, 346–367 (2017)
- [16] Ma, X., Huang, Z.: Torsion pairs in recollements of abelian categories. *Front. Math. China*, **13**(4), 875–892 (2018)
- [17] Ma, X., Zhao T., Huang, Z.: Gorenstein algebras and recollements. *Comm. Algebra*, **47**, 3527–3538 (2019)
- [18] Ma, X., Xie, Z., Zhao, T.: Support τ -tilting modules and recollements. *Colloq. Math.*, **167**, 303–328 (2022)
- [19] Oppermann, S.: Lower bounds for Auslander’s representation dimension. *Duke Math. J.*, **148**, 211–249 (2009)
- [20] Orlov, D. O.: Remarks on generators and dimensions of triangulated categories. *Moscow Math. J.*, **9**, 143–149 (2009)
- [21] Psaroudakis, C.: Homological theory of recollements of abelian categories. *J. Algebra*, **398**, 63–110 (2014)
- [22] Psaroudakis, C., Skartsætherhagen, Ø., Solberg, Ø. Gorenstein categories, singular equivalences and finite generation of cohomology rings in recollements. *Trans. Amer. Math. Soc. Ser. B*, **1**, 45–95 (2014)
- [23] Psaroudakis, C., Vitória, J.: Recollements of module categories. *Appl. Categ. Struct.*, **22**, 579–593 (2014)
- [24] Qin, Y., Han, Y.: Reducing homological conjectures by n -recollements. *Algebr. Represent Theory*, **19**, 377–395 (2016)
- [25] Rouquier, R.: Representation dimension of exterior algebras. *Invent. Math.*, **165**, 357–367 (2006)
- [26] Rouquier, R.: Dimensions of triangulated categories. *J. K-theory*, **1**, 193–256 (2008)
- [27] Wei, J.: Finitistic dimension and Igusa–Todorov algebras. *Adv. Math.*, **222**, 2215–2226 (2009)
- [28] Wei, J.: Finitistic dimension conjecture and conditions on ideals. *Forum Math.*, **23**, 549–564 (2011)
- [29] Wiedemann, A.: On stratifications of derived module categories. *Canad. Math. Bull.*, **34**(2), 275–280 (1991)
- [30] Xi, C.: On the finitistic dimension conjecture I: Related to representation-finite algebras. *J. Pure Appl. Algebra*, **193**, 287–305 (2004)
- [31] Xu, D.: Idealized extensions of Artin algebras and finitistic dimensions. *Comm. Algebra*, **44**, 965–976 (2016)
- [32] Zheng, J., Huang, Z.: The derived and extension dimensions of abelian categories. *J. Algebra*, **606**, 243–265 (2022)
- [33] Zheng, J., Ma, X., Huang, Z.: The extension dimension of abelian categories. *Algebr. Represent. Theory*, **23**, 693–713 (2020)