

# ICE-closed subcategories, wide $\tau$ -tilting modules and recollements <sup>\*†</sup>

Weili Gu<sup>1</sup>, Zhaoyong Huang<sup>1</sup>, Xin Ma<sup>2</sup>

1. School of Mathematics, Nanjing University, Nanjing 210093, P.R. China

2. College of Science, Henan University of Engineering, Zhengzhou 451191, P.R. China

E-mail: weiligu20@126.com (Gu), huangzy@nju.edu.cn (Huang), maxin@haue.edu.cn (Ma)

## Abstract

Let  $(\mathcal{A}', \mathcal{A}, \mathcal{A}'')$  be a recollement of abelian categories. We establish a bijection between certain ICE-closed subcategories of  $\mathcal{A}$  and those of  $\mathcal{A}''$ . As an application, when  $\Lambda', \Lambda$  and  $\Lambda''$  are artin algebras such that  $(\text{mod } \Lambda', \text{mod } \Lambda, \text{mod } \Lambda'')$  is a recollement of abelian categories, we establish a bijection between certain doubly functorially finite ICE-closed subcategories of  $\text{mod } \Lambda$  and those of  $\text{mod } \Lambda''$ . Furthermore, we provide some constructions of wide  $\tau$ -tilting modules in a recollement.

## 1 Introduction

Wide subcategories and torsion classes play an important role in the representation theory of rings and algebras, and they are closely related to each other, see [10, 23, 27] and references therein. Recently, the notion of ICE-closed subcategories of an abelian category was introduced and studied by Enomoto [9], and such subcategories are closed under images, cokernels and extensions. This notion generalizes torsion classes and wide subcategories. Later on, Enomoto and Sakai [10] introduced the notion of wide  $\tau$ -tilting modules, which generalizes that of support  $\tau$ -tilting modules introduced by Adachi, Iyama and Reiten [1]. It was proved in [1] that there exists a bijection between support  $\tau$ -tilting modules and functorially finite torsion classes. As a generalization, Enomoto and Sakai established a bijection between wide  $\tau$ -tilting modules and doubly functorially finite ICE-closed subcategories [10].

Recollements of abelian and triangulated categories were introduced by Beilinson, Bernstein and Deligne [4] in connection with derived categories of sheaves on topological spaces with the idea that one triangulated category may be “glued together” from two others, which play an important role in representation theory of algebras. Gluing and reduction techniques with respect to a recollement have been investigated; for instance, for a given recollement of triangulated categories or abelian categories, Chen [6] glued cotorsion pairs; Liu, Vitória and Yang presented the construction of gluing of silting objects [16]; Ma and Huang [19] glued torsion pairs and torsion classes; Ma and Zhao [21] glued tilting modules; Zhang glued wide subcategories [30] and support  $\tau$ -tilting modules [31]. For more references, see [2, 5, 13, 14, 17, 22, 24, 25, 26, 28, 29]. On the other hand, the reduction of  $\tau$ -tilting modules provides an effective method for constructing support  $\tau$ -tilting modules [8, 12, 15, 32].

---

\*2020 Mathematics Subject Classification: 18E10, 16G10.

†Keywords: ICE-closed subcategories, wide  $\tau$ -tilting modules, support  $\tau$ -tilting modules, torsion classes, recollements.

In this paper, we use  $\text{mod } \Lambda$  to denote the category of finitely generated left  $\Lambda$ -module for an artin algebra  $\Lambda$ . Let  $\Lambda'$ ,  $\Lambda$  and  $\Lambda''$  be artin algebras such that

$$\text{mod } \Lambda' \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \text{mod } \Lambda \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \text{mod } \Lambda''$$

is a recollement of abelian categories.

The aim of this paper is to glue and reduce ICE-closed subcategories in a recollement. As applications, we provide some constructions and reductions of wide  $\tau$ -tilting modules in a recollement. Our main result is as follows.

**Theorem 1.1.** (Corollary 4.9) *Let  $(\text{mod } \Lambda', \text{mod } \Lambda, \text{mod } \Lambda'')$  be a recollement of abelian categories as above. If  $i^!$  is exact, then we have the following commutative diagram with all horizontal maps bijective, where the dashed arrows are derived from solid ones. See Section 2 for the meaning of the notations in the following diagram.*

$$\begin{array}{ccc} \{ \mathcal{C} \in \text{df-ice}_{i_*(\text{mod } \Lambda')} \Lambda \} & \xleftrightarrow{\quad} & \text{df-ice } \Lambda'' \\ \uparrow & \swarrow & \uparrow \\ \{ \mathcal{C} \in \text{f-tors}_{i_*(\text{mod } \Lambda')} \Lambda \} & \xleftrightarrow{\quad} & \text{f-tors } \Lambda'' \\ \uparrow & \swarrow & \uparrow \\ \{ M \in \text{w}\tau\text{-tilt } \Lambda \\ \text{with } i_*(\text{mod } \Lambda') \subseteq \mathcal{C}_M \} & \xleftrightarrow{\quad} & \text{w}\tau\text{-tilt } \Lambda'' \\ \uparrow & \swarrow & \uparrow \\ \{ M \in \text{s}\tau\text{-tilt } \Lambda \\ \text{with } i_*(\text{mod } \Lambda') \subseteq \text{Fac } M \} & \xleftrightarrow{\quad} & \text{s}\tau\text{-tilt } \Lambda'' \end{array}$$

This paper is organized as follows. In Section 2, we recall some notions and preliminary results. In Section 3, let  $(\mathcal{A}', \mathcal{A}, \mathcal{A}'')$  be a recollement of abelian categories. We establish a bijection between certain ICE-closed subcategories of  $\mathcal{A}$  and those of  $\mathcal{A}''$ . In particular, we get a bijection between certain torsion classes of  $\mathcal{A}$  and those of  $\mathcal{A}''$ . In Section 4, as applications of Section 3, we focus on wide  $\tau$ -tilting modules and support  $\tau$ -tilting modules over artin algebras. Let  $\Lambda', \Lambda$  and  $\Lambda''$  be artin algebras such that  $(\text{mod } \Lambda', \text{mod } \Lambda, \text{mod } \Lambda'')$  is a recollement of abelian categories. We establish a bijection between certain doubly functorially finite ICE-closed subcategories of  $\text{mod } \Lambda$  and those of  $\text{mod } \Lambda''$ . In particular, we establish a bijection between certain functorially finite torsion classes of  $\text{mod } \Lambda$  and those of  $\text{mod } \Lambda''$ . Furthermore, using the bijection between doubly functorially finite ICE-closed subcategories and wide  $\tau$ -tilting modules and the bijection between functorially finite torsion classes and support  $\tau$ -tilting modules, we provide some constructions of wide  $\tau$ -tilting modules and support  $\tau$ -tilting modules in a recollement. Finally, in Section 5, we give some examples to illustrate the obtained results.

## 2 Preliminaries

Throughout this paper, all abelian categories have enough projective and injective objects and all subcategories involved are full, additive and closed under isomorphisms.

**Definition 2.1.** Let  $\mathcal{A}$  be an abelian category and  $\mathcal{C}$  a subcategory of  $\mathcal{A}$ .

- (1)  $\mathcal{C}$  is *closed under extensions* if for any exact sequence

$$0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0$$

in  $\mathcal{A}$ ,  $L, N \in \mathcal{C}$  implies  $M \in \mathcal{C}$ .

- (2)  $\mathcal{C}$  is *closed under quotients (resp. subobjects)* in  $\mathcal{A}$  if for each object  $C \in \mathcal{C}$ , any quotient (resp. subobject) of  $C$  in  $\mathcal{A}$  belongs to  $\mathcal{C}$ .
- (3)  $\mathcal{C}$  is a *torsion class (resp. torsion-free class)* in  $\mathcal{A}$  if  $\mathcal{C}$  is closed under extensions and quotients (resp. subobjects).
- (4)  $\mathcal{C}$  is *closed under images (resp. kernels, cokernels)* if for any map  $f : C_1 \rightarrow C_2$  with  $C_1, C_2 \in \mathcal{C}$ ,  $\text{Im } f \in \mathcal{C}$  (resp.  $\text{Ker } f \in \mathcal{C}$ ,  $\text{Coker } f \in \mathcal{C}$ ).
- (5)  $\mathcal{C}$  is a *wide subcategory* of  $\mathcal{A}$  if  $\mathcal{C}$  is closed under extensions, kernels and cokernels.
- (6)  $\mathcal{C}$  is an *ICE-closed subcategory* of  $\mathcal{A}$  if  $\mathcal{C}$  is closed under images, cokernels and extensions.
- (7)  $\mathcal{C}$  is a *contravariantly finite subcategory* if for any object  $A \in \mathcal{A}$ , there exists a morphism  $C \rightarrow A$  with  $C \in \mathcal{C}$  such that  $\text{Hom}_{\mathcal{A}}(\mathcal{C}, C) \rightarrow \text{Hom}_{\mathcal{A}}(\mathcal{C}, A) \rightarrow 0$  is exact; dually, a *covariantly finite subcategory* is defined. The subcategory  $\mathcal{C}$  is *functorially finite* if it is both contravariantly finite and covariantly finite.

Note that all torsion classes and wide subcategories are ICE-closed subcategories.

Let  $\mathcal{C}$  be a subcategory of  $\mathcal{A}$ . We use the following notations:

- $\text{tors } \mathcal{A}$  denotes the set of torsion classes in  $\mathcal{A}$ .
- $\text{ice } \mathcal{A}$  denotes the set of ICE-closed subcategories of  $\mathcal{A}$ .
- $\text{tors}_{\mathcal{C}} \mathcal{A}$  denotes the set of torsion classes in  $\mathcal{A}$  containing  $\mathcal{C}$ .
- $\text{ice}_{\mathcal{C}} \mathcal{A}$  denotes the set of ICE-closed subcategories of  $\mathcal{A}$  containing  $\mathcal{C}$ .

Let  $\mathcal{C}$  be a subcategory of an abelian category  $\mathcal{A}$ . We use  $\text{add } \mathcal{C}$  to denote the subcategory of  $\mathcal{A}$  consisting of direct summands of finite direct sums of objects in  $\mathcal{C}$ .

**Definition 2.2.** Let  $\mathcal{C}$  be a subcategory of  $\mathcal{A}$  closed under extensions.

- (1) An object  $P \in \mathcal{C}$  is *Ext-projective in  $\mathcal{C}$*  if  $\text{Ext}_{\mathcal{A}}^1(P, C) = 0$  for any  $C \in \mathcal{C}$ . We denote by  $\mathcal{P}(\mathcal{C})$  the subcategory of  $\mathcal{C}$  consisting of Ext-projective objects in  $\mathcal{C}$ .
- (2) The subcategory  $\mathcal{C}$  is said to *have enough Ext-projectives* if for any  $C \in \mathcal{C}$ , there exists an exact sequence

$$0 \longrightarrow C' \longrightarrow P \longrightarrow C \longrightarrow 0$$

in  $\mathcal{A}$  such that  $P \in \mathcal{P}(\mathcal{C})$  and  $C' \in \mathcal{C}$ .

- (3) The subcategory  $\mathcal{C}$  is said to *have an Ext-progenerator*  $P$  if  $\mathcal{C}$  has enough Ext-projectives and  $\mathcal{P}(\mathcal{C}) = \text{add } P$  holds.

If  $\mathcal{C}$  has an Ext-progenerator, then we denote by  $P(\mathcal{C})$  the unique basic Ext-progenerator, or equivalently,  $P(\mathcal{C})$  is a direct sum of non-isomorphic indecomposable Ext-projective objects in  $\mathcal{C}$ .

**Definition 2.3.** ([10, Definition 4.8]) Let  $\Lambda$  be an artin algebra and let  $\mathcal{C}$  be an ICE-closed subcategory of  $\text{mod } \Lambda$ . The subcategory  $\mathcal{C}$  is *doubly functorially finite* if there exists a functorially finite wide subcategory  $\mathcal{W}$  of  $\text{mod } \Lambda$  such that  $\mathcal{C}$  is a functorially finite torsion class in  $\mathcal{W}$ .

Let  $\mathcal{C}$  be a subcategory of  $\text{mod } \Lambda$ . We use the following notations:

- $\text{df-ice } \Lambda$ : the set of doubly functorially finite ICE-closed subcategories of  $\text{mod } \Lambda$ .
- $\text{df-ice}_{\mathcal{C}} \Lambda$ : the set of doubly functorially finite ICE-closed subcategories of  $\text{mod } \Lambda$  containing  $\mathcal{C}$ .
- $\text{f-tors } \Lambda$ : the set of functorially finite torsion classes in  $\text{mod } \Lambda$ .
- $\text{f-tors}_{\mathcal{C}} \Lambda$ : the set of functorially finite torsion classes in  $\text{mod } \Lambda$  containing  $\mathcal{C}$ .

## 2.1 Recollements

**Definition 2.4.** ([11]) A recollement, denoted by  $(\mathcal{A}', \mathcal{A}, \mathcal{A}'')$ , of abelian categories is a diagram

$$\begin{array}{ccccc} \longleftarrow i^* \longrightarrow & \longleftarrow j_! \longrightarrow & & & \\ \mathcal{A}' \xrightarrow{i_*} \mathcal{A} & \xrightarrow{j^*} \mathcal{A}'' & & & \\ \longleftarrow i^! \longrightarrow & \longleftarrow j_* \longrightarrow & & & \end{array}$$

of abelian categories and additive functors such that

- (1)  $(i^*, i_*)$ ,  $(i_*, i^!)$ ,  $(j_!, j^*)$  and  $(j^*, j_*)$  are adjoint pairs.
- (2)  $i_*$ ,  $j_!$  and  $j_*$  are fully faithful.
- (3)  $\text{Im } i_* = \text{Ker } j^*$ .

We list some properties of recollements of abelian categories in [11, 19, 20, 25], which will be used in the sequel.

**Lemma 2.5.** *Let  $(\mathcal{A}', \mathcal{A}, \mathcal{A}'')$  be a recollement of abelian categories as in the Definition 2.4.*

- (1) *The functors  $i_*$  and  $j^*$  are exact, the functors  $i^*$  and  $j_!$  are right exact, and the functors  $i^!$  and  $j_*$  are left exact.*
- (2) *All natural transformations  $i^*i_* \rightarrow \text{Id}_{\mathcal{A}'}$ ,  $\text{Id}_{\mathcal{A}'} \rightarrow i^!i_*$ ,  $\text{Id}_{\mathcal{A}''} \rightarrow j^*j_!$  and  $j^*j_* \rightarrow \text{Id}_{\mathcal{A}''}$  are natural isomorphisms.*
- (3) *For any object  $A \in \mathcal{A}$ , there exist the following exact sequences*

$$0 \rightarrow i_*i^!(A) \rightarrow A \rightarrow j_*j^*(A) \rightarrow i_*(A_1) \rightarrow 0,$$

$$0 \rightarrow i_*(A_2) \rightarrow j_!j^*(A) \rightarrow A \rightarrow i_*i^*(A) \rightarrow 0$$

*in  $\mathcal{A}$  with  $A_1, A_2 \in \mathcal{A}'$ .*

(4) For any object  $A \in \mathcal{A}$ , if  $i^!$  is exact, then we have the following exact sequence

$$0 \longrightarrow i_* i^!(A) \longrightarrow A \longrightarrow j_* j^*(A) \longrightarrow 0$$

in  $\mathcal{A}$ .

(5) If  $i^!$  is exact, then there exists an exact sequence of natural transformations

$$0 \longrightarrow i_* i^! j_! \longrightarrow j_! \longrightarrow j_* \longrightarrow 0.$$

## 2.2 $\tau$ -tiltings

Let  $\Lambda$  be an artin algebra and  $M \in \text{mod } \Lambda$ . We use  $|M|$  to denote the number of pairwise nonisomorphic indecomposable direct summands of  $M$ . In particular,  $|\Lambda|$  denotes the number of pairwise nonisomorphic simple modules in  $\text{mod } \Lambda$ .

**Definition 2.6.** ([1, Definitions 0.1 and 0.3, and Proposition 2.3]) Let  $\Lambda$  be an artin algebra and  $M \in \text{mod } \Lambda$ .

- (1)  $M$  is  $\tau$ -rigid if  $\text{Hom}_\Lambda(M, \tau M) = 0$ , and  $M$  is  $\tau$ -tilting if it is  $\tau$ -rigid and  $|M| = |\Lambda|$ .
- (2) For  $P \in \text{proj } \Lambda$ , a pair  $(M, P)$  is a *support  $\tau$ -tilting pair* if  $M$  is  $\tau$ -rigid,  $\text{Hom}_\Lambda(P, M) = 0$  and  $|M| + |P| = |\Lambda|$ .
- (3)  $M$  is *support  $\tau$ -tilting* if there exists some  $P \in \text{proj } \Lambda$  such that  $(M, P)$  is a support  $\tau$ -tilting pair.

We denote by  $s\tau\text{-tilt } \Lambda$  the set of isomorphism classes of basic support  $\tau$ -tilting  $\Lambda$ -modules.

Let  $\Lambda$  be an artin algebra. A wide subcategory  $\mathcal{W}$  of  $\text{mod } \Lambda$  is functorially finite if and only if there exists an artin algebra  $\Gamma$  such that  $\mathcal{W}$  is equivalent to  $\text{mod } \Gamma$  [9, Proposition 4.12].

**Definition 2.7.** ([10, Definition 4.11]) Let  $\Lambda$  be an artin algebra.

- (1) For a functorially finite wide subcategory  $\mathcal{W}$  of  $\text{mod } \Lambda$  and  $M \in \mathcal{W}$ , fix an equivalence  $F : \mathcal{W} \simeq \text{mod } \Gamma$  for an artin algebra  $\Gamma$ . We say that  $M$  is  $\tau_{\mathcal{W}}$ -tilting if  $F(M)$  is a  $\tau$ -tilting  $\Gamma$ -module.
- (2)  $M$  is a *wide  $\tau$ -tilting module* if there exists some functorially finite wide subcategory  $\mathcal{W}$  such that  $M$  is  $\tau_{\mathcal{W}}$ -tilting.

We denote by  $w\tau\text{-tilt } \Lambda$  the set of isomorphism classes of basic wide  $\tau$ -tilting modules.

Following [10, Proposition 4.12], wide  $\tau$ -tilting modules can be regarded as a generalization of support  $\tau$ -tilting modules. Enomoto and Sakai [10, Theorem 4.13] established a bijection between wide  $\tau$ -tilting modules and doubly functorially finite ICE-closed subcategories in  $\text{mod } \Lambda$ , which generalizes the bijection between support  $\tau$ -tilting modules and functorially finite torsion classes [1, Theorem 2.7].

Let  $\Lambda$  be an artin algebra and  $M$  a  $\Lambda$ -module. Set

$$\text{cok } M := \{ A \in \text{mod } \Lambda \mid \text{there exists an exact sequence } M_1 \longrightarrow M_2 \longrightarrow A \longrightarrow 0 \text{ in } \text{mod } \Lambda \\ \text{with } M_1, M_2 \in \text{add } M \}.$$

$$\text{Fac } M := \{ A \in \text{mod } \Lambda \mid \text{there exists an exact sequence } \widetilde{M} \longrightarrow A \longrightarrow 0 \text{ in } \text{mod } \Lambda \text{ with } \widetilde{M} \in \text{add } M \}.$$

**Theorem 2.8.** ([10, Corollary 4.15]) *For an artin algebra  $\Lambda$ , there exists the following commutative diagram with all horizontal maps bijective.*

$$\begin{array}{ccc}
 \text{w}\tau\text{-tilt } \Lambda & \xrightleftharpoons[\text{cok}]{P(-)} & \text{df-ice } \Lambda \\
 \uparrow & & \uparrow \\
 \text{s}\tau\text{-tilt } \Lambda & \xrightleftharpoons[\text{Fac}]{P(-)} & \text{f-tors } \Lambda
 \end{array}$$

By this result, if  $M$  is a support  $\tau$ -tilting  $\Lambda$ -module, then  $\text{Fac } M = \text{cok } M$ . For a wide  $\tau$ -tilting module  $M$  in  $\text{mod } \Lambda$ , we use  $\mathcal{C}_M := \text{cok } M$  to denote the doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda$ .

### 3 ICE-closed subcategories and recollements

In this section, we study how to construct ICE-closed subcategories in a recollement. Assume that

$$\begin{array}{ccccc}
 & \longleftarrow i^* & & \longleftarrow j_! & \\
 \mathcal{A}' & \xrightleftharpoons[i_*]{i^!} & \mathcal{A} & \xrightleftharpoons[j_*]{j^!} & \mathcal{A}'' \\
 & \longleftarrow i^! & & \longleftarrow j_* & 
 \end{array}$$

is a recollement of abelian categories.

**Proposition 3.1.** *If  $\mathcal{C}$  is an ICE-closed subcategory of  $\mathcal{A}$  with  $i_*i^*(\mathcal{C}) \subseteq \mathcal{C}$  (resp.  $i_*i^!(\mathcal{C}) \subseteq \mathcal{C}$ ), then  $i^*(\mathcal{C})$  (resp.  $i^!(\mathcal{C})$ ) is an ICE-closed subcategory of  $\mathcal{A}'$ .*

*Proof.* We only prove that  $i^*(\mathcal{C})$  is an ICE-closed subcategory of  $\mathcal{A}'$ , the other assertion follows similarly.

Let  $f : M \rightarrow N$  be a morphism in  $\mathcal{A}'$  with  $M, N \in i^*(\mathcal{C})$ . Consider the following commutative diagram

$$\begin{array}{ccccccc}
 M & \xrightarrow{f} & N & \longrightarrow & \text{Coker } f & \longrightarrow & 0 \\
 & \searrow & \nearrow & & & & \\
 & & \text{Im } f & & & & 
 \end{array}$$

Since  $i_*$  is exact, we have the following commutative diagram

$$\begin{array}{ccccccc}
 i_*(M) & \xrightarrow{i_*(f)} & i_*(N) & \longrightarrow & i_*(\text{Coker } f) & \longrightarrow & 0 \\
 & \searrow & \nearrow & & & & \\
 & & i_*(\text{Im } f) & & & & 
 \end{array}$$

Notice that  $i_*(M), i_*(N) \in i_*i^*(\mathcal{C}) \subseteq \mathcal{C}$ , so  $i_*(\text{Coker } f), i_*(\text{Im } f) \in \mathcal{C}$  by the assumption that  $\mathcal{C}$  is closed under cokernels and images. Then  $\text{Coker } f \cong i^*i_*(\text{Coker } f) \in i^*(\mathcal{C})$  and  $\text{Im } f \cong i^*i_*(\text{Im } f) \in i^*(\mathcal{C})$ , and thus  $i^*(\mathcal{C})$  is closed under cokernels and images.

Let

$$0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$$

be an exact sequence in  $\mathcal{A}'$  with  $L, N \in i^*(\mathcal{C})$ . Applying the exact functor  $i_*$  to the above exact sequence yields the following exact sequence

$$0 \rightarrow i_*(L) \rightarrow i_*(M) \rightarrow i_*(N) \rightarrow 0$$

in  $\mathcal{A}$ . Notice that  $i_*(L), i_*(N) \in i_*i^*(\mathcal{C}) \subseteq \mathcal{C}$ , so  $i_*(M) \in \mathcal{C}$  by the assumption that  $\mathcal{C}$  is closed under extensions. Then  $M \cong i_*i^*(M) \in i^*(\mathcal{C})$ , and so  $i^*(\mathcal{C})$  is closed under extensions. Thus  $i^*(\mathcal{C})$  is ICE-closed.  $\square$

**Lemma 3.2.** *If  $\mathcal{C}$  is an ICE-closed subcategory of  $\mathcal{A}$  with  $i_*(\mathcal{A}') \subseteq \mathcal{C}$ , then it holds that*

$$(1) \ i_*i^*(\mathcal{C}) \subseteq \mathcal{C}, \ i_*i^!(\mathcal{C}) \subseteq \mathcal{C} \text{ and } j_*j^*(\mathcal{C}) \subseteq \mathcal{C}.$$

$$(2) \text{ If } j!j^*(\mathcal{C}) \subseteq \mathcal{C} \text{ and } C \in \mathcal{A} \text{ such that } j^*(C) \in j^*(\mathcal{C}), \text{ then } C \in \mathcal{C}.$$

*Proof.* (1) It is clear that  $i_*i^*(\mathcal{C}) \subseteq \mathcal{C}$  and  $i_*i^!(\mathcal{C}) \subseteq \mathcal{C}$ .

Let  $C \in \mathcal{C}$ . By Lemma 2.5, there exists an exact sequence

$$0 \longrightarrow i_*i^!(C) \longrightarrow C \longrightarrow j_*j^*(C) \longrightarrow i_*(A_1) \longrightarrow 0$$

in  $\mathcal{A}$  with  $A_1 \in \mathcal{A}'$ . Notice that  $i_*(A_1) \in \mathcal{C}$ ,  $i_*i^!(C) \in \mathcal{C}$  and  $C \in \mathcal{C}$ , so  $j_*j^*(C) \in \mathcal{C}$  by the assumption that  $\mathcal{C}$  is an ICE-closed subcategory of  $\mathcal{A}$ . Thus  $j_*j^*(\mathcal{C}) \subseteq \mathcal{C}$ .

(2) Let  $C \in \mathcal{C}$ . By Lemma 2.5, there exists an exact sequence

$$0 \longrightarrow i_*(A_2) \longrightarrow j!j^*(C) \longrightarrow C \longrightarrow i_*i^*(C) \longrightarrow 0$$

in  $\mathcal{A}$  with  $A_2 \in \mathcal{A}'$ . Notice that  $j!j^*(C) \in j!j^*(\mathcal{C}) \subseteq \mathcal{C}$ ,  $i_*(A_2) \in \mathcal{C}$  and  $i_*i^*(C) \in \mathcal{C}$ , so  $C \in \mathcal{C}$  by the assumption that  $\mathcal{C}$  is an ICE-closed subcategory of  $\mathcal{A}$ .  $\square$

If  $\mathcal{C}$  is a wide subcategory of  $\mathcal{A}$  with  $i_*(\mathcal{A}') \subseteq \mathcal{C}$ , then  $j!j^*(\mathcal{C}) \subseteq \mathcal{C}$  by [30, Lemma 3.1]. For an ICE-closed subcategory  $\mathcal{C}$ , the assertion “ $j!j^*(\mathcal{C}) \subseteq \mathcal{C}$ ” does not hold true in general (see Example 5.1(3)). However, we have the following easy observation.

**Proposition 3.3.** *Let  $\mathcal{C}$  be an ICE-closed subcategory of  $\mathcal{A}$ . If  $i^!$  is exact and  $i_*(\mathcal{A}') \subseteq \mathcal{C}$ , then  $j!j^*(\mathcal{C}) \subseteq \mathcal{C}$ .*

*Proof.* Suppose  $C \in \mathcal{C}$ . Then  $j^*(C) \in \mathcal{A}''$ . By Lemma 2.5, we have an exact sequence

$$0 \longrightarrow i_*i^!j!j^*(C) \longrightarrow j!j^*(C) \longrightarrow j_*j^*(C) \longrightarrow 0$$

in  $\mathcal{A}$ . Since  $j_*j^*(C) \in j_*j^*(\mathcal{C}) \subseteq \mathcal{C}$  by Lemma 3.2 and since  $i_*i^!j!j^*(C) \in i_*(\mathcal{A}') \subseteq \mathcal{C}$  by assumption, we have  $j!j^*(C) \in \mathcal{C}$  and  $j!j^*(\mathcal{C}) \subseteq \mathcal{C}$ .  $\square$

**Proposition 3.4.** *If  $\mathcal{C}$  is an ICE-closed subcategory of  $\mathcal{A}$  with  $i_*(\mathcal{A}') \subseteq \mathcal{C}$  and  $j!j^*(\mathcal{C}) \subseteq \mathcal{C}$ , then  $j^*(\mathcal{C})$  is an ICE-closed subcategory of  $\mathcal{A}'$ .*

*Proof.* Let  $f : M \longrightarrow N$  be a morphism in  $\mathcal{A}''$  with  $M, N \in j^*(\mathcal{C})$ . Consider the following commutative diagram

$$\begin{array}{ccccccc} j_*(M) & \xrightarrow{j_*(f)} & j_*(N) & \longrightarrow & \text{Coker } j_*(f) & \longrightarrow & 0. \\ & \searrow & \nearrow & & & & \\ & & \text{Im } j_*(f) & & & & \end{array}$$

in  $\mathcal{A}$ . By Lemma 3.2(1), we have  $j_*(M), j_*(N) \in j_*j^*(\mathcal{C}) \subseteq \mathcal{C}$ . Since  $\mathcal{C}$  is closed under cokernels and images, we have  $\text{Coker } j_*(f), \text{Im } j_*(f) \in \mathcal{C}$ . Applying the exact functor  $j^*$  to the above exact sequence yields the following commutative diagram

$$\begin{array}{ccccccc}
 j^*j_*(M) & \xrightarrow{j^*j_*(f)} & j^*j_*(N) & \longrightarrow & j^*(\text{Coker } j_*(f)) & \longrightarrow & 0. \\
 \downarrow \cong & \searrow & \downarrow \cong & & \downarrow \text{---} & & \\
 & & j^*(\text{Im } j_*(f)) & & & & \\
 & & \downarrow f^! & & & & \\
 M & \xrightarrow{f} & N & \longrightarrow & \text{Coker } f & \longrightarrow & 0. \\
 \downarrow \cong & \searrow & \downarrow \cong & & & & \\
 & & \text{Im } f & & & & 
 \end{array}$$

Then  $\text{Coker } f \cong j^*(\text{Coker } j_*(f)) \in j^*(\mathcal{C})$  and  $\text{Im } f \cong j^*(\text{Im } j_*(f)) \in j^*(\mathcal{C})$ . Thus  $j^*(\mathcal{C})$  is closed under cokernels and images.

Let

$$0 \longrightarrow L \xrightarrow{g} M \longrightarrow N \longrightarrow 0$$

be an exact sequence in  $\mathcal{A}$  with  $L, N \in j^*(\mathcal{C})$ . Since  $j_*$  is left exact, one can get the following exact sequence

$$0 \longrightarrow j_*(L) \xrightarrow{j_*(g)} j_*(M) \longrightarrow \text{Coker } j_*(g) \longrightarrow 0 \quad (3.1)$$

in  $\mathcal{A}$ . Applying the exact functor  $j^*$  to the above exact sequence yields the following commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & j^*j_*(L) & \longrightarrow & j^*j_*(M) & \longrightarrow & j^*(\text{Coker } j_*(g)) \longrightarrow 0 \\
 & & \downarrow \cong & & \downarrow \cong & & \downarrow \text{---} \\
 0 & \longrightarrow & L & \longrightarrow & M & \longrightarrow & N \longrightarrow 0.
 \end{array}$$

Notice that  $j^*(\text{Coker } j_*(g)) \cong N \in j^*(\mathcal{C})$ , so  $\text{Coker } j_*(g) \in \mathcal{C}$  by Lemma 3.2(2). Since  $j_*(L) \in j_*j^*(\mathcal{C}) \subseteq \mathcal{C}$ , it follows from the exact sequence (3.1) that  $j_*(M) \in \mathcal{C}$ . Then  $M \cong j^*j_*(M) \in j^*(\mathcal{C})$  and  $j^*(\mathcal{C})$  is closed under extensions. Thus  $j^*(\mathcal{C})$  is an ICE-closed subcategory of  $\mathcal{A}$ .  $\square$

The following result provides a construction of ICE-closed subcategories of  $\mathcal{A}$  from those of  $\mathcal{A}''$ .

**Proposition 3.5.** *If  $\mathcal{C}''$  is an ICE-closed subcategory of  $\mathcal{A}''$ , then*

$$\mathcal{C} := \{A \in \mathcal{A} \mid j^*(A) \in \mathcal{C}''\}$$

*is an ICE-closed subcategory of  $\mathcal{A}$  with  $i_*(\mathcal{A}') \subseteq \mathcal{C}$ . In particular,  $j_!(\mathcal{C}'') \subseteq \mathcal{C}$ ,  $j_*(\mathcal{C}'') \subseteq \mathcal{C}$ ,  $j^*(\mathcal{C}) = \mathcal{C}''$  and  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ .*

*Proof.* Let  $f : M \longrightarrow N$  be a morphism in  $\mathcal{A}$  with  $M, N \in \mathcal{C}$ . Consider the following commutative diagram

$$\begin{array}{ccccccc}
 M & \xrightarrow{f} & N & \longrightarrow & \text{Coker } f & \longrightarrow & 0. \\
 & \searrow & \downarrow \cong & & & & \\
 & & \text{Im } f & & & & 
 \end{array}$$

Since  $j^*$  is exact, we have the following commutative diagram

$$\begin{array}{ccccccc} j^*(M) & \xrightarrow{j^*(f)} & j^*(N) & \longrightarrow & j^*(\text{Coker } f) & \longrightarrow & 0. \\ & \searrow & \nearrow & & & & \\ & & j^*(\text{Im } f) & & & & \end{array}$$

Since  $j^*(M), j^*(N) \in \mathcal{C}''$  and  $\mathcal{C}''$  is closed under cokernels and images, we have  $j^*(\text{Coker } f), j^*(\text{Im } f) \in \mathcal{C}''$ . Then  $\text{Coker } f, \text{Im } f \in \mathcal{C}$ . Hence  $\mathcal{C}$  is closed under cokernels and images.

Let

$$0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0$$

be an exact sequence in  $\mathcal{A}$  with  $L, N \in \mathcal{C}$ . Applying the exact functor  $j^*$  to the above exact sequence yields the following exact sequence

$$0 \longrightarrow j^*(L) \longrightarrow j^*(M) \longrightarrow j^*(N) \longrightarrow 0$$

in  $\mathcal{A}$ . Notice that  $j^*(L), j^*(N) \in \mathcal{C}''$  and  $\mathcal{C}''$  is closed under extensions by assumption, so  $j^*(M) \in \mathcal{C}''$ . Then  $M \in \mathcal{C}$  and  $\mathcal{C}$  is closed under extensions. Thus  $\mathcal{C}$  is an ICE-closed subcategory of  $\mathcal{A}$ .

It is clear that  $i_*(\mathcal{A}') \subseteq \mathcal{C}$  since  $j^*i_* = 0$ . The last assertion follows from the fact that  $j^*j_! \cong \text{Id}_{\mathcal{A}'}$  and  $j^*j_* \cong \text{Id}_{\mathcal{A}''}$   $\square$

**Theorem 3.6.** *There exists a bijection*

$$\{\mathcal{C} \in \text{ice}_{i_*(\mathcal{A}')} \mathcal{A} \text{ with } j_!j^*(\mathcal{C}) \subseteq \mathcal{C}\} \xrightleftharpoons[\Psi]{\Phi} \text{ice } \mathcal{A}'' ,$$

which is given by  $\Phi(\mathcal{C}) = j^*(\mathcal{C})$  and  $\Psi(\mathcal{C}'') = \{A \in \mathcal{A} \mid j^*(A) \in \mathcal{C}''\}$  for  $\mathcal{C} \in \text{ice}_{i_*(\mathcal{A}')} \mathcal{A}$  with  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ , and  $\mathcal{C}'' \in \text{ice } \mathcal{A}''$ .

*Proof.* By Propositions 3.4 and 3.5, it suffices to prove  $\Phi\Psi = \text{Id}$  and  $\Psi\Phi = \text{Id}$ .

Let  $\mathcal{C}'' \in \text{ice } \mathcal{A}''$ . Write  $\tilde{\mathcal{C}} := \{A \in \mathcal{A} \mid j^*(A) \in \mathcal{C}''\}$ . Then

$$\Phi\Psi(\mathcal{C}'') = \Phi(\tilde{\mathcal{C}}) = j^*(\tilde{\mathcal{C}}).$$

Clearly, we have  $j^*(\tilde{\mathcal{C}}) \subseteq \mathcal{C}''$ . Since  $j_!(\mathcal{C}'') \subseteq \tilde{\mathcal{C}}$  by Proposition 3.5, we have  $\mathcal{C}'' \cong j^*j_!(\mathcal{C}'') \subseteq j^*(\tilde{\mathcal{C}})$ , and so  $j^*(\tilde{\mathcal{C}}) = \mathcal{C}''$ . Thus  $\Phi\Psi(\mathcal{C}'') = \mathcal{C}''$  and  $\Phi\Psi = \text{Id}$ .

Let  $\mathcal{C} \in \text{ice}_{i_*(\mathcal{A}')} \mathcal{A}$  with  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ . Then

$$\Psi\Phi(\mathcal{C}) = \Psi(j^*(\mathcal{C})) = \{A \in \mathcal{A} \mid j^*(A) \in j^*(\mathcal{C})\}.$$

Clearly, we have  $\mathcal{C} \subseteq \{A \in \mathcal{A} \mid j^*(A) \in j^*(\mathcal{C})\}$ . Now suppose  $C \in \{A \in \mathcal{A} \mid j^*(A) \in j^*(\mathcal{C})\}$ , that is,  $j^*(C) \in j^*(\mathcal{C})$ . Then  $C \in \mathcal{C}$  by Lemma 3.2(2), and so  $\{A \in \mathcal{A} \mid j^*(A) \in j^*(\mathcal{C})\} \subseteq \mathcal{C}$  and  $\{A \in \mathcal{A} \mid j^*(A) \in j^*(\mathcal{C})\} = \mathcal{C}$ . Thus  $\Psi\Phi(\mathcal{C}) = \mathcal{C}$  and  $\Psi\Phi = \text{Id}$ .  $\square$

In particular, restricting Propositions 3.4 and 3.5 and Theorem 3.6 to wide subcategories, which recover [30, Propositions 3.2 and 3.3 and Theorem 3.4], respectively. Restricting Propositions 3.4 and 3.5 to torsion classes, one can get the following two results.

**Proposition 3.7.** *If  $\mathcal{C}$  is a torsion class in  $\mathcal{A}$  with  $i_*(\mathcal{A}') \subseteq \mathcal{C}$  and  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ , then  $j^*(\mathcal{C})$  is a torsion class in  $\mathcal{A}''$ .*

*Proof.* By Proposition 3.4, we only need to prove that  $j^*(\mathcal{C})$  is closed under quotients. Let  $f : M \rightarrow N$  be an epimorphism in  $\mathcal{A}''$  with  $M \in j^*(\mathcal{C})$ . Since  $j_!$  is right exact, we have that  $j_!(M) \rightarrow j_!(N) \rightarrow 0$  is an exact sequence in  $\mathcal{A}$ . Since  $j_!(M) \in j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$  and  $\mathcal{C}$  is closed under quotients by assumption, we have  $j_!(N) \in \mathcal{C}$ . Then  $N \cong j^*j_!(N) \in j^*(\mathcal{C})$  and  $j^*(\mathcal{C})$  is closed under quotients.  $\square$

**Proposition 3.8.** *If  $\mathcal{C}''$  is a torsion class in  $\mathcal{A}''$ , then*

$$\mathcal{C} := \{A \in \mathcal{A} \mid j^*(A) \in \mathcal{C}''\}$$

*is a torsion class in  $\mathcal{A}$  with  $i_*(\mathcal{A}') \subseteq \mathcal{C}$ . In particular,  $j_!(\mathcal{C}'') \subseteq \mathcal{C}$ ,  $j_*(\mathcal{C}'') \subseteq \mathcal{C}$ ,  $j^*(\mathcal{C}) = \mathcal{C}''$  and  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ .*

*Proof.* By Proposition 3.5, we only need to prove that  $\mathcal{C}$  is closed under quotients. Let  $f : M \rightarrow N$  be an epimorphism in  $\mathcal{A}$  with  $M \in \mathcal{C}$ . Since  $j^*$  is exact, we have that  $j^*(M) \rightarrow j^*(N) \rightarrow 0$  is an exact sequence in  $\mathcal{A}''$ . Since  $j^*(M) \in \mathcal{C}''$  and  $\mathcal{C}''$  is closed under quotients by assumption, we have  $j^*(N) \in \mathcal{C}''$ . Then  $N \in \mathcal{C}$  and  $\mathcal{C}$  is closed under quotients.  $\square$

Restricting Theorem 3.6 to torsion classes, one can get the following result.

**Theorem 3.9.** *There exists a bijection*

$$\{\mathcal{C} \in \text{tors}_{i_*(\mathcal{A}')} \mathcal{A} \text{ with } j_!j^*(\mathcal{C}) \subseteq \mathcal{C}\} \begin{array}{c} \xrightarrow{\Phi} \\ \xleftarrow{\Psi} \end{array} \text{tors } \mathcal{A}'' ,$$

*which is given by  $\Phi(\mathcal{C}) = j^*(\mathcal{C})$  and  $\Psi(\mathcal{C}'') = \{A \in \mathcal{A} \mid j^*(A) \in \mathcal{C}''\}$  for  $\mathcal{C} \in \text{tors}_{i_*(\mathcal{A}')} \mathcal{A}$  with  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ , and  $\mathcal{C}'' \in \text{tors } \mathcal{A}''$ .*

*Proof.* It follows from Propositions 3.7 and 3.8 and Theorem 3.6.  $\square$

## 4 Wide $\tau$ -tilting modules and recollements

In this section, we establish some correspondences with respect to doubly functorially finite ICE-closed subcategories and functorially finite torsion classes, and provide some constructions of wide  $\tau$ -tilting modules and support  $\tau$ -tilting modules in a recollement.

Let  $F : \mathcal{A} \rightarrow \mathcal{B}$  be an additive functor between abelian categories  $\mathcal{A}$  and  $\mathcal{B}$ , and let  $\mathcal{C}$  and  $\mathcal{D}$  be subcategories of  $\mathcal{A}$  and  $\mathcal{B}$ , respectively. If  $F(\mathcal{C}) \subseteq \mathcal{D}$ , we denote  $\overline{F} := F|_{\mathcal{C}}$ . The following two results provide certain construction of a recollement from wide subcategories.

**Theorem 4.1.** ([30, Theorem 3.8]) *Let  $(\mathcal{A}', \mathcal{A}, \mathcal{A}'')$  be a recollement of abelian categories as in the Definition 2.4. If  $\mathcal{C}$  is a wide subcategory of  $\mathcal{A}$  with  $i_*(\mathcal{A}') \subseteq \mathcal{C}$ , then*

$$\mathcal{A}' \begin{array}{c} \xleftarrow{\overline{i_*}} \\ \xrightarrow{\overline{i_*}} \\ \xleftarrow{\overline{j_!}} \end{array} \mathcal{C} \begin{array}{c} \xleftarrow{\overline{j_!}} \\ \xrightarrow{\overline{j_*}} \\ \xleftarrow{\overline{j_*}} \end{array} j^*(\mathcal{C})$$

*is a recollement of abelian categories.*

**Corollary 4.2.** *Let  $(\mathcal{A}', \mathcal{A}, \mathcal{A}'')$  be a recollement of abelian categories as in the Definition 2.4. If  $\mathcal{C}''$  is a wide subcategory of  $\mathcal{A}''$ , then*

$$\begin{array}{ccccc} \mathcal{A}' & \begin{array}{c} \xleftarrow{\overline{i^*}} \\ \xrightarrow{i_*} \\ \xleftarrow{\overline{i^!}} \end{array} & \mathcal{C} & \begin{array}{c} \xleftarrow{\overline{j^!}} \\ \xrightarrow{j^*} \\ \xleftarrow{\overline{j^*}} \end{array} & \mathcal{C}'' \end{array}$$

*is a recollement of abelian categories, where  $\mathcal{C} := \{A \in \mathcal{A} \mid j^*(A) \in \mathcal{C}''\}$ .*

*Proof.* It follows from [30, Proposition 3.3], Proposition 3.5 and Theorem 4.1.  $\square$

Recall from [7] that a pair of subcategories  $(\mathcal{T}, \mathcal{F})$  of an abelian category  $\mathcal{A}$  is called a *torsion pair* if the following conditions are satisfied.

- (1)  $\text{Hom}_{\mathcal{A}}(\mathcal{T}, \mathcal{F}) = 0$ ; that is,  $\text{Hom}_{\mathcal{A}}(T, F) = 0$  for any  $T \in \mathcal{T}$  and  $F \in \mathcal{F}$ .
- (2) For any object  $A \in \mathcal{A}$ , there exists an exact sequence

$$0 \longrightarrow T \longrightarrow A \longrightarrow F \longrightarrow 0$$

in  $\mathcal{A}$  with  $T \in \mathcal{T}$  and  $F \in \mathcal{F}$ .

For a torsion pair  $(\mathcal{T}, \mathcal{F})$  in  $\mathcal{A}$ , we have that  $\mathcal{T}$  is a torsion class in  $\mathcal{A}$ . For a length abelian category  $\mathcal{A}$ , that is,  $\mathcal{A}$  is an abelian category in which every object has finite length, it is well known that  $\mathcal{T}$  is a torsion class in  $\mathcal{A}$  if and only if  $(\mathcal{T}, \mathcal{T}^\perp)$  is a torsion pair in  $\mathcal{A}$ , where  $\mathcal{T}^\perp := \{A \in \mathcal{A} \mid \text{Hom}_{\mathcal{A}}(T, A) = 0 \text{ for any } T \in \mathcal{T}\}$ .

**Proposition 4.3.** *Let  $\mathcal{A}'$ ,  $\mathcal{A}$  and  $\mathcal{A}''$  be length abelian categories. If  $\mathcal{C}$  is a functorially finite torsion class in  $\mathcal{A}$  with  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ , then  $j^*(\mathcal{C})$  is a functorially finite torsion class in  $\mathcal{A}''$ .*

*Proof.* See [22, Theorem 3.7(1)].  $\square$

**Proposition 4.4.** *Let  $\mathcal{A}'$ ,  $\mathcal{A}$  and  $\mathcal{A}''$  be length abelian categories, and let  $\mathcal{C}'$  and  $\mathcal{C}''$  be functorially finite torsion classes in  $\mathcal{A}'$  and  $\mathcal{A}''$ , respectively. Set  $\mathcal{C} := \{A \in \mathcal{A} \mid i^*(A) \in \mathcal{C}' \text{ and } j^*(A) \in \mathcal{C}''\}$ . If  $i^!$  is exact, then the following assertions hold.*

- (1) *If  $i_*i^!(\mathcal{C}) \subseteq \mathcal{C}$ , then  $\mathcal{C}$  is a functorially finite torsion class in  $\mathcal{A}$ .*
- (2) *If  $i_*(\mathcal{A}') \subseteq \mathcal{C}$ , then  $\mathcal{C} = \{A \in \mathcal{A} \mid j^*(A) \in \mathcal{C}''\}$  and  $\mathcal{C}$  is a functorially finite torsion class in  $\mathcal{A}$ .*

*Proof.* (1) See [22, Theorem 3.3].

(2) If  $i_*(\mathcal{A}') \subseteq \mathcal{C}$ , then  $\mathcal{A}' \cong i^*i_*(\mathcal{A}') \subseteq i^*(\mathcal{C}) \subseteq \mathcal{C}' \subseteq \mathcal{A}'$ , and so  $i^*(\mathcal{C}) = \mathcal{C}' = \mathcal{A}'$ . Thus  $i^*(\mathcal{A}) \subseteq \mathcal{A}' = \mathcal{C}'$ . It is easy to check that  $\mathcal{C} = \{A \in \mathcal{A} \mid j^*(A) \in \mathcal{C}''\}$ . On the other hand, it is obvious that  $i_*i^!(\mathcal{C}) \subseteq i_*(\mathcal{A}') \subseteq \mathcal{C}$ . Thus  $\mathcal{C}$  is a functorially finite torsion class in  $\mathcal{A}$  by (1).  $\square$

From now on, assume that  $\Lambda'$ ,  $\Lambda$  and  $\Lambda''$  are artin algebras such that

$$\text{mod } \Lambda' \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \text{mod } \Lambda \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j^*} \end{array} \text{mod } \Lambda''$$

is a recollement of abelian categories.

**Proposition 4.5.** *If  $\mathcal{C}$  is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda$  with  $i_*(\text{mod } \Lambda') \subseteq \mathcal{C}$  and  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ , then  $j^*(\mathcal{C})$  is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda''$ .*

*Proof.* By assumption, there exists a functorially finite wide subcategory  $\mathcal{W}$  of  $\text{mod } \Lambda$  such that  $\mathcal{C}$  is a functorially finite torsion class in  $\mathcal{W}$ . By [30, Proposition 3.2], we have that  $j^*(\mathcal{W})$  is a wide subcategory of  $\text{mod } \Lambda''$ . It follows from [22, Lemmas 3.5 and 3.6] that  $j^*(\mathcal{W})$  is functorially finite in  $\text{mod } \Lambda''$ . By Theorem 4.1, we have that

$$\text{mod } \Lambda' \begin{array}{c} \xleftarrow{\overline{i^*}} \\ \xrightarrow{\overline{i_*}} \\ \xleftarrow{\overline{i^!}} \end{array} \mathcal{W} \begin{array}{c} \xleftarrow{\overline{j^!}} \\ \xrightarrow{\overline{j^*}} \\ \xleftarrow{\overline{j_*}} \end{array} j^*(\mathcal{W})$$

is a recollement of abelian categories. By Proposition 4.3, we know that  $j^*(\mathcal{C})$  is a functorially finite torsion class in  $j^*(\mathcal{W})$ . Thus  $j^*(\mathcal{C})$  is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda''$ .  $\square$

**Proposition 4.6.** *Assume that  $i^!$  is exact. If  $\mathcal{C}'' \subseteq \text{mod } \Lambda''$  is a doubly functorially finite ICE-closed subcategory, then*

$$\mathcal{C} := \{A \in \text{mod } \Lambda \mid j^*(A) \in \mathcal{C}''\}$$

*is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda$  with  $i_*(\text{mod } \Lambda') \subseteq \mathcal{C}$ . In particular,  $j_!(\mathcal{C}'') \subseteq \mathcal{C}$ ,  $j_*(\mathcal{C}'') \subseteq \mathcal{C}$ ,  $j^*(\mathcal{C}) = \mathcal{C}''$  and  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ .*

*Proof.* By assumption, there exists a functorially finite wide subcategory  $\mathcal{W}''$  of  $\text{mod } \Lambda''$  such that  $\mathcal{C}''$  is a functorially finite torsion class in  $\mathcal{W}''$ . By [30, Proposition 3.3], we have that  $\mathcal{W} := \{A \in \text{mod } \Lambda \mid j^*(A) \in \mathcal{W}''\}$  is a wide subcategory of  $\text{mod } \Lambda$  and  $\mathcal{C} \subseteq \mathcal{W}$ . It follows from [31, Proposition 3.1] that  $\mathcal{W}$  is covariantly finite in  $\text{mod } \Lambda$ . Then  $\mathcal{W}$  is functorially finite in  $\text{mod } \Lambda$  by [9, Proposition 4.12], and thus

$$\text{mod } \Lambda' \begin{array}{c} \xleftarrow{\overline{i^*}} \\ \xrightarrow{\overline{i_*}} \\ \xleftarrow{\overline{i^!}} \end{array} \mathcal{W} \begin{array}{c} \xleftarrow{\overline{j^!}} \\ \xrightarrow{\overline{j^*}} \\ \xleftarrow{\overline{j_*}} \end{array} \mathcal{W}''$$

is a recollement of abelian categories by Corollary 4.2. Since  $i^!$  is exact, we have that  $\overline{i^!}$  is exact. It follows from Proposition 4.4 that  $\mathcal{C}$  is a functorially finite torsion class in  $\mathcal{W}$ . Thus  $\mathcal{C}$  is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda$ .  $\square$

**Theorem 4.7.** *If  $i^!$  is exact, then there exists a bijection*

$$\{\mathcal{C} \in \text{df-ice}_{i_*(\text{mod } \Lambda')} \Lambda\} \begin{array}{c} \xleftarrow{\Phi} \\ \xrightarrow{\Psi} \end{array} \text{df-ice } \Lambda'',$$

*which is given by  $\Phi(\mathcal{C}) = j^*(\mathcal{C})$  and  $\Psi(\mathcal{C}'') = \{A \in \text{mod } \Lambda \mid j^*(A) \in \mathcal{C}''\}$  for  $\mathcal{C} \in \text{df-ice}_{i_*(\text{mod } \Lambda')} \Lambda$  and  $\mathcal{C}'' \in \text{df-ice } \Lambda''$ .*

*Proof.* Since  $i^!$  is exact by assumption, we have that  $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$  for any  $\mathcal{C} \in \text{df-ice}_{i_*(\text{mod } \Lambda')} \Lambda$  by Proposition 3.3. The assertion follows from Propositions 4.5 and 4.6, and Theorem 3.6.  $\square$

Clearly, a functorially finite torsion class is a doubly functorially finite ICE-closed subcategory. Restricting Theorem 4.7 to functorially finite torsion classes, we get the following result.

**Theorem 4.8.** *If  $i^!$  is exact, then there exists a bijection*

$$\{\mathcal{C} \in \text{f-tors}_{i_*(\text{mod } \Lambda')} \Lambda\} \xrightleftharpoons[\Psi]{\Phi} \text{f-tors } \Lambda'',$$

which is given by  $\Phi(\mathcal{C}) = j^*(\mathcal{C})$  and  $\Psi(\mathcal{C}'') = \{A \in \text{mod } \Lambda \mid j^*(A) \in \mathcal{C}''\}$  for  $\mathcal{C} \in \text{f-tors}_{i_*(\text{mod } \Lambda')} \Lambda$  and  $\mathcal{C}'' \in \text{f-tors } \Lambda''$ .

*Proof.* It follows from Propositions 4.3 and 4.4 and Theorem 4.7.  $\square$

Combing Theorems 2.8, 4.7 and 4.8, we immediately get the following result.

**Corollary 4.9.** *If  $i^!$  is exact, then we have the following commutative diagram with all horizontal maps bijective, where the dashed arrows derive from solid arrows.*

$$\begin{array}{ccc} \{\mathcal{C} \in \text{df-ice}_{i_*(\text{mod } \Lambda')} \Lambda\} & \xrightleftharpoons{\quad} & \text{df-ice } \Lambda'' \\ \uparrow & \swarrow & \uparrow \\ \{\mathcal{C} \in \text{f-tors}_{i_*(\text{mod } \Lambda')} \Lambda\} & \xrightleftharpoons{\quad} & \text{f-tors } \Lambda'' \\ \uparrow & \swarrow & \uparrow \\ \left\{ \begin{array}{l} M \in \text{w}\tau\text{-tilt } \Lambda \\ \text{with } i_*(\text{mod } \Lambda') \subseteq \mathcal{C}_M \end{array} \right\} & \xrightleftharpoons{\quad} & \text{w}\tau\text{-tilt } \Lambda'' \\ \uparrow & \swarrow & \uparrow \\ \left\{ \begin{array}{l} M \in \text{s}\tau\text{-tilt } \Lambda \\ \text{with } i_*(\text{mod } \Lambda') \subseteq \text{Fac } M \end{array} \right\} & \xrightleftharpoons{\quad} & \text{s}\tau\text{-tilt } \Lambda'' \end{array}$$

The following result shows that the converse of [10, Lemma 4.4] holds true, which is useful in the sequel.

**Lemma 4.10.** *Let  $\mathcal{C}$  be an ICE-closed subcategory of an abelian category  $\mathcal{A}$  with  $M$  an Ext-projective object in  $\mathcal{C}$ . Then  $\mathcal{C} = \text{cok } M$  if and only if  $M$  is an Ext-progenerator of  $\mathcal{C}$ .*

*Proof.* The sufficiency follows from [10, Lemma 4.4]. In the following, we prove the necessity.

Let  $P$  be an indecomposable Ext-projective object in  $\mathcal{C}$ . By assumption, there exists an exact sequence

$$M_1 \xrightarrow{f} M_2 \rightarrow P \rightarrow 0$$

in  $\mathcal{A}$  with  $M_1, M_2 \in \text{add } M$ . Since  $\mathcal{C}$  is ICE-closed, we have  $\text{Im } f \in \mathcal{C}$ . Then the following exact sequence

$$0 \rightarrow \text{Im } f \rightarrow M_2 \rightarrow P \rightarrow 0$$

splits, and so  $P$  is a direct summand of  $M_2$ . It follows that  $P \in \text{add } M$  and  $\mathcal{P}(\mathcal{C}) \subseteq \text{add } M$ . Clearly,  $\text{add } M \subseteq \mathcal{P}(\mathcal{C})$ . Thus  $\mathcal{P}(\mathcal{C}) = \text{add } M$ . On the other hand, for any  $C \in \mathcal{C}$ , there exists an exact sequence

$$\widetilde{M}_1 \xrightarrow{g} \widetilde{M}_2 \rightarrow C \rightarrow 0$$

in  $\mathcal{A}$  with  $\widetilde{M}_1, \widetilde{M}_2 \in \text{add } M$ . Notice that  $\text{Im } g \in \mathcal{C}$  since  $\mathcal{C}$  is ICE-closed, thus  $M$  is an Ext-progenerator of  $\mathcal{C}$ .  $\square$

As an application of Proposition 4.5, we give a construction of wide  $\tau$ -tilting  $\Lambda''$ -modules from wide  $\tau$ -tilting  $\Lambda$ -modules.

**Proposition 4.11.** *Assume that  $j_*$  is exact. If  $M$  is a wide  $\tau$ -tilting  $\Lambda$ -module such that  $i_*(\text{mod } \Lambda') \subseteq \mathcal{C}_M$  and  $j_!j^*(\mathcal{C}_M) \subseteq \mathcal{C}_M$ , then  $j^*(M)$  is a wide  $\tau$ -tilting module in  $\text{mod } \Lambda''$ .*

*Proof.* By assumption, we have that  $\mathcal{C}_M$  is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda$ . It follows from Proposition 4.5 that  $j^*(\mathcal{C}_M)$  is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda''$ .

We claim that  $\text{cok } j^*(M) = j^*(\mathcal{C}_M)$ . Clearly,  $j^*(M) \in j^*(\mathcal{C}_M)$  holds. It follows that  $\text{cok } j^*(M) \subseteq j^*(\mathcal{C}_M)$  from the fact that  $j^*(\mathcal{C}_M)$  is closed under cokernels. Now suppose  $A_1 \in j^*(\mathcal{C}_M)$ . Then there exists an object  $C \in \mathcal{C}_M$  such that  $A_1 \cong j^*(C)$ . Since  $\mathcal{C}_M = \text{cok } M$ , there exists an exact sequence

$$M_1 \longrightarrow M_2 \longrightarrow C \longrightarrow 0$$

in  $\text{mod } \Lambda$  with  $M_1, M_2 \in \text{add } M$ . Since  $j^*$  is exact, we get the following exact sequence

$$j^*(M_1) \longrightarrow j^*(M_2) \longrightarrow j^*(C) (\cong A_1) \longrightarrow 0$$

in  $\text{mod } \Lambda''$  with  $j^*(M_1), j^*(M_2) \in \text{add } j^*(M)$ . Then  $A_1 \in \text{cok } j^*(M)$  and  $j^*(\mathcal{C}_M) \subseteq \text{cok } j^*(M)$ . Thus  $\text{cok } j^*(M) = j^*(\mathcal{C}_M)$ . The claim is proved.

By Theorem 2.8, we have that  $M'' := P(j^*(\mathcal{C}_M))$  is a wide  $\tau$ -tilting module with  $\text{cok } M'' = j^*(\mathcal{C}_M)$ , then  $\text{cok } M'' = j^*(\mathcal{C}_M) = \text{cok } j^*(M)$ . On the other hand, since  $j_*$  is exact by assumption and  $j_*j^*(\mathcal{C}_M) \subseteq \mathcal{C}_M$  by Lemma 3.2(1), we have

$$\text{Ext}_{\Lambda''}^1(j^*(M), j^*(\mathcal{C}_M)) \cong \text{Ext}_{\Lambda}^1(M, j_*j^*(\mathcal{C}_M)) = 0$$

by [22, Proposition 2.8]. Then  $j^*(M)$  is an Ext-projective object in  $j^*(\mathcal{C}_M)$ . So  $\text{add } M'' = \text{add } j^*(M)$  by Lemma 4.10, and hence  $j^*(M)$  is a wide  $\tau$ -tilting  $\Lambda''$ -module.  $\square$

In the following result, we provide a construction of wide  $\tau$ -tilting modules in  $\text{mod } \Lambda$  from those in  $\text{mod } \Lambda''$ .

**Proposition 4.12.** *Let  $M''$  be a wide  $\tau$ -tilting  $\Lambda''$ -module. If  $i^!$  and  $j_!$  are exact, then  $i_*(\Lambda') \oplus j_!(M'')$  is a wide  $\tau$ -tilting  $\Lambda$ -module.*

*Proof.* By assumption, we have that  $\mathcal{C}_{M''}$  is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda''$ . It follows from Proposition 4.6 that  $\mathcal{C} := \{A \in \text{mod } \Lambda \mid j^*(A) \in \mathcal{C}_{M''}\}$  is a doubly functorially finite ICE-closed subcategory of  $\text{mod } \Lambda$ . By Theorem 2.8, there exists a wide  $\tau$ -tilting module  $M := P(\mathcal{C})$  such that  $\text{cok } M = \mathcal{C}$ . Since  $j^*i_* = 0$  and  $j^*j_! \cong \text{Id}$ , we have  $i_*(\Lambda') \oplus j_!(M'') \in \mathcal{C}$ , and so  $\text{cok}(i_*(\Lambda') \oplus j_!(M'')) \subseteq \mathcal{C}$  by the fact that  $\mathcal{C}$  is closed under cokernels.

On the other hand, assume  $A \in \mathcal{C}$ . Then  $j^*(A) \in \mathcal{C}_{M''} = \text{cok } M''$  and there exists the following exact sequence

$$M_1'' \xrightarrow{f} M_2'' \longrightarrow j^*(A) \longrightarrow 0$$

in  $\text{mod } \Lambda''$  with  $M_1'', M_2'' \in \text{add } M'' \subseteq \mathcal{C}_{M''}$ . Thus  $\text{Im } f \in \mathcal{C}_{M''}$ . Since  $j_!$  is exact by assumption, we get the following exact sequence

$$0 \longrightarrow j_!(\text{Im } f) \longrightarrow j_!(M_2'') \longrightarrow j_!j^*(A) \longrightarrow 0$$

in  $\text{mod } \Lambda$ . Since  $i^!$  is exact by assumption, we further get the following exact sequence

$$0 \longrightarrow i_*i^!j_!j^*(A) \longrightarrow j_!j^*(A) \longrightarrow j_*j^*(A) \longrightarrow 0$$

in  $\text{mod } \Lambda$  by Lemma 2.5. Consider the following pull-back diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & \downarrow & & \downarrow & & \\ & & j_!(\text{Im } f) & \text{===} & j_!(\text{Im } f) & & \\ & & \downarrow & & \downarrow & & \\ 0 & \text{---} & \downarrow & & \downarrow & & \\ 0 & \text{---} & K & \text{---} & j_!(M_2'') & \text{---} & j_*j^*(A) & \text{---} & 0 \\ & & \downarrow & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & i_*i^!j_!j^*(A) & \longrightarrow & j_!j^*(A) & \longrightarrow & j_*j^*(A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \parallel & & \\ & & 0 & & 0 & & & & \end{array}$$

Note that  $i_*i^!j_!j^*(A) \in i_*(\text{mod } \Lambda') \subseteq \mathcal{C}$  and  $j_!(\text{Im } f) \in j_!(\mathcal{C}_{M''}) \subseteq \mathcal{C}$  by Proposition 4.6. One can get  $K \in \mathcal{C}$  since  $\mathcal{C}$  is closed under extensions. Because  $i^!(A) \in \text{mod } \Lambda'$ , there exists an exact sequence

$$0 \longrightarrow K' \longrightarrow P' \longrightarrow i^!(A) \longrightarrow 0$$

in  $\text{mod } \Lambda'$  with  $P' \in \text{add } \Lambda'$ . Since  $i_*$  is exact, we get the following exact sequence

$$0 \longrightarrow i_*(K') \longrightarrow i_*(P') \longrightarrow i_*i^!(A) \longrightarrow 0$$

in  $\text{mod } \Lambda$ . Since  $j^*i_* = 0$  and  $j_!$  is exact, we have

$$\text{Ext}_{\Lambda}^1(j_!(M_2''), i_*i^!(A)) \cong \text{Ext}_{\Lambda''}^1(M_2'', j_*i_*i^!(A)) = 0$$

by [22, Proposition 2.8]. Consider the following commutative diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & i_*(K') & \text{---} & A' & \text{---} & K & \text{---} & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & i_*(P') & \longrightarrow & i_*(P') \oplus j_!(M_2'') & \longrightarrow & j_!(M_2'') & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & i_*i^!(A) & \longrightarrow & A & \longrightarrow & j_*j^*(A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & 0 & & \end{array}$$

Notice that  $i_*(K') \in \mathcal{C}$  and  $K \in \mathcal{C}$ , so  $A' \in \mathcal{C}$ . Repeating this process, we get an exact sequence

$$i_*(\widetilde{P}') \oplus j_!(\widetilde{M}_2'') \longrightarrow i_*(P') \oplus j_!(M_2'') \longrightarrow A \longrightarrow 0$$

in  $\text{mod } \Lambda$  with  $i_*(\widetilde{P}') \oplus j_!(\widetilde{M}_2''), i_*(P') \oplus j_!(M_2'') \in \text{add}(i_*(\Lambda') \oplus j_!(M''))$ . So  $A \in \text{cok}(i_*(\Lambda') \oplus j_!(M''))$ , and hence  $\mathcal{C} \subseteq \text{cok}(i_*(\Lambda') \oplus j_!(M''))$ . Thus we conclude that  $\text{cok}(i_*(\Lambda') \oplus j_!(M'')) = \mathcal{C} = \text{cok } M$ .

Since  $j^*(\mathcal{C}) = \mathcal{C}_{M''}$  by Proposition 4.6, we have

$$\begin{aligned} \text{Ext}_\Lambda^1(i_*(\Lambda') \oplus j_!(M''), \mathcal{C}) &\cong \text{Ext}_\Lambda^1(i_*(\Lambda'), \mathcal{C}) \oplus \text{Ext}_\Lambda^1(j_!(M''), \mathcal{C}) \\ &\cong \text{Ext}_{\Lambda'}^1(\Lambda', i^!(\mathcal{C})) \oplus \text{Ext}_{\Lambda''}^1(M'', j^*(\mathcal{C})) \quad (\text{by [22, Proposition 2.8]}) \\ &= 0. \end{aligned}$$

So  $i_*(\Lambda') \oplus j_!(M'')$  is an Ext-projective object in  $\mathcal{C}$ . It follows from Lemma 4.10 that  $\text{add}(i_*(\Lambda') \oplus j_!(M'')) = \text{add } M$ . Thus  $i_*(\Lambda') \oplus j_!(M'')$  is a wide  $\tau$ -tilting module in  $\text{mod } \Lambda$ .  $\square$

If  $M'$  and  $M''$  are wide  $\tau$ -tilting modules in  $\text{mod } \Lambda'$  and  $\text{mod } \Lambda''$ , respectively, then  $i_*(M') \oplus j_!(M'')$  is not necessarily a wide  $\tau$ -tilting  $\Lambda$ -module in general (see Example 5.1(4)).

The following result provides a construction of support  $\tau$ -tilting modules in a recollement.

**Proposition 4.13.** *Let  $M''$  be a support  $\tau$ -tilting  $\Lambda''$ -module. If  $i^!$  and  $j_!$  are exact, then  $i_*(\Lambda') \oplus j_!(M'')$  is a support  $\tau$ -tilting  $\Lambda$ -module.*

*Proof.* Suppose that  $(\Lambda', 0)$  and  $(M'', P'')$  are support  $\tau$ -tilting pair in  $\text{mod } \Lambda'$  and  $\text{mod } \Lambda''$ , respectively. Then  $\text{Fac } \Lambda' = \text{mod } \Lambda'$  and  $\text{Fac } M''$  are functorially finite torsion classes in  $\text{mod } \Lambda'$  and  $\text{mod } \Lambda''$ , respectively. It follows from Proposition 4.4 that  $\mathcal{C} := \{A \in \text{mod } \Lambda \mid j^*(A) \in \text{Fac } M''\}$  is a functorially finite torsion class in  $\text{mod } \Lambda$ .

We claim that  $(i_*(\Lambda') \oplus j_!(M''), j_!(P''))$  is a support  $\tau$ -tilting pair in  $\text{mod } \Lambda$ . Note that  $j_!(P'')$  is a projective  $\Lambda$ -module by [22, Proposition 2.5(2)].

Step 1: Since  $j^*(\mathcal{C}) = \text{Fac } M''$  by Proposition 3.8, we have

$$\begin{aligned} \text{Ext}_\Lambda^1(i_*(\Lambda') \oplus j_!(M''), \mathcal{C}) &\cong \text{Ext}_\Lambda^1(i_*(\Lambda'), \mathcal{C}) \oplus \text{Ext}_\Lambda^1(j_!(M''), \mathcal{C}) \\ &\cong \text{Ext}_{\Lambda'}^1(\Lambda', i^!(\mathcal{C})) \oplus \text{Ext}_{\Lambda''}^1(M'', j^*(\mathcal{C})) \quad (\text{by [22, Proposition 2.8]}) \\ &= 0. \end{aligned}$$

Then  $i_*(\Lambda') \oplus j_!(M'') \in \text{add } P(\mathcal{C})$  is a  $\tau$ -rigid  $\Lambda$ -module.

Step 2: Since  $j^*i_* = 0$ , we have

$$\begin{aligned} &\text{Hom}_\Lambda(j_!(P''), i_*(\Lambda') \oplus j_!(M'')) \\ &= \text{Hom}_\Lambda(j_!(P''), i_*(\Lambda')) \oplus \text{Hom}_\Lambda(j_!(P''), j_!(M'')) \\ &\cong \text{Hom}_{\Lambda''}(P'', j^*i_*(\Lambda')) \oplus \text{Hom}_{\Lambda''}(P'', M'') \\ &= 0. \end{aligned}$$

Step 3: Since  $\text{Hom}_\Lambda(j_!(M''), i_*(\Lambda')) \cong \text{Hom}_{\Lambda''}(M'', j^*i_*(\Lambda')) = 0$ , we have that  $i_*(\Lambda')$  and  $j_!(M'')$  have no common direct summands. Thus we have

$$\begin{aligned}
& |i_*(\Lambda') \oplus j_!(M'')| + |j_!(P'')| \\
&= |i_*(\Lambda')| + |j_!(M'')| + |j_!(P'')| \\
&= |\Lambda'| + |M''| + |P''| \quad (\text{by [22, Lemma 2.12]}) \\
&= |\Lambda'| + |\Lambda''| \\
&= |\Lambda|. \quad (\text{by [22, Lemma 2.13]})
\end{aligned}$$


Thus  $(i_*(\Lambda') \oplus j_!(M''), j_!(P''))$  is a support  $\tau$ -tilting pair in  $\text{mod } \Lambda$ , and therefore  $i_*(\Lambda') \oplus j_!(M'')$  is a support  $\tau$ -tilting module in  $\text{mod } \Lambda$ .  $\square$

If  $M'$  and  $M''$  are support  $\tau$ -tilting modules in  $\text{mod } \Lambda'$  and  $\text{mod } \Lambda''$ , respectively, then  $i_*(M') \oplus j_!(M'')$  is not necessarily a support  $\tau$ -tilting  $\Lambda$ -module in general (see Example 5.1(5)).

## 5 Examples


In this section, we give some examples to illustrate the obtained results. Let  $\Lambda', \Lambda''$  be artin algebras and let  ${}_{\Lambda'}M_{\Lambda''}$  be a  $(\Lambda', \Lambda'')$ -bimodule such that  ${}_{\Lambda'}M$  and  $M_{\Lambda''}$  are finitely generated, and let  $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$  be the triangular matrix algebra. Then  $\Lambda$  is an artin algebra [3, Proposition III.2.1]. A module in  $\text{mod } \Lambda$  can be uniquely written as a triple  $\begin{pmatrix} X \\ Y \end{pmatrix}_f$  with  $X \in \text{mod } \Lambda', Y \in \text{mod } \Lambda''$  and  $f \in \text{Hom}_{\Lambda'}(M \otimes_{\Lambda''} Y, X)$  [3, p.76].

**Example 5.1.** Let  $\Lambda'$  be a finite dimensional algebra given by the quiver  $1 \longrightarrow 2$  and let  $\Lambda''$  be a finite dimensional algebra given by the quiver  $3 \xrightarrow{\alpha} 4 \xrightarrow{\beta} 5$  with the relation  $\beta\alpha = 0$ . Define a triangular matrix algebra  $\Lambda = \begin{pmatrix} \Lambda' & M \\ 0 & \Lambda'' \end{pmatrix}$ , where  $M = \Lambda'$  and the right  $\Lambda''$ -module structure on  $\Lambda'$  is induced by the unique algebra surjective homomorphism  $\Lambda'' \xrightarrow{\phi} \Lambda'$  satisfying  $\phi(e_3) = e_1, \phi(e_4) = e_2$  and  $\phi(e_5) = 0$ . Then  $\Lambda$  is a finite dimensional algebra given by the quiver



with the relations  $\gamma\alpha = \delta\epsilon$  and  $\beta\alpha = 0$ .

The Auslander–Reiten quiver of  $\Lambda$  is



Following [25, Example 2.12], we have that

$$\text{mod } \Lambda' \begin{array}{c} \longleftarrow i^* \text{ ---} \\ \xrightarrow{i_*} \\ \longleftarrow i^! \text{ ---} \end{array} \text{mod } \Lambda \begin{array}{c} \longleftarrow j^! \text{ ---} \\ \xrightarrow{j_*} \\ \longleftarrow j^* \text{ ---} \end{array} \text{mod } \Lambda''$$

is a recollement of module categories, where


$$\begin{aligned} i^*\left(\begin{pmatrix} X \\ Y \end{pmatrix}_f\right) &= \text{Coker } f, & i_*(X) &= \begin{pmatrix} X \\ 0 \end{pmatrix}, & i^!\left(\begin{pmatrix} X \\ Y \end{pmatrix}_f\right) &= X, \\ j_!(Y) &= \begin{pmatrix} M \otimes_{\Lambda''} Y \\ Y \end{pmatrix}_1, & j^*\left(\begin{pmatrix} X \\ Y \end{pmatrix}_f\right) &= Y, & j_*(Y) &= \begin{pmatrix} 0 \\ Y \end{pmatrix}. \end{aligned}$$

By [18, Lemma 3.2], we know that  $i^!$  admits a right adjoint  $\tilde{i}^!$  and  $j_*$  admits a right adjoint  $\tilde{j}_*$ , where


$$\tilde{i}^!(X) = \begin{pmatrix} X \\ \text{Hom}_{\Lambda'}(M, X) \end{pmatrix}, \quad \tilde{j}_*\left(\begin{pmatrix} X \\ Y \end{pmatrix}_f\right) = \text{Ker}(Y \rightarrow \text{Hom}_{\Lambda'}(M, X)).$$

Thus  $i^!$  and  $j_*$  are exact. Notice that  $M_{\Lambda''}$  is projective, so  $j_!$  is exact.

By [10, Examples 1.1 and 6.4], the Hasse quiver of  $\text{ice } \Lambda'$  and  $\text{ice } \Lambda''$  are as follows, where the vertices are the corresponding wide  $\tau$ -tilting modules, and the red vertices are wide  $\tau$ -tilting modules which are not support  $\tau$ -tilting modules.



Hasse(ice  $\Lambda''$ ) :



- (1) By Propositions 4.12 and 4.13, we have the following table. It shows some wide  $\tau$ -tilting modules (resp. support  $\tau$ -tilting modules) in  $\text{mod } \Lambda$  constructed from those in  $\text{mod } \Lambda''$ , where the black marks are both wide  $\tau$ -tilting and support  $\tau$ -tilting modules and the red ones are wide  $\tau$ -tilting modules but not support  $\tau$ -tilting.

$M'' \in \text{w}\tau\text{-tilt } \Lambda''$	$i_*(\Lambda') \oplus j_!(M'') \in \text{w}\tau\text{-tilt } \Lambda$
$P(3) \oplus P(4) \oplus P(5)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ P(3) \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ P(4) \end{pmatrix} \oplus \begin{pmatrix} 0 \\ P(5) \end{pmatrix}$
$P(3) \oplus P(4) \oplus S(4)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ P(3) \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ P(4) \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ S(4) \end{pmatrix}$
$P(3) \oplus S(3) \oplus P(5)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ P(3) \end{pmatrix} \oplus \begin{pmatrix} S(1) \\ S(3) \end{pmatrix} \oplus \begin{pmatrix} 0 \\ P(5) \end{pmatrix}$
$P(4) \oplus P(5)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ P(4) \end{pmatrix} \oplus \begin{pmatrix} 0 \\ P(5) \end{pmatrix}$
$P(4) \oplus S(3)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ P(4) \end{pmatrix} \oplus \begin{pmatrix} S(1) \\ S(3) \end{pmatrix}$
$P(3) \oplus S(4)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ P(3) \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ S(4) \end{pmatrix}$
$S(3) \oplus P(5)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} S(1) \\ S(3) \end{pmatrix} \oplus \begin{pmatrix} 0 \\ P(5) \end{pmatrix}$
$P(3) \oplus P(5)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ P(3) \end{pmatrix} \oplus \begin{pmatrix} 0 \\ P(5) \end{pmatrix}$
$P(3) \oplus S(3)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ P(3) \end{pmatrix} \oplus \begin{pmatrix} S(1) \\ S(3) \end{pmatrix}$
$P(4) \oplus S(4)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ P(4) \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ S(4) \end{pmatrix}$
$P(5)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} 0 \\ P(5) \end{pmatrix}$
$P(3)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ P(3) \end{pmatrix}$
$P(4)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ P(4) \end{pmatrix}$
$S(3)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} S(1) \\ S(3) \end{pmatrix}$
$S(4)$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} S(2) \\ S(4) \end{pmatrix}$
$0$	$\begin{pmatrix} S(2) \\ 0 \end{pmatrix} \oplus \begin{pmatrix} P(1) \\ 0 \end{pmatrix}$

- (2) The condition “ $i_*i^!(\mathcal{C}) \subseteq \mathcal{C}$ ” in Proposition 3.1 is necessary. Taking an ICE-closed subcategory

$\mathcal{C} = \text{add}(\binom{S(2)}{0} \oplus \binom{S(1)}{S(3)})$  of  $\text{mod } \Lambda$ , we have that  $i^!(\mathcal{C}) = \text{add}(S(1) \oplus S(2))$  is not an ICE-closed subcategory of  $\text{mod } \Lambda'$  and  $i_*i^!(\mathcal{C}) = \text{add}(\binom{S(1)}{0} \oplus \binom{S(2)}{0}) \subsetneq \mathcal{C}$ .

- (3) For an ICE-closed subcategory  $\mathcal{C}$  of  $\text{mod } \Lambda$ , the assertion “ $j_!j^*(\mathcal{C}) \subseteq \mathcal{C}$ ” does not hold true in general. In fact,  $\mathcal{C} = \text{add}(\binom{S(1)}{0} \oplus \binom{S(1)}{P(3)} \oplus \binom{0}{P(3)} \oplus \binom{S(1)}{S(3)} \oplus \binom{0}{S(3)})$  is an ICE-closed subcategory of  $\text{mod } \Lambda$  with  $j_!j^*(\mathcal{C}) = \text{add}(\binom{P(1)}{P(3)} \oplus \binom{S(1)}{S(3)}) \subsetneq \mathcal{C}$ .
- (4) If  $M'$  and  $M''$  are wide  $\tau$ -tilting modules in  $\text{mod } \Lambda'$  and  $\text{mod } \Lambda''$ , respectively, then  $i_*(M') \oplus j_!(M'')$  is not necessarily a wide  $\tau$ -tilting module in general. Take wide  $\tau$ -tilting modules  $M' = P(1)$  and  $M'' = S(3) \oplus P(4)$  in  $\text{mod } \Lambda'$  and  $\text{mod } \Lambda''$ , respectively. We have that  $i_*(M') \oplus j_!(M'') = \binom{P(1)}{0} \oplus \binom{S(2)}{P(4)} \oplus \binom{S(1)}{S(3)}$  is not a wide  $\tau$ -tilting  $\Lambda$ -module. In fact,  $\text{cok}(i_*(M') \oplus j_!(M'')) = \text{add}(\binom{P(1)}{0} \oplus \binom{S(2)}{P(4)} \oplus \binom{S(1)}{S(3)})$  is not an ICE-closed subcategory of  $\text{mod } \Lambda$ .
- (5) If  $M'$  and  $M''$  are support  $\tau$ -tilting modules in  $\text{mod } \Lambda'$  and  $\text{mod } \Lambda''$ , respectively, then  $i_*(M') \oplus j_!(M'')$  is not necessarily a support  $\tau$ -tilting module in general. Take support  $\tau$ -tilting modules  $M' = S(1)$  and  $M'' = P(3) \oplus P(4) \oplus P(5)$  in  $\text{mod } \Lambda'$  and  $\text{mod } \Lambda''$ , respectively. We have that  $M := i_*(M') \oplus j_!(M'') = \binom{S(1)}{0} \oplus \binom{P(1)}{P(3)} \oplus \binom{S(2)}{P(4)} \oplus \binom{0}{P(5)}$  is not a support  $\tau$ -tilting  $\Lambda$ -module. In fact, there does not exist a projective  $\Lambda$ -module  $P$  such that  $(M, P)$  is a support  $\tau$ -tilting pair.

**Funding** This work was supported partially by National Natural Science Foundation of China (Grant No. 12371038), Central Plains Science and Technology Innovation Youth Top-notch Talent (Grant No. 2024ZYBJRC002) and Natural Science Foundation of Henan Province (Grant No. 262300421832). The authors thank the referees for useful suggestions.

## Declarations

**Competing interests** The authors have no relevant competing interests to declare.

## References

- [1] T. Adachi, O. Iyama and I. Reiten,  *$\tau$ -tilting theory*, *Compos. Math.* **150** (2014), 415–452.
- [2] L. Angeleri Hügel, S. Koenig and Q.H. Liu, *Recollements and tilting objects*, *J. Pure Appl. Algebra*, **215**(4) (2011), 420–438.
- [3] M. Auslander, I. Reiten and S.O. Smalø, *Representation Theory of Artin Algebras*, Cambridge Stud. in Adv. Math. **36**, Cambridge Univ. Press, Cambridge, 1995.
- [4] A.A. Beilinson, J. Bernstein and P. Deligne, *Faisceaux pervers*, In *Analysis and topology on singular spaces, I (Luminy, 1981)*, Astérisque **100**, Soc. Math. France, Paris, 1982, pp.5–171.
- [5] H.X. Chen and C.C. Xi, *Recollements of derived categories III: finitistic dimensions*, *J. Lond. Math. Soc.* **95** (2017), 633–658.
- [6] J.M. Chen, *Cotorsion pairs in a recollement of triangulated categories*, *Comm. Algebra* **41** (2013), 2903–2915.
- [7] S.E. Dickson, *A torsion theory for Abelian categories*, *Trans. Amer. Math. Soc.* **121** (1966), 223–235.

- [8] F. Eisele, G. Janssens and T. Raedschelders, *A reduction theorem for  $\tau$ -rigid modules*, Math. Z. **290** (2018) 1377–1413.
- [9] H. Enomoto, *Rigid modules and ICE-closed subcategories in quiver representations*, J. Algebra **594** (2022), 364–388.
- [10] H. Enomoto and A. Sakai, *ICE-closed subcategories and wide  $\tau$ -tilting modules*, Math. Z. **300** (2022), 541–577.
- [11] V. Franjou and T. Pirashvili, *Comparison of abelian categories recollements*, Doc. Math. **9** (2004), 41–56.
- [12] H.P. Gao,  *$\tau$ -tilting modules over one-point extensions by a simple module at a source point*, J. Algebra Appl. **21** (2022), 2250122.
- [13] N. Gao, *Gorensteinness, homological invariants and Gorenstein derived categories*, Sci. China Math. **60** (2017), 431–438.
- [14] D. Happel, *Reduction techniques for homological conjectures*, Tsukuba J. Math. **17** (1993), 115–130.
- [15] G. Jasso, *Reduction of  $\tau$ -tilting modules and torsion pairs*, Int. Math. Res. Not. IMRN **16** (2015), 7190–7237.
- [16] Q.H. Liu, J. Vitória and D. Yang, *Gluing silting objects*, Nagoya Math. J. **216** (2014), 117–151.
- [17] Y. Liu and P.Y. Zhou, *Gluing  $n$ -tilting and  $n$ -cotilting subcategories*, Bull. Malays. Math. Sci. Soc. **46** (2023), Paper No. 16, 20 pp.
- [18] M. Lu, *Gorenstein defect categories of triangular matrix algebras*, J. Algebra **480** (2017), 346–367.
- [19] X. Ma and Z.Y. Huang, *Torsion pairs in recollements of abelian categories*, Front. Math. China **13**(4) (2018), 875–892.
- [20] X. Ma, Y.Y. Peng and Z.Y. Huang, *The extension dimension of subcategories and recollements of abelian categories*, Acta Math. Sin. (Engl. Ser.) **40** (2024), 1042–1058.
- [21] X. Ma and T.W. Zhao, *Recollements and tilting modules*, Comm. Algebra **48** (2020), 5163–5175.
- [22] X. Ma, Z.Z. Xie and T.W. Zhao, *Support  $\tau$ -tilting modules and recollements*, Colloq. Math. **167** (2022), 303–328.
- [23] F. Marks and J. Šťovíček, *Torsion classes, wide subcategories and localisations*, Bull. Lond. Math. Soc. **49** (2017), 405–416.
- [24] C.E. Parra and J. Vitória, *Properties of abelian categories via recollements*, J. Pure Appl. Algebra **223** (2019), 3941–3963.
- [25] C. Psaroudakis, *Homological theory of recollements of abelian categories*, J. Algebra **398** (2014), 63–110.
- [26] Y.Y. Qin and Y. Han, *Reducing homological conjectures by  $n$ -recollements*, Algebr. Represent. Theory **19** (2016), 377–395.
- [27] A. Sota and P. Calvin, *Wide subcategories and lattices of torsion classes*, Algebr. Represent. Theory **25** (2022), 1611–1629.
- [28] Y.L. Sun and Y.H. Zhang, *Gluing simple-minded collections in triangulated categories*, J. Algebra **645** (2024), 54–85.
- [29] A. Wiedemann, *On stratifications of derived module categories*, Canad. Math. Bull. **34** (1991), 275–280.
- [30] Y.Y. Zhang, *Reduction of wide subcategories and recollements*, Algebra Colloq. **30** (2023), 713–720.

- [31] Y.Y. Zhang, *Gluing support  $\tau$ -tilting modules via symmetric ladders of height 2*, *Sci. China Math.* **67** (2024), 2217–2236.
- [32] P. Suarez,  *$\tau$ -tilting modules over one-point extensions by a projective module*, *Algebr. Represent. Theory* **21** (2018), 769–786.