Coproper Coresolutions and Direct Limits

Yucheng Wang and Zhaoyong Huang*

School of Mathematics, Nanjing University, Nanjing 210093, Jiangsu Province, P.R. China E-mail: wangyucheng2358@163.com (Wang), huangzy@nju.edu.cn (Huang)

Abstract

Let R be a ring and let M be a left R-module such that $M = \lim_{i \in I} M_i$ with I a directed index set. For a class \mathscr{X} of left R-modules, we construct certain coresolution of M from strong coproper \mathscr{X} -coresolutions of all M_i . As a consequence, we get that if \mathscr{X} is coresolving and closed under direct limits, then the supremum of \mathscr{X} -injective dimensions of all left R-modules and that of all finitely presented left R-modules coincide. Some known results are obtained as corollaries. Moreover, we get some equivalent characterizations of weakly Gorenstein algebras.

1. Introduction

It is well known that the notion of direct limits is fundamental in homological theory, which plays a very important role in studying the structure and classification of modules and rings. For example, any module is a direct limit of finitely presented modules or its finitely generated submodules, and any flat module is a direct limit of finitely generated projective modules [24, 29]. These results provide useful tools for investigating the transfer of certain homological properties between infinitely generated modules and finitely generated ones. In addition, a ring R is left Noetherian if and only if any direct limit of injective left R-modules is injective [5], and a ring R is left coherent if and only if any direct limit of FP-injective left R-modules is FP-injective [32]. These are partial classical results about direct limits. Recently, many authors studied certain properties of direct limits in (relative) homological theory, see [14, 16, 19, 21, 25, 30, 31, 35] and references therein. In particular, it was proved in [19] that if R is a left Noetherian ring and M is a left R-module such that $M = \underset{\longrightarrow}{\lim} M_i$ with I a directed index set, then a (minimal) injec $i \in I$ tive coresolution of M can be constructed from those of all M_i . The aim of this paper is to generalize this result to a much more general setting and give some applications. This paper is organized as follows.

The research was partially supported by NSFC (Grant Nos. 12371038, 12171207).

²⁰²⁰ Mathematics Subject Classification: 18G25, 16E05, 16E10.

Keywords: Strong coproper coresolutions, direct limits, relative injective dimension, weakly Gorenstein algebras.

^{*} Corresponding author.

In Section 2, some terminology and preliminary results are given. Let R be a ring and let M be a left R-module such that $M = \lim_{i \in I} M_i$ with I a directed index set. In Section 3, we prove that certain coresolution of M can be constructed from injective coresolutions of all M_i ; and if \mathscr{X} is a class of left R-modules closed under direct limits, then a strong coproper \mathscr{X} -coresolution of M can be constructed from those of all M_i (Theorem 3.5). Let R be a left Noetherian ring, and let \mathscr{X} be a subclass of ModR and $M = \lim_{i \in I} M_i$ with all M_i finitely generated. As a consequence of the above result, we get that if both \mathscr{X} and the left 1-orthogonal class of \mathscr{X} are closed under direct limits and each M_i admits a strong coproper $n - \mathscr{X}'$ -coresolution, where \mathscr{X}' is the class of finitely generated submodules of modules in \mathscr{X} , then M admits a strong coproper $n - \mathscr{X}$ -coresolution (Proposition 3.6).

In Section 4, we give some applications of the results obtained in Section 3. We prove that if \mathscr{X} is a class of left *R*-modules which is coresolving and closed under direct limits, then the supremum of \mathscr{X} -injective dimensions of all left *R*-modules and that of all finitely presented left *R*-modules coincide (Theorem 4.1). Some known results are obtained as corollaries. Finally, we obtain some equivalent characterizations of weakly Gorenstein algebras (Theorem 4.8).

2. Preliminaries

In this paper, R is an arbitrary associative ring with unit. We use ModR to denote the class of left R-modules, and use modR to denote the class of finitely presented left R-modules. Let \mathscr{C} be a subclass of ModR. We write

$${}^{\perp_1}\mathscr{C} := \{ M \in \operatorname{Mod} R \mid \operatorname{Ext}_R^1(M, C) = 0 \text{ for any } C \in \mathscr{C} \},$$

$${}^{\perp}\mathscr{C} := \{ M \in \operatorname{Mod} R \mid \operatorname{Ext}_R^{\geq 1}(M, C) = 0 \text{ for any } C \in \mathscr{C} \},$$

$${}^{\mathscr{C}^{\perp_1}} := \{ M \in \operatorname{Mod} R \mid \operatorname{Ext}_R^1(C, M) = 0 \text{ for any } C \in \mathscr{C} \},$$

$${}^{\mathscr{C}^{\perp}} := \{ M \in \operatorname{Mod} R \mid \operatorname{Ext}_R^{\geq 1}(C, M) = 0 \text{ for any } C \in \mathscr{C} \}.$$

Definition 2.1. ([10, 11]) Let $\mathscr{C} \subseteq \mathscr{D}$ be two subclasses of Mod*R*. A homomorphism $f: D \to C$ in Mod*R* with $D \in \mathscr{C}$ and $C \in \mathscr{C}$ is called a \mathscr{C} -preenvelope of D if $\operatorname{Hom}_R(f, C')$ is epic for any $C' \in \mathscr{C}$. A homomorphism $f: D \to C$ in Mod*R* is called *left minimal* if any endomorphism $h: C \to C$ is an automorphism whenever f = hf. A \mathscr{C} -preenvelope $f: D \to C$ of D is called a \mathscr{C} -envelope of D if f is left minimal. A \mathscr{C} -preenvelope $f: D \to C$ of D is called *special* if f is monic and $\operatorname{Coker} f \in {}^{\perp_1}\mathscr{C}$. The subclass \mathscr{C} is said to be (pre)enveloping in \mathscr{D} if any module in \mathscr{D} admits a \mathscr{C} -(pre)envelope, and it is said to be *special preenveloping* in \mathscr{D} if any module in \mathscr{D} admits a special \mathscr{C} -preenvelope. Dually, the notions of a (*special*) \mathscr{C} -precover of D and a *special preeovering subclass* are defined.

By the Wakamatsu lemma (cf. [11, Proposition 7.2.4]), if \mathscr{C} is closed under extensions, then any monic \mathscr{C} -envelope of a left *R*-module *M* is a special \mathscr{C} -preenvelope of *M*, and any epic \mathscr{C} -cover of a left *R*-module *M* is a special \mathscr{C} -precover of *M*.

Let \mathscr{C} be a subclass of ModR. The \mathscr{C} -injective dimension \mathscr{C} -idM of M is defined as $\inf\{n \mid \text{there exists an exact sequence}\}$

$$0 \to M \to C^0 \to C^1 \to \dots \to C^n \to 0$$

in ModR with all $C^i \in \mathscr{C}$ }, and set \mathscr{C} -id $M = \infty$ if no such integer exists. Recall that \mathscr{C} is called *coresolving* if \mathscr{C} contains all injective left R-modules, and \mathscr{C} is closed under extensions and cokernels of monomorphisms. A sequence

$$\mathbb{S}:\cdots\to S_1\to S_2\to S_3\to\cdots$$

in ModR is called Hom_R($-, \mathscr{C}$)-exact if Hom_R(\mathbb{S}, C) is exact for any $C \in \mathscr{C}$; dually, the notion of Hom_R($\mathscr{C}, -$)-exact sequences is defined [11].

Definition 2.2. Let \mathscr{C} be a subclass of ModR and $n \ge 0$. A module $M \in \text{Mod}R$ is said to *admit a coproper n-\mathscr{C}-coresolution* if there exists a $\text{Hom}_R(-, \mathscr{C})$ -exact exact sequence

$$0 \to M \xrightarrow{f^0} C^0 \xrightarrow{f^1} C^1 \xrightarrow{f^2} \cdots \xrightarrow{f^n} C^n$$
(2.1)

in ModR with all C^i in \mathscr{C} }, and M is said to admit a coproper ∞ - \mathscr{C} -coresolution if M admits a coproper n- \mathscr{C} -coresolution for all $n \geq 0$.

A coproper *n*- \mathscr{C} -coresolution of M as in (2.1) is called *minimal* if all $\operatorname{Im} f^i \to C^i$ are left minimal. A coproper *n*- \mathscr{C} -coresolution of M as in (2.1) is called *strong* if all $\operatorname{Coker} f^i$ are in ${}^{\perp_1}\mathscr{C}$. If (2.1) is a strong coproper *n*- \mathscr{C} -coresolution of M, then

$$0 \to M \xrightarrow{f^0} C^0 \xrightarrow{f^1} C^1 \xrightarrow{f^2} \cdots \xrightarrow{f^n} C^n \to \operatorname{Coker} f^n \to 0$$

is called a *partial strong coproper* n-C-coresolution of M.

Dually, the notions of proper ∞ - \mathscr{C} -resolutions and (partial) strong proper n- \mathscr{C} -resolutions are defined.

It is easy to see that (2.1) is a coproper *n*- \mathscr{C} -coresolution of M if and only if each $\operatorname{Im} f^i \to C^i$ is a monic \mathscr{C} -preenvelope of $\operatorname{Im} f^i$, and that (2.1) is a strong coproper *n*- \mathscr{C} -coresolution of M if and only if each $\operatorname{Im} f^i \to C^i$ is a special \mathscr{C} -preenvelope of $\operatorname{Im} f^i$.

The following observation might be known.

Lemma 2.3. Let \mathscr{C} be an enveloping class of left *R*-modules and $n \ge 0$. If *M* admits a coproper *n*- \mathscr{C} -coresolution

$$0 \to M \xrightarrow{f^0} C^0 \to C^1 \to \dots \to C^n, \qquad (2.2)$$

then M admits a minimal coproper n-C-coresolution in the following form:

$$0 \to M \xrightarrow{f'^0} C'^0 \xrightarrow{f'^1} C'^1 \xrightarrow{f'^2} \cdots \xrightarrow{f'^n} C'^n, \qquad (2.3)$$

where C'^{i} is a direct summand of C^{i} for any $0 \leq i \leq n$.

Proof. Since \mathscr{C} is an enveloping class, we have that M admits a monic \mathscr{C} -envelope $f'^0: M \to C'^0$ by (2.2). Set $M^1 := \operatorname{Coker} f^0$ and $M'^1 := \operatorname{Coker} f'^0$. Then we get an exact commutative diagram:



By [11, Proposition 6.1.2], the middle column splits (and hence C'^0 is a direct summand of C^0). It follows that the rightmost column splits. Thus we get a \mathscr{C} -preenvelope $M'^1 \rightarrow C^1$, which is the composition $M'^1 \xrightarrow{g^1} M^1 \hookrightarrow C^1$. Similar to above, we get a monic \mathscr{C} -envelope $f'^1 : M'^1 \to C'^1$ of M'^1 such that C'^1 is a direct summand of C^1 . Continuing this procedure, the assertion follows.

We write

 $\mathcal{P}(ModR) :=$ the class of projective left *R*-modules,

 $\mathcal{P}(\mathrm{mod}R) :=$ the class of finitely generated projective left *R*-modules,

 $\mathcal{I}(ModR) :=$ the class of injective left *R*-modules.

Definition 2.4. ([11])

- (1) A module $M \in ModR$ is called *Gorenstein projective* if $M \in {}^{\perp}\mathcal{P}(ModR)$ and M admits a coproper $\infty \mathcal{P}(ModR)$ -coresolution.
- (2) A module $N \in \operatorname{Mod} R^{op}$ is called *Gorenstein injective* if $N \in \mathcal{I}(\operatorname{Mod} R^{op})^{\perp}$ and N admits a proper $\infty \mathcal{I}(\operatorname{Mod} R^{op})$ -resolution.

We write

 $\mathcal{GP}(ModR) :=$ the class of Gorenstein projective left *R*-modules,

 $\mathcal{GP}(\mathrm{mod}R) :=$ the class of finitely generated Gorenstein projective left *R*-modules,

 $\mathcal{GI}(\mathrm{Mod}R^{op}) :=$ the class of Gorenstein injective right *R*-modules,

 $\mathcal{GI}(\mathrm{mod}R^{op}) :=$ the class of finitely generated Gorenstein injective right *R*-modules.

3. Constructions of coproper coresolutions

We begin with the following lemma.

Lemma 3.1. Let κ be a limit ordinal number. Suppose that

$$\mathbb{S}_{\alpha} := 0 \to M_{\alpha} \to X^{0}_{\alpha} \to X^{1}_{\alpha} \to \cdots \xrightarrow{f^{n}_{\alpha}} X^{n}_{\alpha}$$

is an exact sequence in ModR for any $\alpha \leq \kappa$ and $\{\mathbb{S}_{\alpha}, F_{\beta\alpha} : \mathbb{S}_{\alpha} \to \mathbb{S}_{\beta} \mid \alpha \leq \beta < \kappa\}$ is a direct system of exact sequences. If there exists a chain map from $\varinjlim_{\alpha < \kappa} \mathbb{S}_{\alpha}$ to \mathbb{S}_{κ} , that is,



Diagram (3.1)

then $\{\mathbb{S}_{\alpha}, F_{\beta\alpha} : \mathbb{S}_{\alpha} \to \mathbb{S}_{\beta} \mid \alpha \leq \beta \leq \kappa\}$ is also a direct system of exact sequences.

Proof. Consider the following commutative diagram:

where $G_{\kappa\alpha}$ is the colimit map and H_{κ} is obtained by assumption. For each $\alpha < \kappa$, set $F_{\kappa\alpha} := H_{\kappa}G_{\kappa\alpha}$. It follows that $F_{\kappa\alpha} = F_{\kappa\beta}F_{\beta\alpha}$ for any $\alpha \leq \beta < \kappa$. As a consequence, $\{\mathbb{S}_{\alpha}, F_{\beta\alpha} : \mathbb{S}_{\alpha} \to \mathbb{S}_{\beta} \mid \alpha \leq \beta \leq \kappa\}$ is a direct system.

As a consequence, we obtain the following result, which plays a crucial role in proving the main result.

Lemma 3.2. Let \mathscr{X} be a subclass of ModR, and let κ be an ordinal number. Suppose that $\{M_{\alpha}, f_{\beta\alpha} : M_{\alpha} \to M_{\beta} \mid \alpha \leq \beta < \kappa\}$ is a direct system in ModR and

$$\mathbb{S}_{\alpha} := 0 \to M_{\alpha} \xrightarrow{\varphi_{\alpha}^{0}} X_{\alpha}^{0} \to X_{\alpha}^{1} \to \cdots \xrightarrow{\varphi_{\alpha}^{n}} X_{\alpha}^{n}$$

is an exact sequence in ModR with all X^i_{α} in \mathscr{X} . If one of the conditions is satisfied:

- (1) \mathbb{S}_{α} is an injective coresolution of M_{α} ,
- (2) \mathbb{S}_{α} is a strong coproper n- \mathscr{X} -coresolution of M_{α} and both \mathscr{X} and $^{\perp_1}\mathscr{X}$ are closed under direct limits,

then these exact sequences \mathbb{S}_{α} are the members of a direct system indexed by $\alpha < \kappa$ in such a way that if $\alpha \leq \beta < \kappa$, the map from the sequence indexed by α into that indexed by β with the origin map $f_{\beta\alpha} : M_{\alpha} \to M_{\beta}$. In particular, we obtain an exact sequence

$$\lim_{\alpha < \kappa} \mathbb{S}_{\alpha} : 0 \to \lim_{\alpha < \kappa} M_{\alpha} \to \lim_{\alpha < \kappa} X^{0}_{\alpha} \to \lim_{\alpha < \kappa} X^{1}_{\alpha} \to \dots \to \lim_{\alpha < \kappa} X^{n}_{\alpha} \to \lim_{\alpha < \kappa} \operatorname{Coker} \varphi^{n}_{\alpha} \to 0$$

In Case (2), the sequence $\lim_{\alpha < \kappa} \mathbb{S}_{\alpha}$ is a partial strong coproper $n - \mathscr{X}$ -coresolution of $\lim_{\alpha < \kappa} M_{\alpha}$.

Proof. We need construct a direct system $\mathbb{S} = \{\mathbb{S}_{\alpha}, F_{\beta\alpha} : \mathbb{S}_{\alpha} \to \mathbb{S}_{\beta} \mid \alpha \leq \beta < \kappa\}$ indexed by κ , where each \mathbb{S}_{α} is a coproper *n*- \mathscr{X} -coresolution of M_{α} and $F_{\beta\alpha}$ is a sequence of maps $(f_{\beta\alpha}, f^{0}_{\beta\alpha}, \cdots, f^{n}_{\beta\alpha})$ such that the following diagram

$$\begin{aligned}
 \mathbb{S}_{\alpha} : & 0 \longrightarrow M_{\alpha} \longrightarrow X^{0}_{\alpha} \longrightarrow X^{1}_{\alpha} \longrightarrow \cdots \longrightarrow X^{n}_{\alpha} \\
 \downarrow_{F_{\beta\alpha}} & \downarrow_{f_{\beta\alpha}} & \downarrow_{f_{\beta\alpha}}^{\dagger} & \downarrow_{f_{\beta\alpha}}^{\dagger} & \downarrow_{f_{\beta\alpha}}^{\dagger} \\
 \mathbb{S}_{\beta} : & 0 \longrightarrow M_{\beta} \longrightarrow X^{0}_{\beta} \longrightarrow X^{1}_{\beta} \longrightarrow \cdots \longrightarrow X^{n}_{\beta}
 \end{aligned}$$

commutes. In the following, we use transfinite induction on $\beta < \kappa$ to construct $F_{\beta\alpha}$: $\mathbb{S}_{\alpha} \to \mathbb{S}_{\beta}$ with $\alpha \leq \beta < \kappa$.

(i) For the successor case, assume that we have constructed $F_{\gamma\alpha}$ for any $\alpha \leq \gamma \leq \beta$. Since \mathbb{S}_{β} is a strong coproper \mathscr{X} -*n*-coresolution of M_{β} , there exists $f^{i}_{\beta+1,\beta}: X^{i}_{\beta} \to X^{i}_{\beta+1}$ for any $0 \leq i \leq n$, such that the following diagram

$$\begin{split} \mathbb{S}_{\beta} : & 0 \longrightarrow M_{\beta} \longrightarrow X^{0}_{\beta} \longrightarrow X^{1}_{\beta} \longrightarrow \cdots \longrightarrow X^{n}_{\beta} \\ & \downarrow^{F_{\beta+1,\beta}} & \downarrow^{f_{\beta+1,\beta}} & \downarrow^{f_{\beta+1,\beta}} & \downarrow^{f_{\beta+1,\beta}} \\ \mathbb{S}_{\beta+1} : & 0 \longrightarrow M_{\beta+1} \longrightarrow X^{0}_{\beta+1} \longrightarrow X^{1}_{\beta+1} \longrightarrow \cdots \longrightarrow X^{n}_{\beta+1}. \end{split}$$

commutes. Let $F_{\beta+1,\beta} := (f_{\beta+1,\beta}, f^0_{\beta+1,\beta}, f^1_{\beta+1,\beta}, \cdots, f^n_{\beta+1,\beta})$ and $F_{\beta+1,\alpha} := F_{\beta+1,\beta}F_{\beta\alpha}$ for any ordinal $\alpha < \beta$. Then we complete the proof for the successor case.

(ii) For the limit case, let $\beta < \kappa$ be a limit ordinal. Assume that we have constructed $F_{\gamma\alpha}$ for any $\alpha \leq \gamma < \beta$. Now we need construct $F_{\beta\alpha}$ for any $\alpha < \beta$. Note that $\{\mathbb{S}_{\alpha}, F_{\gamma\alpha} : \mathbb{S}_{\alpha} \to \mathbb{S}_{\gamma} \mid \alpha \leq \gamma < \beta\}$ is a direct subsystem of \mathbb{S} . We need to find the chain map in Diagram (3.1).

For Case (1), since X^i_{β} is injective, it is clear.

For Case (2), we get an exact sequence $\lim_{\alpha < \beta} \mathbb{S}_{\alpha}$ and a colimt map $G_{\beta\alpha} = (g_{\beta\alpha}, g^{0}_{\beta\alpha}, g^{1}_{\beta\alpha}, \cdots, g^{n}_{\beta\alpha})$. Set $K^{1}_{\alpha} := \operatorname{Coker} \varphi^{0}_{\alpha}$ for any $\alpha < \beta$. Then $\{K^{1}_{\alpha}\}_{\alpha < \beta}$ is also a direct system. Since $K_{\alpha} \in {}^{\perp_{1}} \mathscr{X}$ and ${}^{\perp_{1}} \mathscr{X}$ is closed under direct limits, we have $\lim_{\alpha < \beta} K_{\alpha} \in {}^{\perp_{1}} \mathscr{X}$ and $\lim_{\alpha < \beta} M_{\alpha} \rightarrow \lim_{\alpha < \beta} X^{0}_{\alpha}$ is a special \mathscr{X} -preenvelope of $\lim_{\alpha < \beta} M_{\alpha}$. Since $K_{\alpha}^{1} \rightarrow X_{\alpha}^{1}$ is a special \mathscr{X} -preenvelope of K_{α}^{1} , we get that $\lim_{\alpha < \beta} K_{\alpha}^{1} \rightarrow \lim_{\alpha < \beta} X_{\alpha}^{1}$ is a special \mathscr{X} -preenvelope of $\lim_{\alpha < \beta} K_{\alpha}^{1}$ by using an argument similar to that as above. Continuing this procedure, we have that $\lim_{\alpha < \beta} S_{\alpha}$ is a strong coproper *n*- \mathscr{X} -coresolution. By using the universal property of $\lim_{\alpha < \beta} M_{\alpha}$, there exists a unique h_{β} such that $h_{\beta}g_{\beta\alpha} = f_{\beta\alpha}$, and we get $H_{\beta} = (h_{\beta}, h_{\beta}^{0}, \cdots, h_{\beta}^{n})$ induced by the origin map h_{β} . Then by Lemma 3.1 and transfinite induction, we get the desired system S. \Box

Definition 3.3. ([22]) Let β be an ordinal number. A set S is called a *continuous union* of a family of subsets indexed by ordinals α with $\alpha < \beta$ if for each such α we have a subset $S_{\alpha} \subset S$ such that if $\alpha \leq \alpha'$ then $S_{\alpha} \subset S_{\alpha'}$, and such that if $\gamma < \beta$ is a limit ordinal then $S_{\gamma} = \bigcup_{\alpha < \gamma} S_{\alpha}$.

The following lemma is [12, Proposition 2.3] (cf. [15, Lemma 2.14]).

Lemma 3.4. If \mathscr{X} is a class of left *R*-modules closed under direct limits of well-ordered chains, then \mathscr{X} is closed under direct limits.

In Lemma 3.4, if $M = \underset{i \in I}{\underset{i : i I}{\underset{i \in I}{\underset{i : I}{\underset{i \in I}{\underset{i : : I}{\underset{i : : I}{\underset{i : : I}{\underset{i : I}{\underset{i : I}{\underset{i : : I}{\underset{i : : I}{\underset{i : I}{\underset{i : I}{\underset{i : I}{\underset{i : I}{\underset{i : : I}{\underset{i : I}{I}{I}{I}{I}{I}{I}{I}{I}{I}{I$

Our main result is the following theorem.

Theorem 3.5. Let $M \in ModR$ such that $M = \underset{i \in I}{\underset{i \in I}{\lim}} M_i$ with I a directed index set. Keep the notations as above.

(1) If

 $\mathbb{S}_i := 0 \to M_i \to E_i^0 \to E_i^1 \to \cdots \xrightarrow{f_i^n} E_i^n$

is an injective coresolution of M_i for any $i \in I$, then we have an exact sequence

$$\lim_{i \in I_{\alpha}} \mathbb{S}_{i} := 0 \to N_{\alpha} \to \lim_{i \in I_{\alpha}} E_{i}^{0} \to \lim_{i \in I_{\alpha}} E_{i}^{1} \to \dots \to \lim_{i \in I_{\alpha}} E_{i}^{n} \to \lim_{i \in I_{\alpha}} \operatorname{Coker} f_{i}^{n} \to 0.$$
(3.1)

Furthermore, if

$$\mathbb{S}'_{\alpha} := 0 \to N_{\alpha} \to E^0_{\alpha} \to E^1_{\alpha} \to \cdots \xrightarrow{f^n_{\alpha}} E^n_{\alpha}$$

is an injective coresolution of N_{α} , then we have the following exact sequence

$$\mathbb{S} := 0 \to M \to \varinjlim_{\alpha < \beta} E^0_{\alpha} \to \varinjlim_{\alpha < \beta} E^1_{\alpha} \to \dots \to \varinjlim_{\alpha < \beta} E^n_{\alpha} \to \varinjlim_{\alpha < \beta} \operatorname{Coker} f^n_{\alpha} \to 0.$$
(3.2)

(2) Let X be a subclass of ModR such that both X and ^{⊥1} X are closed under direct limits. If

$$\mathbb{S}_i := 0 \to M_i \to X_i^0 \to X_i^1 \to \cdots \xrightarrow{g_i^n} X_i^n$$

is a strong coproper $n-\mathscr{X}$ -coresolution of M_i for any $i \in I$, then N_{α} admits a partial strong coproper $n-\mathscr{X}$ -coresolution

$$\lim_{i \in I_{\alpha}} \mathbb{S}_{i} := 0 \to N_{\alpha} \to \lim_{i \in I_{\alpha}} X_{i}^{0} \to \lim_{i \in I_{\alpha}} X_{i}^{1} \to \dots \to \lim_{i \in I_{\alpha}} X_{i}^{n} \to \lim_{i \in I_{\alpha}} \operatorname{Coker} g_{i}^{n} \to 0; \qquad (3.3)$$

furthermore, M admits a partial strong coproper n- \mathscr{X} -coresolution

$$\mathbb{S} := 0 \to M \to \varinjlim_{\alpha < \beta} Y^0_{\alpha} \to \varinjlim_{\alpha < \beta} Y^1_{\alpha} \to \dots \to \varinjlim_{\alpha < \beta} Y^n_{\alpha} \to \varinjlim_{\alpha < \beta} C^n_{\alpha} \to 0, \tag{3.4}$$

where
$$Y_{\alpha}^{j} = \lim_{i \in I_{\alpha}} X_{i}^{j}$$
 and $C_{\alpha}^{n} = \lim_{i \in I_{\alpha}} \operatorname{Coker} g_{i}^{n}$ for any $0 \leq j \leq n$.

Proof. We prove it by transfinite induction on |I|. The case for $|I| < \infty$ is clear.

Suppose that $|I| = \aleph_0$ and $I = \{i_n \mid n \in \mathbb{N}\}$ with \mathbb{N} the set of non-negative integers. We construct a sequence j_0, j_1, \cdots of elements in I by letting $j_0 = i_0$, then we choose j_1 such that $j_1 \geq j_0, i_1$ by the upper directed set I. By induction, we choose $j_n \geq j_{n-1}, i_n$. Let $J = \{j_n \mid n \in \mathbb{N}\}$. Then J is cofinal well-ordered subset of I and

$$M = \lim_{\overrightarrow{i \in I}} M_i = \lim_{\overrightarrow{j \in J}} M_j.$$

The assertions follow from Lemma 3.2.

When $|I| > \aleph_0$, using Lemma 3.4, we may write $I = \bigcup_{\alpha < \beta} I_{\alpha}$ for some ordinal β and we have

$$M = \lim_{\overrightarrow{i \in I}} M_i = \lim_{\overrightarrow{\alpha < \beta}} N_{\alpha},$$

where $N_{\alpha} = \lim_{i \in I_{\alpha}} M_i$. Since $|I_{\alpha}| < |I|$ for each α , we get (3.1) and (3.3) by induction hypothesis.

For (1), there exists a chain map from $\lim_{i \in I_{\alpha}} \mathbb{S}_i$ to \mathbb{S}'_{α} as follows:

For (2), note that each N_{α} admits a strong coproper *n*- \mathscr{X} -coresolution. Thus we get (3.2) and (3.4) from Lemma 3.2.

Let R be a left Noetherian but not left Artinian ring with global dimension at least two (for example, the polynomial ring in n indeterminates over the ring of integers with $n \ge 1$). Then there exists a flat left R-module M which is not torsionless by [8, Theorem 4.1]. Note that $M = \underset{i \in I}{\lim M_i}$ with all M_i finitely generated projective left R-modules by [29, Thmorem 5.40]. Set $\mathscr{X} := \mathscr{P}(ModR)$. It is easy to see that each M_i admits a strong coproper 0- \mathscr{X} -coresolution. But M is not torsionless, so M does not admit a (strong) coproper 0- \mathscr{X} -coresolution. On the other hand, we have that R is not left perfect by [1, Corollary 15.23 and Theorem 28.4], and hence \mathscr{X} is not closed under direct limits. This means that the condition that the class \mathscr{X} is closed under direct limits in Theorem 3.5(2) is necessary.

Proposition 3.6. Let R be a left Noetherian ring, and let \mathscr{X} be a subclass of ModR and $M \in \text{Mod}R$ such that $M = \varinjlim_{i \in I} M_i$ with all M_i in modR. Set $\mathscr{X}' := \{ all \text{ finitely generated submodules of modules in } \mathscr{X} \}$. If both \mathscr{X} and $^{\perp_1}\mathscr{X}$ are closed under direct limits and each M_i admits a strong coproper $n \cdot \mathscr{X}'$ -coresolution, then M admits a strong coproper $n \cdot \mathscr{X}'$ -coresolution.

Proof. Let

$$0 \to M_i \xrightarrow{\varphi_i^0} X_i^0 \xrightarrow{\varphi_i^1} X_i^1 \longrightarrow \cdots \xrightarrow{\varphi_i^n} X_i^n$$
(3.5)

be a strong coproper $n \mathscr{X}'$ -coresolution of M_i for any $i \in I$. Set $K_i^j := \operatorname{Coker} \varphi_i^j$ for any $0 \leq j \leq n$. Then all K_i^j are in $^{\perp_1} \mathscr{X}'$. Notice that all K_i^j are in mod R, they are in $^{\perp_1} \mathscr{X}$ by [15, Lemma 6.6]. Thus (3.5) is a strong coproper $n \mathscr{X}$ -coresolution of M_i for any $i \in I$. Now the assertion follows from Theorem 3.5(2).

Note that any module is a direct limit of its finitely generated submodules. Thus by Theorem 3.5(2) and Proposition 3.6, we obtain the following result.

Corollary 3.7. Let R be a left Noetherian ring, and let \mathscr{X} be a subclass of ModR such that both \mathscr{X} and ${}^{\perp_1}\mathscr{X}$ are closed under direct limits. If one of the following conditions is satisfied, then \mathscr{X} is special preenveloping in ModR.

- (1) Any module in $\operatorname{mod} R$ admits a special \mathscr{X} -preenvelope.
- (2) \mathscr{X}' is special preenveloping in mod R, where $\mathscr{X}' = \{all \text{ finitely generated submodules of modules in } \mathscr{X}\}.$

In the following, we list the dual counterparts of Lemma 3.2, Theorem 3.5(2), Proposition 3.6 and Corollary 3.7. Since their proofs are completely dual to those of the previous corresponding results, we omit them.

Lemma 3.8. (Dual to Lemma 3.2) Let \mathcal{X} be a subclass of ModR, and let κ be an ordinal number. Suppose that $\{M_{\alpha}, f_{\beta\alpha} : M_{\alpha} \to M_{\beta} \mid \alpha \leq \beta < \kappa\}$ is a direct system in ModR and

$$\mathbb{S}_{\alpha} := X_{\alpha}^{n} \xrightarrow{\varphi_{\alpha}^{n}} \cdots \to X_{\alpha}^{1} \to X_{\alpha}^{0} \xrightarrow{\varphi_{\alpha}^{0}} M_{\alpha} \to 0$$

is an exact sequence in ModR with all X_{α}^{i} in \mathcal{X} . If \mathbb{S}_{α} is a strong proper n- \mathcal{X} -resolution of M_{α} and both \mathcal{X} and $\mathcal{X}^{\perp_{1}}$ are closed under direct limits, then these exact sequences \mathbb{S}_{α} are the members of a direct system indexed by $\alpha < \kappa$ in such a way that if $\alpha \leq \beta < \kappa$, the map from the sequence indexed by α into that indexed by β with the origin map $f_{\beta\alpha} : M_{\alpha} \to M_{\beta}$. In particular, we obtain a partial strong proper n- \mathcal{X} -resolution of $\lim M_{\alpha}$:

$$\lim_{\alpha < \kappa} \mathbb{S}_{\alpha} : 0 \to \lim_{\alpha < \kappa} \operatorname{Ker} \varphi_{\alpha}^{n} \to \lim_{\alpha < \kappa} X_{\alpha}^{n} \to \cdots \to \lim_{\alpha < \kappa} X_{\alpha}^{1} \to \lim_{\alpha < \kappa} X_{\alpha}^{0} \to \lim_{\alpha < \kappa} M_{\alpha} \to 0.$$

$$\mathbb{S}_i := X_i^n \xrightarrow{g_i^n} \cdots \to X_i^1 \to X_i^0 \to M_i \to 0$$

is a strong proper $n-\mathcal{X}$ -resolution of M_i for any $i \in I$, then N_{α} admits a partial strong proper $n-\mathcal{X}$ -resolution

$$\lim_{i \in I_{\alpha}} \mathbb{S}_{i} := 0 \to \lim_{i \in I_{\alpha}} \operatorname{Ker} g_{i}^{n} \to \lim_{i \in I_{\alpha}} X_{i}^{0} \to \cdots \to \lim_{i \in I_{\alpha}} X_{i}^{1} \to \lim_{i \in I_{\alpha}} X_{i}^{0} \to N_{\alpha} \to 0;$$

furthermore, M admits a partial strong proper $n-\mathcal{X}$ -resolution

$$\mathbb{S} := 0 \to \varinjlim_{\alpha < \beta} C^n_{\alpha} \to \varinjlim_{\alpha < \beta} Y^0_{\alpha} \to \dots \to \varinjlim_{\alpha < \beta} Y^1_{\alpha} \to \varinjlim_{\alpha < \beta} Y^n_{\alpha} \to M \to 0,$$

where $Y_{\alpha}^{j} = \underset{i \in I_{\alpha}}{\lim} X_{i}^{j}$ and $C_{\alpha}^{n} = \underset{i \in I_{\alpha}}{\lim} \operatorname{Ker} g_{i}^{n}$ for any $0 \leq j \leq n$.

Note that all finitely generated modules are pure injective over Artinian algebras.

Proposition 3.10. (Dual to Proposition 3.6) Let R be an Artinian algebra and \mathcal{X} be a subclass of ModR which is closed under direct limits, and let $M \in \text{Mod}R$ such that $M = \lim_{i \in I} M_i$ with all M_i in modR. Set $\mathcal{X}' := \{ all \text{ finitely generated submodules of modules} in <math>\mathcal{X} \}$. If \mathcal{X}^{\perp_1} is closed under direct limits and each M_i admits a strong proper $n-\mathcal{X}'$ resolution, then M admits a strong proper $n-\mathcal{X}$ -resolution.

Corollary 3.11. (Dual to Corollary 3.7) Let R be an Artinian algebra, \mathcal{X} be a subclass of ModR such that both \mathcal{X} and \mathcal{X}^{\perp_1} are closed under direct limits. If one of the following conditions is satisfied, then \mathcal{X} is special precovering in ModR.

- (1) Any module in mod R admits a special \mathcal{X} -precover.
- (2) \mathcal{X}' is special precovering in mod R, where $\mathcal{X}' = \{all \text{ finitely generated submodules of modules in } \mathcal{X}\}.$

In the final of this section, we raise the following question.

Question 3.12. Is there a dual counterpart of Theorem 3.5(1)?

4. Applications

In this section, we give some applications of the results obtained in Section 3.

4.1. Relative injective dimension. As an application of Theorem 3.5(1), we get the following result.

Theorem 4.1. Let \mathscr{X} be a subclass of ModR which is coresolving and closed under direct limits. Then

$$\sup\{\mathscr{X} - \mathrm{id}M \mid M \in \mathrm{Mod}R\} = \sup\{\mathscr{X} - \mathrm{id}M \mid M \in \mathrm{mod}R\},\$$

Proof. It is trivial that $\sup \{\mathscr{X} - \operatorname{id} M \mid M \in \operatorname{Mod} R\} \ge \sup \{\mathscr{X} - \operatorname{id} M \mid M \in \operatorname{mod} R\}.$

Now suppose $\sup \{\mathscr{X} - \operatorname{id} M \mid M \in \operatorname{mod} R\} = n < \infty$. Let $M \in \operatorname{Mod} R$. Then $M = \lim_{i \in I} M_i$ with all M_i finitely presented left *R*-modules by [15, Lemma 2.5]. For any $i \in I$, we have $\mathscr{X} - \operatorname{id} M_i \leq n$. Since \mathscr{X} is coresolving, there exists an exact sequence

$$0 \to M_i \to E_i^0 \to E_i^1 \to \dots \to E_i^n \to X_i^{n+1} \to 0$$

in ModR with all E_i^j injective and X_i^{n+1} in \mathscr{X} by the dual version of [36, Lemma 2.1] (cf. the dual version of [4, Lemma 3.12]). Keep the notations N_{α} and I_{α} as in Theorem 3.5(1). Then we get an exact sequence

$$0 \to N_{\alpha} \to \varinjlim_{i \in I_{\alpha}} E_i^0 \to \varinjlim_{i \in I_{\alpha}} E_i^1 \to \dots \to \varinjlim_{i \in I_{\alpha}} E_i^n \to \varinjlim_{i \in I_{\alpha}} X_i^{n+1} \to 0.$$

Since \mathscr{X} is closed under direct limits, we have that all $\lim_{i \in I_{\alpha}} E_{i}^{j}$ and $\lim_{i \in I_{\alpha}} X_{i}^{n+1}$ are in \mathscr{X} , and thus \mathscr{X} -id $N_{\alpha} \leq n$. As above, there exists an exact sequence

$$0 \to N_{\alpha} \to E_{\alpha}^{0} \to E_{\alpha}^{1} \to \dots \to E_{\alpha}^{n} \to X_{\alpha}^{n+1} \to 0$$

in ModR with all E^j_{α} injective and X^{n+1}_{α} in \mathscr{X} , which induces an exact sequence

$$0 \to M \to \varinjlim_{\alpha < \beta} E^0_{\alpha} \to \varinjlim_{\alpha < \beta} E^1_{\alpha} \to \dots \to \varinjlim_{\alpha < \beta} E^n_{\alpha} \to \varinjlim_{\alpha < \beta} X^{n+1}_{\alpha} \to 0$$

with all $\varinjlim_{\alpha < \beta} E_{\alpha}^{j}$ and $\varinjlim_{i \in I_{\alpha}} X_{\alpha}^{n+1}$ are in \mathscr{X} . Thus \mathscr{X} -id $M \leq n$, and the assertion follows. \Box

Recall that a module $M \in ModR$ is called *weak injective* [13], or *absolutely clean* [6], if $\operatorname{Ext}^1_R(A, M) = 0$ for any left *R*-module *A* admitting a degreewise finite *R*-projective resolution. We use $\mathcal{WI}(ModR)$ to denote the class of weak injective left *R*-modules. Recall from [32] that a module $M \in ModR$ is called *FP-injective* (or *absolutely pure*) if $\operatorname{Ext}^1_R(A, M) = 0$ for any finitely presented left *R*-module *A*. If *R* is a left Noetherian ring, then the class $\mathcal{I}(ModR)$ of injective left *R*-modules coincides with $\mathcal{WI}(ModR)$, and if *R* is a left coherent ring, then the class $\mathcal{FI}(ModR)$ of FP-injective left *R*-modules coincides with $\mathcal{WI}(ModR)$. Recall that a ring R is called *left* Π -*coherent* if any finitely generated torsionless left R-module is finitely presented, and a module $M \in \text{Mod}R$ is called *FGT-injective* if $\text{Ext}^1_R(A, M) = 0$ for any finitely generated torsionless left R-module A [7, 9]. We use $\mathcal{FTI}(\text{Mod}R)$ to denote the class of FGT-injective left R-modules.

Let R and S be rings. An (R, S)-bimodule ${}_{R}C_{S}$ is called *semidualizing* if the following conditions are satisfied: (1) ${}_{R}C$ admits a degreewise finite R-projective resolution and C_{S} admits a degreewise finite S^{op} -projective resolution, (2) the homothety maps ${}_{R}R_{R} \xrightarrow{R\gamma}$ $\operatorname{Hom}_{S^{op}}(C, C)$ and ${}_{S}S_{S} \xrightarrow{\gamma_{S}} \operatorname{Hom}_{R}(C, C)$ are isomorphisms, and (3) $\operatorname{Ext}_{R}^{\geq 1}(C, C) = 0 =$ $\operatorname{Ext}_{S^{op}}^{\geq 1}(C, C)$. The Bass class $\mathcal{B}_{C}(R)$ with respect to C consists of all left R-modules Msatisfying the following conditions: (1) $\operatorname{Ext}_{R}^{\geq 1}(C, M) = 0$, (2) $\operatorname{Tor}_{\geq 1}^{S}(C, \operatorname{Hom}_{R}(C, M)) = 0$, and (3) the canonical evaluation homomorphism $\theta_{M} : C \otimes_{S} \operatorname{Hom}_{R}(C, M) \to M$ defined by $\theta_{M}(x \otimes f) = f(x)$ for any $x \in C$ and $f \in \operatorname{Hom}_{R}(C, M)$ is an isomorphism of left R-modules [17].

We collect some known facts that we need to use.

Fact 4.2. It holds that

- (1) By [6, Lemma 2.7(3)(4)], the class $\mathcal{WI}(ModR)$ is coresolving and closed under direct limits.
- (2) If R is a left Π -coherent ring, then the class $\mathcal{FTI}(\mathrm{Mod}R)$ is coresolving and closed under direct limits by [9, Propositions 1.4 and 2.2].
- (3) The class $\mathcal{GI}(ModR)$ of Gorenstein injective left *R*-modules is coresolving by [16, Theorem 2.6]. If *R* is a left Artinian ring such that the injective envelope of every simple left *R*-module is finitely generated (in particular, if *R* is an Artinian algebra), then $\mathcal{GI}(ModR)$ is closed under direct limits by [21, Theorem 2] and [25, Theorem 2.3].
- (4) Recall that a module $T \in \text{Mod}R$ is called *tilting* if the following conditions are satisfied: (i) the projective dimension of T is finite; (ii) $\text{Ext}_R^{\geq 1}(T, T^{(I)}) = 0$ for any set I; and (iii) there exists an exact sequence

$$0 \to R \to T^0 \to T^1 \to \dots \to T^n \to 0$$

in ModR with all T^i direct summands of direct sums of copies of T. Let $T \in ModR$ be tilting. Then T^{\perp} is clearly coresolving, and it is closed under direct limits by [15, Corollary 13.42].

- (5) Let *PP*(*R*) be the class of pure projective left *R*-modules, then *PP*(*R*)[⊥] is coresolving by [33, Proposition 39]. If *R* is left coherent, then *PP*(*R*)[⊥] is closed under direct limits [33, Theorem 47].
- (6) The Bass class $\mathcal{B}_C(R)$ with respect to a semidualizing bimodule ${}_RC_S$ is coresolving and closed under direct limits [17, Theorem 6.2 and Proposition 4.2(a)].

Following the usual customary notation, we write

$$w\text{-}\mathrm{id}_R M := \mathcal{WI}(\mathrm{Mod}R)\text{-}\mathrm{id}M, \quad \mathrm{id}_R M := \mathcal{I}(\mathrm{Mod}R)\text{-}\mathrm{id}M,$$
$$FP\text{-}\mathrm{id}_R M := \mathcal{FI}(\mathrm{Mod}R)\text{-}\mathrm{id}M, \quad FGT\text{-}\mathrm{id}_R M := \mathcal{FII}(\mathrm{Mod}R)\text{-}\mathrm{id}M,$$
$$G\text{-}\mathrm{id}_R M := \mathcal{GI}(\mathrm{Mod}R)\text{-}\mathrm{id}M.$$

Let $M \in ModR$. If R is a left Noetherian ring, then w-id_RM = id_RM. If R is a left coherent ring, then w-id_RM = FP-id_RM.

By Theorem 4.1 and Fact 4.2, we obtain the following result.

Corollary 4.3. It holds that

(1) $\sup\{\text{w-id}_R M \mid M \in \text{Mod}R\} = \sup\{\text{w-id}_R M \mid M \in \text{mod}R\}$. In particular, we have (a) ([27, Theorem C]) If R is a left Noetherian ring, then

 $\sup\{\mathrm{id}_R M \mid M \in \mathrm{Mod}R\} = \sup\{\mathrm{id}_R M \mid M \in \mathrm{mod}R\}.$

(b) ([32, Theorem 3.3]) If R is a left coherent ring, then

 $\sup\{\operatorname{FP-id}_R M \mid M \in \operatorname{Mod} R\} = \sup\{\operatorname{FP-id}_R M \mid M \in \operatorname{mod} R\}.$

(2) If R is a left Π -coherent ring, then

 $\sup\{\operatorname{FGT-id}_R M \mid M \in \operatorname{Mod} R\} = \sup\{\operatorname{FGT-id}_R M \mid M \in \operatorname{mod} R\}.$

(3) If R is a left Artinian ring such that the injective envelope of every simple left R-module is finitely generated (in particular, if R is an Artinian algebra), then

 $\sup\{\operatorname{G-id}_R M \mid M \in \operatorname{Mod} R\} = \sup\{\operatorname{G-id}_R M \mid M \in \operatorname{mod} R\}.$

(4) If $T \in ModR$ is a tilting module, then

 $\sup\{T^{\perp} \operatorname{-id} M \mid M \in \operatorname{Mod} R\} = \sup\{T^{\perp} \operatorname{-id} M \mid M \in \operatorname{mod} R\}.$

(5) If R is a left coherent ring, then

$$\sup\{\mathcal{PP}(R)^{\perp} \operatorname{-id} M \mid M \in \operatorname{Mod} R\} = \sup\{\mathcal{PP}(R)^{\perp} \operatorname{-id} M \mid M \in \operatorname{mod} R\}.$$

(6) We have

$$\sup\{\mathcal{B}_C(R) \text{-}\mathrm{id}M \mid M \in \mathrm{Mod}R\} = \sup\{\mathcal{B}_C(R) \text{-}\mathrm{id}M \mid M \in \mathrm{mod}R\}.$$

Recall from [26] that a module $M \in \text{Mod}R$ is called *strong Gorenstein injective*, which is usually called *Ding injective* [14, 21], if $M \in \mathcal{FI}(\text{Mod}R)^{\perp}$ and there exists a $\text{Hom}_R(\mathcal{FI}(\text{Mod}R), -)$ -exact exact sequence

$$0 \to M \to E^0 \to E^1 \to \dots \to E^i \to \dots$$

in ModR with all E^i in $\mathcal{I}(ModR)$. Recall from [11] that a module $N \in ModR^{op}$ is called Gorenstein flat if there exists an exact sequence

$$\cdots \to F_i \to \cdots \to F_1 \to F_0 \to F^0 \to F^1 \to \cdots \to F^i \to \cdots$$

in Mod R^{op} with all F^i flat, such that it remains exact after applying the functor $-\otimes_R E$ for any $E \in \mathcal{I}(ModR)$, and $N \cong Im(F_0 \to F^0)$. For a module $M \in ModR$, we call $Hom_{\mathbb{Z}}(M, \mathbb{Q}/\mathbb{Z})$ its *character module*, where \mathbb{Z} is the additive group of integers and \mathbb{Q} is the additive group of rational numbers.

Note that the class SGI(ModR) of strong Gorenstein injective left *R*-modules is coresolving by [18, Remark 4.4(4)(b)]. About its direct limit closure, we have the following result, which extends [21, Theorem 2].

Proposition 4.4. The following statements are equivalent.

- (1) $\mathcal{SGI}(ModR)$ is closed under direct limits.
- (2) $\mathcal{GI}(ModR)$ is closed under direct limits.
- (3) *R* is a left Noetherian ring and the character module of any Gorenstein injective left *R*-module is Gorenstein flat.

Proof. The equivalence $(2) \iff (3)$ has been proved in [21, Theorem 2]. It is well known that $\mathcal{FI}(ModR) = \mathcal{I}(ModR)$ if R is a left Noetherian ring, thus we have $(2) + (3) \Longrightarrow (1)$.

Now suppose that the assertion (1) holds true, to prove that (2) also holds true, it suffices to prove that R is a left Noetherian ring by the above argument. Since SGI(ModR)is closed under direct limits by (1), we have that SGI(ModR) is closed under direct products and pure submodules by [14, Theorem 44] and [30, Theorem 3.5]. Notice that the direct sum of modules is a pure submodule of the direct product of the modules, so SGI(ModR) is closed under direct sums.

Let $\{E_i \mid i \in I\}$ be a family of injective left *R*-modules. Then $\bigoplus_{i \in I} E_i \in SGI(ModR)$, and thus there exists a $\operatorname{Hom}_R(\mathcal{FI}(ModR), -)$ -exact exact sequence

$$E \xrightarrow{\varphi} \bigoplus_{i \in I} E_i \to 0$$

in Mod R with $E \in \mathcal{I}(Mod R)$. For each standard embedding $\lambda_i : E_i \to \bigoplus_{i \in I} E_i$, there exists $f_i \in \operatorname{Hom}_R(E_i, E)$ such that $\varphi f_i = \lambda_i$. By the universal property of direct sums, there exists $\varphi' \in \operatorname{Hom}_R(\bigoplus_{i \in I} E_i, E)$ such that $\varphi' \lambda_i = f_i$, and thus

$$(\varphi\varphi')\lambda_i = \varphi f_i = \lambda_i.$$

It yields that $\varphi \varphi'$ is the identity homomorphism of $\bigoplus_{i \in I} E_i$ and φ is a split epimorphism. So $\bigoplus_{i \in I} E_i$ is a direct summand of E, and hence it is injective. It follows from [5, Theorem 1.1] that R is a left Noetherian ring.

For a module $M \in ModR$, we use $fd_R M$ to denote the flat dimension of M. The assertion (1) in the following result generalizes [19, Theorem 3.1].

Proposition 4.5. Let $M \in \text{Mod}R$ such that $M = \underset{i \in I}{\underset{i I}{\underset{i \in I}{\underset{i \in I}{\underset{i \in I}{\underset{i I}{\underset{i \in I}{\underset{i I}{\underset$

$$0 \to M_i \to X_i^0 \to X_i^1 \to \dots \to X_i^n$$

$$0 \to M \to X^0 \to X^1 \to \dots \to X^n$$

such that for any $j \ge 0$, it holds that

- (1) $\operatorname{fd}_R X^j = \sup\{\operatorname{fd}_R X^j_i \mid i \in I\}.$
- (2) w-id_R X^j = sup{w-id_R $X^j_i | i \in I$ }.

In particular, if \mathscr{X} is enveloping, then a minimal coproper n- \mathscr{X} -coresolution of M as (3.3) also satisfies (1) and (2).

Proof. (1) Since the functor Tor commutes with direct limits, the assertion follows from Theorem 3.5.

(2) For any $X \in \text{Mod}R$ and $n \ge 0$, it is easy to see that w-id_RX = n if and only if $n = \inf\{i \mid \text{Ext}_R^{\ge i+1}(A, X) = 0$ for any left *R*-module *A* admitting a degreewise finite *R*-projective resolution}. Now the assertion follows from Theorem 3.5(2) and [15, Lemma 6.6].

According to (1) and (2), the last assertion follows from Lemma 2.3.

4.2. Weakly Gorenstein algebras. As an application of Proposition 3.6, we have the following result.

Proposition 4.6. If R is a left and right Artinian ring, then the following statements are equivalent.

- (1) $\mathcal{GP}(\mathrm{mod}R) = {}^{\perp}_{R}R \cap \mathrm{mod}R.$
- (2) $\mathcal{GP}(\mathrm{Mod}R) = {}^{\perp}_{R}R.$

Proof. The implication $(2) \Longrightarrow (1)$ is clear.

(1) \Longrightarrow (2) Let $M \in {}^{\perp}{}_{R}R$. Then $M = \lim_{i \in I} M_{i}$ with all M_{i} finitely generated submodules of M. Since R is a right Artinian ring, it follows from [1, Theorem 28.4 and Corollary 28.8] that a left R-module is flat if and only if it is projective, and thus any projective left R-module is pure injective by [34, Lemma 3.1.6]. Then ${}^{\perp_{1}}\mathcal{P}(ModR)$ is closed under direct limits and

$$\lim_{i \in I} \operatorname{Ext}_{R}^{j}(M_{i}, R) \cong \operatorname{Ext}_{R}^{j}(\lim_{i \in I} M_{i}, R) = \operatorname{Ext}_{R}^{j}(M, R) = 0$$

for any $j \ge 1$ by [3, Proposition I.10.1]. So for any $i \in I$ and $j \ge 1$, we have $\operatorname{Ext}_R^j(M_i, R) = 0$, that is, $M_i \in {}^{\perp}_R R \cap \operatorname{mod} R$, and hence $M_i \in \mathcal{GP}(\operatorname{mod} R)$ by (1). Thus each M_i admits a strong coproper ∞ - $\mathcal{P}(\operatorname{mod} R)$ -coresolution. On the other hand, the class $\mathcal{P}(\operatorname{Mod} R)$ is closed under direct limits by [23, Proposition 4.4]. Since R is a left Artinian ring, it follows from Proposition 3.6 that M admits a strong coproper ∞ - $\mathcal{P}(\operatorname{Mod} R)$ -coresolution and $M \in \mathcal{GP}(\operatorname{Mod} R)$.

For an Artinian algebra R, we use \mathbb{D} to denote the usual duality between mod R and mod R^{op} . We need the following easy observation.

Lemma 4.7. Let R be an Artinian algebra, and let $M \in \text{mod}R$ and $M' \in \text{Mod}R$. Then for any $i \ge 1$, it holds that

$$\operatorname{Ext}_{R^{op}}^{i}(\mathbb{D}(M'),\mathbb{D}(M)) \cong \mathbb{D}^{2}\operatorname{Ext}_{R}^{i}(M,M'),$$

in particular, if $M' \in \text{mod}R$, then

$$\operatorname{Ext}_{R^{op}}^{i}(\mathbb{D}(M'), \mathbb{D}(M)) \cong \operatorname{Ext}_{R}^{i}(M, M').$$

Proof. For any $M \in \text{mod}R$, $M' \in \text{Mod}R$ and $i \ge 1$, we have

$$\begin{aligned} &\operatorname{Ext}_{R^{op}}^{i}(\mathbb{D}(M'),\mathbb{D}(M)) \\ &\cong \mathbb{D}\operatorname{Tor}_{i}^{R}(\mathbb{D}(M'),M) \quad \text{(by [15, Lemma 2.16(b)])} \\ &\cong \mathbb{D}^{2}\operatorname{Ext}_{R}^{i}(M,M'). \quad \text{(by [15, Lemma 2.16(d)])} \end{aligned}$$

If $M, M' \in \text{mod}R$, then $\mathbb{D}^2\text{Ext}^i_R(M, M') \cong \text{Ext}^i_R(M, M')$, and thus the latter assertion follows.

Recall from [28] that an Artinian algebra R is called *left weakly Gorenstein* if $\mathcal{GP}(\text{mod}R) = {}^{\perp}_R R \cap \text{mod}R$. A Gorenstein algebra R (that is, $\text{id}_R R = \text{id}_{R^{op}} R < \infty$) is left weakly Gorenstein, but the converse does not holds true in general [20, 28]. In the following result, we give some equivalent characterizations of weakly Gorenstein algebras, which generalizes part of [20, Theorem 4.9] (that is, the equivalence (4) \iff (5) there).

Theorem 4.8. For an Artinian algebra R, the following statements are equivalent.

- (1) R is left weakly Gorenstein, that is, $\mathcal{GP}(\text{mod}R) = {}^{\perp}_{R}R \cap \text{mod}R$.
- (2) $\mathcal{GP}(\mathrm{Mod}R) = {}^{\perp}_{R}R.$
- (3) $\mathcal{GP}(\mathrm{Mod}R) = {}^{\perp}\mathcal{P}(\mathrm{Mod}R).$
- (4) $\mathcal{GI}(\mathrm{mod}R^{op}) = \mathbb{D}(_RR)^{\perp} \cap \mathrm{mod}R^{op}.$
- (5) $\mathcal{GI}(\mathrm{Mod}R^{op}) = \mathbb{D}(_RR)^{\perp}.$
- (6) $\mathcal{GI}(\mathrm{Mod}R^{op}) = \mathcal{I}(\mathrm{Mod}R^{op})^{\perp}$.

Proof. The equivalence $(1) \iff (2)$ follows from Proposition 4.6, and the implication $(5) \implies (4)$ is clear. Since

 $\mathcal{GP}(\mathrm{Mod}R) \subseteq {}^{\perp}\mathcal{P}(\mathrm{Mod}R) \subseteq {}^{\perp}{}_{R}R \text{ and } \mathcal{GI}(\mathrm{Mod}R^{op}) \subseteq \mathcal{I}(\mathrm{Mod}R^{op})^{\perp} \subseteq \mathbb{D}({}_{R}R)^{\perp},$

we have $(2) \Longrightarrow (3)$ and $(5) \Longrightarrow (6)$.

(3) \Longrightarrow (1) Let $M \in {}^{\perp}_{R}R \cap \operatorname{mod} R$. Then $M \in {}^{\perp}\mathcal{P}(\operatorname{Mod} R)$ by [32, Theorem 3.2], and hence $M \in \mathcal{GP}(\operatorname{Mod} R) \cap \operatorname{mod} R = \mathcal{GP}(\operatorname{mod} R)$ by (3).

 $(4) \Longrightarrow (1)$ Let $M \in {}^{\perp}_{R}R \cap \operatorname{mod} R$. Then we have

$$\operatorname{Ext}_{R^{op}}^{i}(\mathbb{D}(_{R}R),\mathbb{D}(M))\cong\operatorname{Ext}_{R}^{i}(M,_{R}R)=0$$

for any $i \ge 1$ by Lemma 4.7, so $\mathbb{D}(M) \in \mathbb{D}(_R R)^{\perp} \cap \operatorname{mod} R^{op} = \mathcal{GI}(\operatorname{mod} R^{op})$ by (4). Thus $M \in \mathcal{GP}(\operatorname{mod} R)$ by [16, Theorem 3.6] and [35, Corollary 3.7]. (1) \Longrightarrow (4) Let $N \in \mathbb{D}(_R R)^{\perp} \cap \operatorname{mod} R^{op}$. Then we have

$$\operatorname{Ext}_{R}^{i}(\mathbb{D}(N), {}_{R}R) \cong \operatorname{Ext}_{R}^{i}(\mathbb{D}(N), \mathbb{D}^{2}({}_{R}R)) \cong \operatorname{Ext}_{R^{op}}^{i}(\mathbb{D}({}_{R}R), N) = 0$$

for any $i \geq 1$ by Lemma 4.7, so $\mathbb{D}(N) \in {}^{\perp}_{R}R \cap \operatorname{mod}R = \mathcal{GP}(\operatorname{mod}R)$ by (1). It follows from [16, Theorem 3.6] and [35, Corollary 3.7] that $N \cong \mathbb{D}^{2}(N) \in \mathcal{GI}(\operatorname{mod}R^{op})$.

(4) \Longrightarrow (5) Let $N \in \mathbb{D}(RR)^{\perp}$. Then $N = \lim_{\substack{i \in I \\ i \in I}} N_i$ with all N_i finitely generated submodulos of N. Since

ules of N. Since

$$\lim_{i \in I} \operatorname{Ext}_{R^{op}}^{j}(\mathbb{D}(_{R}R), N_{i}) \cong \operatorname{Ext}_{R^{op}}^{j}(\mathbb{D}(_{R}R), \lim_{i \in I} N_{i}) = \operatorname{Ext}_{R^{op}}^{j}(\mathbb{D}(_{R}R), N) = 0$$

for any $j \geq 1$ by [15, Lemma 6.6], we have $\operatorname{Ext}_{R^{op}}^{\geq 1}(\mathbb{D}(RR), N_i) = 0$, and hence $N_i \in \mathbb{D}(RR)^{\perp} \cap \operatorname{mod} R^{op} = \mathcal{GI}(\operatorname{mod} R^{op})$ for any $i \in I$ by (4). It follows from Fact 4.2(3) that $N \in \mathcal{GI}(\operatorname{Mod} R^{op})$.

(6) \Longrightarrow (4) Let $N \in \mathbb{D}(R^{n})^{\perp} \cap \operatorname{mod} R^{op}$. Then for any $i \geq 1$, we have

$$\operatorname{Ext}^{i}_{R}(\mathbb{D}(N), {}_{R}R) \cong \operatorname{Ext}^{i}_{R^{op}}(\mathbb{D}({}_{R}R), \mathbb{D}^{2}(N)) \cong \operatorname{Ext}^{i}_{R^{op}}(\mathbb{D}({}_{R}R), N) = 0$$

by Lemma 4.7. It follows from [32, Theorem 3.2] that $\operatorname{Ext}_{R}^{i}(\mathbb{D}(N), \mathcal{P}(\operatorname{Mod} R)) = 0$. Then for any set J, we have

$$\operatorname{Ext}_{R^{op}}^{i}(\mathbb{D}(_{R}R)^{J}, N) \cong \operatorname{Ext}_{R^{op}}^{i}(\mathbb{D}(_{R}R^{(J)}), N) \cong \operatorname{Ext}_{R^{op}}^{i}(\mathbb{D}(_{R}R^{(J)}), \mathbb{D}^{2}(N))$$
$$\cong \mathbb{D}^{2}\operatorname{Ext}_{R}^{i}(\mathbb{D}(N), _{R}R^{(J)}) \quad \text{(by Lemma 4.7)}$$
$$= 0.$$

Since any modules in $\mathcal{I}(\operatorname{Mod} R^{op})$ is a direct summand of $\mathbb{D}({}_{R}R)^{J}$ for some set J, we have that $N \in \mathcal{I}(\operatorname{Mod} R^{op})^{\perp}$, and hence $N \in \mathcal{GI}(\operatorname{Mod} R^{op}) \cap \operatorname{mod} R = \mathcal{GI}(\operatorname{mod} R^{op})$ by (6). \Box

As a consequence, we obtain the following result.

Corollary 4.9. If R is an Artinian algebra with $id_R R < \infty$, then the following statements are equivalent.

- (1) R is Gorenstein.
- (2) R is left weakly Gorenstein.
- (3) $\mathcal{GP}(\mathrm{Mod}R) = {}^{\perp}_{R}R.$
- (4) $\mathcal{GI}(\mathrm{Mod}R^{op}) = \mathbb{D}(_RR)^{\perp}.$

Proof. The assertion $(2) \iff (3) \iff (4)$ follows from Theorem 4.8, and the assertion $(1) \iff (3)$ follows from [2, Proposition 3.10].

References

- F. W. Anderson and K. R. Fuller, Rings and Categories of Modules, Second edition, Graduate Texts in Math. 13, Springer-Verlag, New York-Berlin-Heidelberg, 1992.
- [2] L. Angeleri Hügel, D. Herbera and J. Trlifaj, *Tilting modules and Gorenstein rings*, Forum Math. 18 (2006), 211–229.
- [3] M. Auslander, Functors and morphisms determined by objects, in Representation Theory of Algebras (Proc. Conf., Temple Univ., Philadelphia, Pa., 1976), Lecture Notes in Pure Appl. Math. 37, Dekker, New York, 1978, pp.1–244.
- [4] M. Auslander and M. Bridger, Stable module theory, Memoirs Amer. Math. Soc. 94, Amer. Math. Soc., Providence, RI, 1969.
- [5] H. Bass, Injective dimension in Noetherian rings, Trans. Amer. Math. Soc. 102 (1962), 18–29.
- [6] D. Bravo, J. Gillespie and M. Hovey, *The stable module category of a general ring*, Preprint is available at: arXiv:1405.5768.
- [7] V. Camillo, Coherence for polynomial rings, J. Algebra 132 (1990), 72–76.
- [8] S. U. Chase, Direct products of modules, Trans. Amer. Math. Soc. 97 (1960), 457–473.
- [9] F. C. Cheng, J. Y. Wang, Z. Y. Huang and M. Y. Wang, Π-coherent rings and FGT-injective dimension, Southeast Asian Bull. Math. 19 (1995), 105–112.
- [10] E. E. Enochs, Injective and flat covers, envelopes and resolvents, Israel J. Math. 39 (1981), 189–209.
- [11] E. E. Enochs and O. M. G. Jenda, Relative Homological Algebra, Vol. 1, the second revised and extended edition, de Gruyter Expositions in Math. 30, Walter de Gruyter GmbH & Co. KG, Berlin, 2011.
- [12] E. E. Enochs and J. A. López-Ramos, *Kaplansky classes*, Rend. Sem. Mat. Univ. Padova 107 (2002) 67-79.
- [13] Z. H. Gao and F. G. Wang, Weak injective and weak flat modules, Comm. Algebra 43 (2015), 3857–3868.
- [14] J. Gillespie and A. Iacob, Duality pairs, generalized Gorenstein modules, and Ding injective envelopes, C. R. Math. Acad. Sci. Paris 360 (2022), 381–398.
- [15] R. Göbel and J. Trlifaj, Approximations and Endomorphism Algebras of Modules, the second revised and extended edition, de Gruyter Expositions in Math. 41, Walter de Gruyter GmbH & Co. KG, Berlin, 2012.
- [16] H. Holm, Gorenstein homological dimensions, J. Pure Appl. Algebra 189 (2004), 167–193.
- [17] H. Holm and D. White, Foxby equivalence over associative rings, J. Math. Kyoto Univ. 47 (2007), 781–808.
- [18] Z. Y. Huang, Homological dimensions relative to preresolving subcategories II, Forum Math. 34 (2022), 507–530.
- [19] Z. Y. Huang, On Auslander-type conditions of modules, Publ. Res. Inst. Math. Sci. 59 (2023), 57–88.
- [20] Z. Y. Huang, Auslander-type conditions and weakly Gorenstein algebras, Bull. London Marh. Soc. 56 (2024), 3382–3399.
- [21] A. Iacob, Direct limits of Gorenstein injective modules, arXiv:2308.08699v1.
- [22] C. U. Jensen, Les Foncteurs Dérivés de lim Et Leurs Applications En Théorie Des Modules. Lect. Notes in Math. 254, Springer-Verlag, Berlin, 1972.
- [23] T. Y. Lam, Lectures on Modules and Rings, Grad. Texts in Math. 189, Springer-Verlag, New York, 1999.
- [24] D. Lazard, Sur les modules plats, C. R. Acad. Sci. Paris 258 (1964), 6313-6316.

- [25] Z. K. Liu and X. Y. Yang, Gorenstein projective, injective and flat modules, J. Aust. Math. Soc. 87 (2009), 395–407.
- [26] L. X. Mao and N. Q. Ding, Gorenstein FP-injective and Gorenstein flat modules, J. Algebra Appl. 7 (2008), 491–506.
- [27] B. Osofsky, Global dimension of valuation rings, Trans. Amer. Math. Soc. 127 (1967), 136–149.
- [28] C. M. Ringel and P. Zhang, Gorenstein-projective modules and semi-Gorenstein-projective modules, Algebra Number Theory 14 (2020), 1–36.
- [29] J. J. Rotman, An Introduction to Homological Algebra, the second edition. Universitext, Springer, New York, 2009.
- [30] J. Šaroch and J. Štovíček, The countable telescope conjecture for module categories, Adv. Math. 219 (2008), 1002–1036.
- [31] J. Šaroch and J. Štovíček, Singular compactness and definability for Σ-cotorsion and Gorenstein modules, Selecta Math. (N. S.) 26 (2020), no. 2, Paper No. 23.
- [32] B. Stenström, Coherent rings and FP-injective modules, J. London Math. Soc. 2 (1970), 323–329.
- [33] A. Umamaheswaran, R. Udhayakumar, C. Selvaraj, K. Tamilvanan and M. Jima Kabeto, Existence of covers and envelopes of a left orthogonal class and its right orthogonal class of modules, J. Math. 2022, Article ID 8072134, 12 pp.
- [34] J. Z. Xu, Flat Covers of Modules, Lecture Notes in Math. 1634, Springer-Verlag, Berlin, 1996.
- [35] Y. J. Yang and X. G. Yan, Notes on Gorenstein flat modules, Bull. Iranian Math. Soc. 45 (2019), 337–344.
- [36] X. S. Zhu, Resolving resolution dimensions, Algebr. Represent. Theor. 16 (2013), 1165–1191.