

# Oscillatory Escape in a Duffing Equation with a Polynomial Potential

Mark Levi\*

*Mathematical Sciences, Rensselaer Polytechnic Institute, Troy, New York 12180*

and

Jiangong You

*Department of Mathematics, Nanjing University, Nanjing 210093, People's Republic of China*

Received March 3, 1997; revised April 8, 1997

We show that the time-periodic Hamiltonian systems  $d^2x/dt^2 + x^{2n+1} + a(t)x^{2l+1} = 0$ ,  $2n > 2l > n$ , with a discontinuity in  $a(t)$ , possess unbounded solutions  $x(t)$  which, moreover, oscillate between a finite disk and infinity; in particular  $\liminf_{t \rightarrow \infty} x(t) < \infty$  and  $\limsup_{t \rightarrow \infty} x(t) = \infty$ . As a consequence, the Poincaré map possesses no invariant KAM curves enclosing the origin outside a bounded disk. © 1997 Academic Press

## 1. BACKGROUND

*Earlier stability results.* Stability problem of Hamiltonian dynamics in its most basic but already nontrivial form arises in the study of particles in the force field of a time-periodic potential:

$$\frac{d^2x}{dt^2} + V_x(t, x) = 0, \quad V(t+1, x) = V(t, x). \quad (1)$$

A large class of such systems turns out to be near-integrable at infinity; as a consequence, “most” large-amplitude solutions are quasiperiodic with two frequencies, and furthermore, all solutions are bounded for all time. The main distinguishing feature of this class of systems is the super-quadratic behavior of the potential  $V$  for large  $|x|$ . The near-integrability at infinity is due to the difference in time scales between the time-dependence of the potential on the one hand and the rapid oscillations of high-amplitude solutions on the other. There has been much work on the

\* Research partially supported by an NSF grant.

near-integrability of (1) in various cases, starting with Morris [MO], who considered the quartic potential  $V = x^4 - p(t)x$ . Later this was extended to arbitrary polynomial potentials of even degree [DZ], [Liu], [LL], [WY] and to more general potentials [L].

It might seem surprising at the first glance that no smallness requirements are made on the time-dependence of  $V(x, t)$  which can undergo large changes during one period. For instance, all solutions of both systems

$$\ddot{x} + (2 + \sin t)x^3 = 0, \quad (2)$$

$$\ddot{x} + (2 + \sin t)\cosh x = 0 \quad (3)$$

are bounded, and in fact these systems are near-integrable for large-amplitude solutions. This near-integrability result holds for a large class of systems (1) with  $V$  satisfying the superquadraticity condition

$$\frac{VV_{xx}}{V_x^2} > \frac{1}{2} + c, \quad c > 0 \quad (4)$$

together with some extra growth assumptions on higher derivatives [L].

All stability proofs are based on the reduction to a near-integrable form with a subsequent application of Moser's twist theorem [M3].

*Smoothness and regularity.* For the simplest potential  $V = \frac{1}{4}x^4 - p(t)x$  corresponding to the system

$$\ddot{x} + x^3 = p(t) \quad (5)$$

considered by Morris [MO], stability holds if  $p(t)$  is merely piecewise continuous, or even summable. Dieckerhoff and Zehnder [DZ] proved stability for polynomial potentials with constant leading coefficients; the latter restriction was removed in [LL] with the result that any system (1) with

$$V = \sum_{k=1}^{2n} p_k(t)x^k, \quad n \geq 2 \quad (6)$$

is stable as long as all  $p_k \in C^{5+\varepsilon}$  and  $0 < p_{2n} \in C^{6+\varepsilon}$ . There have been further refinements on the degree of smoothness required of different coefficients of polynomials. By modifying proofs in [DZ] and using some approximation techniques, B. Liu ([Liu]) proved that for the system  $d^2x/dt^2 + x^{2n+1} + a(t)x = p(t)$  the mere continuity of  $a(t)$ ,  $p(t)$  implies near-integrability. Recently Wang and You [WY] proved that the smoothness can be lowered to  $C^2$  in a general polynomial potential.

*Instability results.* If the polynomial nature of the potential is destroyed, the instability can occur even if the potential remains superquadratic in  $x$ . The first such “counter-KAM” result was proven by Littlewood [Lit] in 1966 (see also [LO], [L1]), who constructed a system

$$\ddot{x} + V_x(x) = p(t) \tag{7}$$

with  $V_x/x \rightarrow \infty$  where  $p$  is piecewise continuous, with an unbounded solution. Construction of a similar counterexample with a *continuous*  $p$  is a much more delicate question; it has recently been solved by Zharnitsky [Z]. In the examples of Littlewood and Zharnitsky the potentials are not polynomial. For the polynomial potentials with their nice  $x$ -dependence this leaves open the question of the required smoothness for the coefficients.

## 2. RESULTS

In this paper we prove that in special polynomial equations

$$\frac{d^2x}{dt^2} + x^{2n+1} + a(t)x^{2l+1} = 0, \quad 2l > n, \tag{8}$$

with  $a(t)$  periodic, a jump discontinuity in  $a(t)$  causes “chaos”: no bounding KAM circles exist near infinity in the Poincaré phase plane. Moreover, we will show the existence of solutions which oscillate “chaotically” between infinity and a bounded disk.

Here, in contrast with the equation (5), the discontinuity destroys the near-integrability. The latter is thus linked with the smoothness of coefficients of higher order terms in the polynomial potential.

In this paper we consider the simplest case of a piecewise constant periodic function  $a(t) = k^{[t] \bmod 2}$ , but the proof extends to the general case without difficulty. In summary, a jump in the coefficient  $a(t)$  destroys all KAM curves bounding the origin from infinity near infinity.

### 2.1. Main Theorem

Before stating the theorem we recall that  $a(t) = k^{[t] \bmod 2}$  in the equation (8), so that  $a$  alternates periodically between the values 1 and  $k$ ; we take  $0 < k < 1$  throughout. Introduce the Hamiltonian

$$H_k = \frac{1}{2}y^2 + x^{2n+2} + kx^{2l+2}. \tag{9}$$

**THEOREM 1.** *There exist constants  $\lambda_0$  and  $\kappa > 0$  such that for any binary sequence  $\sigma = (\sigma_0\sigma_1\sigma_2\cdots)$ ,  $\sigma_j = \pm 1$ , satisfying  $\sum_{k=0}^N \sigma_k \geq 0$ ,  $\forall N \in \mathbb{Z}^+$ , there*

is an initial condition  $z = z_\sigma \in \mathbf{R}^2$  such that for the Poincaré map  $P$  of Equation (8) we have

$$H_k(P^i z_\sigma) < H_k(P^{i-1} z_\sigma) - \kappa (H_k(P^{i-1} z_\sigma))^\alpha \quad \text{if } \sigma_i = -1, \quad (10)$$

$$H_k(P^i z_\sigma) > H_k(P^{i-1} z_\sigma) + \kappa (H_k(P^{i-1} z_\sigma))^\alpha \quad \text{if } \sigma_i = 1, \quad (11)$$

where  $\alpha = (l+1)/(n+1)$ .

**COROLLARY 2.** *By different choices of the sequence we can make a point escape to infinity in infinitely many ways. For instance, we can make the iterates escape to infinity in an oscillatory fashion, so that  $\limsup |P^n z_\sigma| = \infty$ , while  $\liminf |P^n z_\sigma| < \infty$ .*

**COROLLARY 3.** *There exists an open interval  $I$  such that for any  $\mu \in I$  there exists a sequence  $\sigma$  depending on  $\mu$  such that  $|P^n z_\sigma| = O(n^\mu)$ .*

**Remark 4.** With a little extra work one can prove a stronger statement: Theorem 1 holds for any periodic  $a(t)$  which is piecewise smooth and has at least one jump (but finitely many). For the proof of this statement one has to make a change of variables in which the vectorfield is near-autonomous between the jumps of  $a$ . Such changes of variables are readily available in the papers mentioned above.

**COROLLARY 5.** *Equation (8) possesses a solution  $(x(t), \dot{x}(t))$  with  $|x(t)| + |\dot{x}(t)| \rightarrow \infty$  as  $t \rightarrow \infty$ , with  $H_k(x(t))$  growing as  $t^{1/(1-\rho)}$ , where  $0 < \rho = \rho(k, l, n) < 1$ .*

**Remark 6.** It follows from the proof that for the choice  $\sigma_i = 1$  for all  $i$  we have  $|P^n z_\sigma| \approx n^{1/(1-\kappa)}$ .

**Remark 7.** It is somewhat easier to prove that a similar result holds for the equation

$$\ddot{x} + a(t) x^{2n+1} + \dots = 0 \quad (12)$$

with the discontinuity in the highest coefficient (here  $\dots$  denote the lower order polynomial terms). In that case the growth estimates (10) and (11) of Theorem 1 are modified so that they give a geometric growth rate.

**Remark 8.** The Poincaré map of (8) is the composition of two monotone twist maps; this justifies the application of the Aubry-Mather theory with the result that Poincaré map of Equation (8) possesses a Mather set of any rotation number  $r$  in the interval  $[r_0, \infty)$  for some  $r_0$ . As a consequence of Theorem 1, none of these Mather sets with  $r > r_0$  forms an invariant curve.

2.2. An Outline of the Proof

Let  $P_k$  be the time one map of the flow with the Hamiltonian  $H_k$  given by (9). We note that the Poincaré map is the composition of two twist maps:

$$P = P_k \circ P_1. \tag{13}$$

The main steps of the proof are illustrated in Fig. 1. We start by choosing a parametrized curve  $AB$  given by  $z = z(s)$ ,  $s \in [a, b]$ , from which all the initial points mentioned in the theorem will be chosen. Referring to Fig. 1, we choose  $AB$  in the first quadrant, far enough from the origin, and  $H_k$ -monotone, in the sense of the following definition.

DEFINITION 9. We will call a smooth curve  $z(s) = (x(s), y(s))$ ,  $s \in [a, b]$   $= I$ , with  $a \leq s \leq b$ ,  $H$ -monotone, if

$$\frac{d}{ds} H(z(s)) \geq 0, \quad \frac{d}{ds} \arg z(s) < 0. \tag{14}$$

Next we consider the iterated curve  $z_1(s) = P_1 z(s)$ , Fig. 1. This curve spirals  $O(\lambda^{(2l-n)/(2n+2)})$  times around the origin for  $a \leq s \leq b$ , where  $\lambda = \min_{0 \leq s \leq \pi/2} H_k(z(s))$ . The intersection of this spiral with the second quadrant  $Q_2 = \{(x, y) : x \leq 0, y \geq 0\}$  contains  $O(\lambda^{(2l-n)/(2n+2)})$  curves. Each of these curves is  $H_1$ -monotone, as we will show later; more precisely, there exist  $O(\lambda^{(2l-n)/(2n+2)})$  intervals  $I_j \subset [a, b]$  such that each of the curves  $z_1(s)$ ,  $s \in I_j$  is  $H_1$ -monotone. In a similar way, each of these latter curves maps under  $P_k$  onto a spiral which cuts through  $Q_1$  in  $O(\lambda^{(2l-n)/(2n+2)})$   $H_k$ -monotone curves. To summarize, we get the existence of subintervals




FIG. 1. Iteration of the initial curve  $AB$  with the Poincaré map.

$I_{ij} \subset [a, b]$  such that each curve  $P_k P_1 z(s)$ ,  $s \in I_{ij}$  is  $H_k$ -monotone. This completes an induction step which produces a “chain reaction”: one  $H_k$ -monotone curve gives rise to many  $H_k$ -monotone curves; amongst these curves some are “much closer” to the origin and some are much further than the original curve, as measured by  $H_k$ : there exist indices  $i$  and  $j$  such that

$$H_k(P(z(s))) - H_k(z(s)) \geq \kappa H_k(z(s))^\alpha \quad \text{if } s \in I_{ij} \quad (15)$$

and there exist other indices  $i$  and  $j$  for which

$$H_k(P(z(s))) - H_k(z(s)) \leq -\kappa H_k(z(s))^\alpha \quad \text{if } s \in I_{ij}, \quad (16)$$

where  $\alpha = (l+1)/(n+1)$  and where  $\kappa > 0$  is independent of the initial choice of  $z(0)$ . We will prove this estimate below. According to this estimate we can control the distance to the origin of the iterate by choosing the initial  $s$  in the proper  $I_{ij}$ . This allows us to satisfy the first  $\sigma_1$  in the binary sequence. Applying the above argument to  $I_{ij}$  we can satisfy  $\sigma_2$ , and so on. The intersection of the resulting sequence of nested intervals contains the point  $s_\sigma$  whose iterates  $P^n z(s_\sigma)$  have the desired itinerary. It remains to mention that in the above argument the distance to the origin has to remain bounded from 0 by a fixed amount for every iterate; this results in the restriction  $\sum_1^N \sigma_i \geq 0$ .

### 3. PROOFS

#### 3.1. Notations and Lemmas

Let  $P_k$  be the time one map of the flow with the Hamiltonian (9), and let  $T_k(\lambda)$  be the period of the solution of the system with  $H_k$  with the energy  $H_k = \lambda$ .

LEMMA 10. *Let  $H_c = \frac{1}{2}y^2 + V_c(x)$ , with  $V_c(x) = x^{2n+2} + cx^{2l+2}$ , with  $0 < c \leq 1$ . If  $c_1 \leq c_2$ , we have*

- (1)  $H_{c_1}$  monotone curve in  $Q_1$  is  $H_{c_2}$  monotone;
- (2)  $H_{c_2}$  monotone curve in  $Q_2$  is  $H_{c_1}$  monotone.

It suffices to prove the case (1). Let  $z(s) = (x(s), y(s))$ ,  $s \in [a, b]$  be an  $H^1$ -monotone curve in  $Q_1$ . Note (14) implies that  $x_s \equiv (d/ds)x(s) > 0$ . Indeed,  $(d/ds) \arg z(s) < 0$  implies  $y_s x < x_s y$ , and the contrary assumption  $x_s < 0$  implies  $y_s < 0$ ; the last two inequalities contradict the energy estimate  $H_s = yy_s + V'x_s > 0$  of (14).

Using this fact we obtain

$$\frac{d}{ds} H_{c_2} = yy_s + V'_{c_2} x_s \geqq yy_s + V'_{c_1} x_s = \frac{d}{ds} H_{c_1} > 0. \tag{17}$$

The proof of (2) is identical.

Q.E.D.

The main theorem of this paper is an easy conclusion of the following main lemma.

LEMMA 11. *Let  $z(s)$  be an  $H_k$ -monotone curve in  $Q_1$ , with  $a \leqq s \leqq b$ . Assume, moreover, that the change in  $H_k(z(s)) \equiv h(s)$  is bounded according to*

$$h(s)|_{s=a}^{s=b} = h(b) - h(a) \leqq \frac{1}{6} h^\alpha(a), \quad \text{where } \alpha = (l+1)/(n+1). \tag{18}$$

*There exist constants  $\lambda_0 = \lambda_0(k, n)$  and  $\kappa > 0$  such that if  $h(a) \geqq \lambda_0$ , then there exist two sub-intervals  $I_1 = [a_1, b_1] \subset [a, b]$  and  $I^1 = [a^1, b^1] \subset [a, b]$  such that  $z_1(s) = P(z(s))|_{s \in I_1}$  and  $z^1(s) = P(z(s))|_{s \in I^1}$  are  $H_k$ -monotone curves in  $Q_1$ . Moreover,*

$$\text{for all } s \in I_1 \quad H_k(P(z(s))) \geqq h(b) + \kappa h^\alpha(b) \tag{19}$$

$$\text{for all } s \in I^1 \quad H_k(P(z(s))) \leqq h(a) - \kappa h^\alpha(a). \tag{20}$$

*In addition,*

$$H_k(P(z(s)))|_{s=a_1}^{s=b_1} \leqq \frac{1}{6} h^\alpha(a_1). \tag{21}$$

$$H_k(P(z(s)))|_{s=a^1}^{s=b^1} \leqq \frac{1}{6} h^\alpha(a^1). \tag{22}$$

Inductive application of this lemma according to a given binary sequence  $\sigma$  yields a sequence of nested intervals,  $I \supset I_1 \supset I_2 \supset \dots$  and a sequence of monotone curves  $z_i(s) \equiv P^i z(s)$ ,  $s \in I_i$  satisfying

$$\begin{aligned} H_k(P^i z_\sigma) &\leqq \min_{s \in I_i} (H_k(P^{i-1} z(s)) - \kappa (H_k(P^{i-1} z(s)))^\alpha) \\ &\leqq H_k(P^{i-1} z_\sigma) - \kappa (H_k(P^{i-1} z_\sigma))^\alpha \quad \text{for negative } \sigma_i, \end{aligned} \tag{23}$$

$$\begin{aligned} H_k(P^i z_\sigma) &\geqq \max_{s \in I_i} (H_k(P^{i-1} z(s)) + \kappa (H_k(P^{i-1} z(s)))^\alpha) \\ &\geqq H_k(P^{i-1} z_\sigma) + \kappa (H_k(P^{i-1} z_\sigma))^\alpha \quad \text{for positive } \sigma_i. \end{aligned} \tag{24}$$

We thus conclude that the iterates of any point  $z^* \in \bigcap_{n=1}^\infty I_n$  escape to infinity<sup>1</sup> in the way described in the Theorem. This completes the proof of the main theorem modulo last lemma.

<sup>1</sup> One can show that this point is in fact unique:  $\bigcap_{n=1}^\infty I_n = \{x^*\}$ .

In order to prove Lemma 11, we first prove:

LEMMA 12. For  $A_1$  sufficiently large, we have, for any  $\lambda > A_1$ ,

$$c_1 \lambda^{-n/(2n+2)} < T_k(\lambda) < c_2 \lambda^{-n/(2n+2)}, T'_k(\lambda) < -c_3 \lambda^{-1-n/(2n+2)}, \quad (25)$$

where  $c_1, c_2$  and  $c_3$  are positive constants depending only on  $n, k$ .

*Proof.* We omit the simple proof. Q.E.D.

LEMMA 13. Fix  $0 < c < 1$ . For any  $H_c$ -monotone curve  $z(s), s \in [a, b]$  which satisfies

$$H_c(z(b)) - H_c(z(a)) \leq \frac{1}{6} H_k(z(a))^{(l+1)/(n+1)}, \quad (26)$$

we have

$$\arg P_c z(s)|_a^b \geq \frac{1}{2} c_3 H_c(z(a))^{-(n+2)/(2n+2)} (H_c(z(b)) - H_c(z(a))) \quad (27)$$

if  $\lambda > A_1$ , where  $A_1, c_3$  are the constants in Lemma 12.

*Proof.* In view of Lemma 12, we have

$$\begin{aligned} & \arg P_c z(b) - \arg P_c z(a) \\ &= \frac{1}{T_c(H_c(z(b)))} - \frac{1}{T_c(H_c(z(a)))} \\ &\geq \frac{T_c(H_c(z(a))) - T_c(H_c(z(b)))}{T_c(H_c(z(a)))^2} \geq \frac{T_c(H_c(z(a))) - T_c(H_c(z(b)))}{T_c(H_c(z(a)))^2} \\ &= -\frac{T'_c(H_c(z(a))) + \xi(H_c(z(b)) - H_c(z(a)))}{T_c(H_c(z(a)))^2} (H_c(z(b)) - H_c(z(a))) \\ &\geq c_3 \frac{(H_c(z(a))) + \frac{1}{6} H_k(z(a))^{(l+1)/(n+1)}^{-1-n/(2n+2)}}{T_c(H_c(z(a)))^2} (H_c(z(b)) - H_c(z(a))) \\ &\geq \frac{1}{2} c_3 H_c(z(a))^{-(n+2)/(2n+2)} (H_c(z(b)) - H_c(z(a))), \end{aligned} \quad (28)$$

if  $H_c(z(a)) > A_1$ . Q.E.D.

*Proof of Lemma 11.* Let  $z(s)$  be an  $H_k$ -monotone curve in the first quadrant which satisfies the assumptions in Lemma 11. That is, denoting

$$H_k(z(a)) = \lambda, \quad (29)$$

we assume

$$H_k(z(b)) = \lambda_1 \leq \lambda + \frac{1}{6} \lambda^\alpha, \tag{30}$$

and  $z(a) = (0, y(a))$ ,  $z(b) = (x(b), 0)$  with  $\lambda > \lambda_0$ ; later we will specify  $\lambda_0$ .

In the following,  $z(a)$ ,  $z(b)$ ,  $z(a')$ ,  $z(b')$ ,  $z(a_1)$ ,  $z(b_1)$ ,  $z(a_+)$  coincide with  $A$ ,  $B$ ,  $A'$ ,  $B'$ ,  $A_1$ ,  $B_1$ ,  $A_+$  in the figures respectively.

Following the outline of the proof above, we first prove that the image of  $z(s)$  under the map  $P_1$  spirals many times around the origin so that we choose the higher  $H_k$ -energy part of it in the second quadrant. In view of Lemma 13, we only need to prove the energy difference between the two end points is sufficiently large.

We first estimate the  $H_1$  energy difference of  $z(a)$  and  $z(b)$ . Note that  $H_k(z(a)) \leq H_k(z(b))$  by the  $H_k$ -monotonicity of  $z(s)$ , and

$$H_1(z(a)) = H_k(z(a)) \leq H_k(z(b)) = x^{2n+2}(b) + kx^{2l+2}(b), \tag{31}$$

$$H_1(z(b)) = x^{2n+2}(b) + x^{2l+2}(b) = H_k(z(b)) + (1 - k) x(b)^{2l+2}, \tag{32}$$

$$x(b) \geq (\frac{1}{2})^{1/(2n+2)} \lambda^{1/(2n+2)} \tag{33}$$

if  $\lambda > 2$ . It follows that for  $\lambda \geq 2$ ,

$$\begin{aligned} H_1(z(b)) - H_1(z(a)) &\geq H_k(z(b)) + (1 - k) x(b)^{2l+2} - H_1(z(a)) \\ &= H_k(z(b)) - H_k(z(a)) + (1 - k) x(b)^{2l+2} \\ &\geq (1 - k) x(b)^{2l+2} \geq \frac{1}{2} (1 - k) \lambda^\alpha. \end{aligned} \tag{34}$$

In view of Lemma 13, for  $\lambda \geq A_1$  we have

$$\begin{aligned} \arg P_1 z(s)|_a^b &\geq \frac{1}{2} (1 - k) \lambda^{(l+1)/(n+1)} \cdot \frac{1}{2} c_3 \lambda^{-(n+2)/(2n+2)} \\ &= \frac{1}{4} (1 - k) c_3 \lambda^{(2l-n)/(2n+2)}. \end{aligned} \tag{35}$$

If  $2l > n$ ,  $\lambda > A_2 = \max\{A_1, (96\pi/(1 - k) c_3)^{(2n+2)/(2l-n)}\}$ , then there exists  $a_+ \in [a, b]$  such that  $\arg P_1 z(s)|_{a_+}^b = 4\pi$  and

$$H_1(z(s))|_{a_+}^b \leq \frac{1}{12} (1 - k) \lambda^\alpha. \tag{36}$$

Note that  $z(s)$  is  $H_1$ -monotone in the first quadrant since it is  $H_k$ -monotone, and  $P_1$  is a energy preserving map implies

$$\frac{d}{ds} H_1(P_1 z(s)) = \frac{d}{ds} H_1(z(s)) \geq 0.$$

By  $H_1$  monotonicity of  $z(s)$ ,  $H_1(z(s')) > H_1(z(s))$  if  $s' > s$ . By the monotone twist estimate of Lemma 12, which states that the point with large energy rotates clockwise further than the point with small energy, it follows that

$$\arg H_1(z(s')) < \arg H_1(z(s)), \tag{37}$$

if  $s' > s$ , i.e.,  $(d/ds) \arg H_1(z(s)) \geq 0$ . By the definition of the  $H_k$ -monotonicity,  $P_1 z(s)$  is  $H_k$ -monotone.

We assure the existence of a subinterval  $I' = [a', b'] \subset [a_+, b]$  such that  $z_1(s) = P_1 z(s)$ ,  $s \in I'$  is  $H_1$ -monotone in the second quadrant  $Q_2$ :

$$z_1(I') = P_1 z(I') \subset Q_2, x_1(a') < 0, y_1(a') = 0; x_1(b') = 0, y_1(b') > 0, \tag{38}$$

and

$$\frac{d}{ds} H_1(z_1(s)) \geq 0, \quad \frac{d}{ds} \arg z_1(s) < 0. \tag{39}$$

From (34) and (36) we obtain, using monotonicity, for  $s \in [a', b']$  the following estimate on the energy gain (see (34)):

$$\begin{aligned} H_1(z(s)) &\geq H_1(z(a_+)) = H_1(z(a)) + H_1(z(s))|_a^b - H_1(z(s))|_a^b \\ &\geq \lambda + \frac{1}{2}(1-k) \lambda^\alpha - \frac{1}{12}(1-k) \lambda^\alpha = \lambda + \frac{5}{12} \lambda^\alpha, \end{aligned} \tag{40}$$

if  $\lambda > \max\{A_2, 2\}$ . To prove that  $P_k z_1(s)$ ,  $s \in I'$  spirals many times around the origin, we observe that  $z_1(s)$ ,  $s \in I'$  climbs steeply with respect to  $P_k$ , see Fig. 1. To be precise,

$$\begin{aligned} H_k(z_1(s))|_{a'}^{b'} &= \frac{y_1^2(b')}{2} - x_1^{2n+2}(a') - kx_1^{2l+2}(a') \\ &= H_1(z_1(b')) - H_1(z_1(a')) + (1-k) x_1^{2l+2}(a') \\ &= H_1(z_1(s))|_{a'}^{b'} + (1-k) x_1^{2l+2}(a') \geq (1-k) x_1^{2l+2}(a') \end{aligned} \tag{41}$$

From  $H_1(z_1(a')) = x_1^{2n+2}(a') + x_1^{2l+2}(a') \geq \lambda$  (see (29)), it follows that  $x_1(a') \geq (\frac{1}{2})^{1/(2n+2)} \lambda^{1/(2n+2)}$  if  $\lambda > 2$ . Thus

$$H_k(z_1(s))|_{a'}^{b'} \geq \frac{1}{2}(1-k) \lambda^\alpha. \tag{42}$$

It follows from Lemma 13 that

$$\begin{aligned} \arg P_k z_1(s)|_{a'}^{b'} &\geq \frac{1}{4}c_3(1-k) \lambda^\alpha H_k(z_1(a'))^{-(n+2)/(2n+2)} \\ &\geq \frac{1}{8}c_3(1-k) \lambda^\alpha H_1(z_1(a'))^{-(n+2)/(2n+2)} \\ &= \frac{1}{8}c_3(1-k) \lambda^\alpha H_1(z(a'))^{-(n+2)/(2n+2)} \\ &\geq \frac{1}{8}(1-k) c_3 \lambda^{(2l-n)/(2n+2)} \end{aligned} \tag{43}$$

if  $\lambda > 2$ .

As before, if  $\lambda \geq 2A_2$  we can choose the higher-energy subinterval of  $I'$  by letting  $a'_+ \in (a', b')$  such that  $\arg P_k z_1(s)|_{a'_+}^{b'} = 4\pi$  and

$$H_k(z_1(s))|_{a'_+}^{b'} \leq \frac{1}{12}(1-k) \lambda^\alpha. \tag{44}$$

Thus there exists a subinterval  $I_1 = [a_1, b_1] \subset [a'_+, b']$  for which  $P_k z_1(s)$  is an  $H_k$ -montone curve in the first quadrant; moreover, for  $s \in I_1$ , using monotonicity, we get

$$\begin{aligned} H_k(Pz(s)) &= H_k(P_k z_1(s)) = H_k(z_1(s)) \\ &\geq H_k(z_1(a'_+)) = H_k(z(b')) - H_k(z_1(s))|_{a'_+}^{b'} \\ &\geq H_1(z_1(b')) - \frac{1}{12}(1-k) \lambda^\alpha \\ &\geq \lambda + \frac{5}{12}(1-k) \lambda^\alpha - \frac{1}{12} \lambda^\alpha = \lambda + \frac{1}{3} \lambda^\alpha. \end{aligned} \tag{45}$$

Together with  $\lambda_1 - \lambda \leq \frac{1}{6} \lambda^\alpha$ , it implies that for  $\lambda > 2$

$$H_k(Pz(s)) \geq \lambda + \frac{1}{3} \lambda^\alpha \geq \lambda_1 + \frac{1}{6} \lambda^\alpha \geq \lambda_1 + \frac{1}{7} \lambda_1^\alpha, \tag{46}$$

as long as  $s \in [a_1, b_1]$ . Meanwhile,  $H_k(Pz(s))|_{a_1}^{b_1} \leq H_k(z_1(s))|_{a'_+}^{b'} \leq \frac{1}{6}(1-k) \lambda^{(l+1)/(n+1)} \leq \frac{1}{6} H_k(Pz(a_1))$  if  $H_k(z(a)) \geq \max\{2, 2A_2\}$ .

Similarly, we can prove the existence of  $I^1$ . We omit the virtually identical details. Q.E.D.

### ACKNOWLEDGMENTS

We thank Forschungsinsitut für mathematik at ETH Zurich, and in particular Professor Jürgen Moser for hospitality.

## REFERENCES

- [A] V. I. Arnold, Proof of A. N. Kolmogorov's theorem on the preservation of quasiperiodic motions under small perturbations of the Hamiltonian, *Usp. Mat. Nauk. SSSR* **18**, No. 5 (1963), 13–40.
- [DZ] R. Dieckerhoff and E. Zehnder, Boundedness of solutions via the twist theorem, *Ann. Scuola. Sup. Pisa Cl. Sci.* **14** (1987), 79–95.
- [GH] J. Guckenheimer and P. Holmes, “Nonlinear Oscillations, Dynamical System and Bifurcations of Vector Fields,” Springer-Verlag, New York, 1983.
- [LL] S. Laederich and M. Levi, Invariant curves and time-dependent potentials, *Ergod. Th. Dynam. Sys.* **11** (1991), 365–378.
- [L] M. Levi, Quasiperiodic motions in superquadratic time-dependent potentials, *Commun. Math. Phys.* **143** (1991), 43–83.
- [L1] M. Levi, On Littlewood's counterexample of unbounded motions in superquadratic potentials, *Dynam. Report (N.S.)* **1** (1997), 113–124.
- [Lit] J. E. Littlewood, Unbounded solutions of an equation  $\ddot{y} + g(y) = p(t)$ , with  $p(t)$  periodic and bounded, and  $g(y)/y \rightarrow \infty$  as  $y \rightarrow \pm\infty$ , *J. Lond. Math. Soc.* **41** (1966), 497–507.
- [Liu] B. Liu, Boundedness for solutions of nonlinear periodic differential equations via Moser's twist theorem, *Acta Math. Sinica (N.S.)* **8** (1992), 91–98.
- [LO] Y. Long, An unbounded solution of a superlinear Duffing's equation, *Acta Math. Sinica (N.S.)* **7**, No. 4 (1991), 360–369.
- [MO] G. Morris, A case of boundedness in Littlewood's problem on oscillatory differential equations, *Bull. Austral. Math. Soc.* **14** (1976), 71–93.
- [M1] J. Moser, “Stable and Random Motions in Dynamical Systems,” *Ann. of Math. Studies* **77**, Princeton Univ. Press, Princeton, NJ, 1973.
- [M2] J. Moser, On invariant curves of area-preserving mappings of annulus, *Nachr. Akad. Wiss. Gottingen Math. Phys. Kl. (II)* (1962), 1–20.
- [WY] Y. Wang and J. You, Boundedness of solutions for polynomial potentials with  $C^2$  time dependent coefficients, *ZAMP* **47** (1996).
- [Z] V. Zharnitsky, Breakdown of stability of motion in superquadratic potentials. Ph.D. thesis, RPI, 1996. To appear in *Comm. Math. Phys.*