

# Homogenization of stationary Navier-Stokes equations in domains with tiny holes

Eduard Feireisl <sup>\*</sup>      Yong Lu <sup>†</sup>

Institute of Mathematics of the Academy of Sciences of the Czech Republic  
Žitná 25, 115 67 Praha 1, Czech Republic  
and

Charles University in Prague, Faculty of Mathematics and Physics, Mathematical Institute  
Sokolovská 83, 186 75 Praha 8, Czech Republic

## Abstract

We consider the homogenization problem for the stationary compressible Navier-Stokes equations describing a steady flow of a compressible Newtonian fluid in a bounded three dimensional domain. We focus on the case where the domain is perforated with very tiny holes for which the diameters are much smaller than their mutual distances. We show that the homogenization process does not change the motion of the fluids: In the asymptotic limit, we obtain again the same system of equations. This coincides with similar results for the stationary Stokes and stationary *incompressible* Navier-Stokes system.

**Key words:**

## 1 Introduction

Homogenization problems in the framework of fluid mechanics have gain a lot of interest. For Stokes and stationary *incompressible* Navier-Stokes equations, Allaire in [1, 2] gave a system study

---

<sup>\*</sup>Eduard Feireisl acknowledges the support of the project LL1202 in the programme ERC-CZ funded by the Ministry of Education, Youth and Sports of the Czech Republic.

<sup>†</sup>Yong Lu acknowledges the support of the project LL1202 in the programme ERC-CZ funded by the Ministry of Education, Youth and Sports of the Czech Republic.

for physical domains perforated by a family of holes of different size. More specifically, consider a system of holes of diameter  $O(\varepsilon^\alpha)$ , where  $\varepsilon$  is their mutual distance. In three spatial dimensions, Allaire showed that when  $\alpha < 3$ , the limit fluid behavior is governed by the classical Darcy's law; when  $\alpha = 3$ , in the limit yields Brinkman law; when  $\alpha > 3$ , the equations do not change in the homogenization process and the limit problem is determined by the same system of Stokes or Navier-Stokes equations. In the case  $\alpha = 1$ , meaning that the size of holes is proportional to their mutual distance, the results have been extended to the incompressible evolutionary Navier-Stokes equations by Mikelić [14], and to the compressible Navier-Stokes system by Masmoudi [13], and to the complete Navier-Stokes-Fourier system in [9]. In all the aforementioned cases, the asymptotic limit gives rise to Darcy's law.

In this paper, we focus on the homogenization process for the stationary compressible Navier-Stokes equations in a domain perforated by very tiny holes, where the diameter of the holes is taken to be of size  $O(\varepsilon^\alpha)$ , with  $\alpha > 3$ , where  $\varepsilon$  denotes the mutual distance between the holes. We start by a precise description of the physical domain. We consider a bounded domain  $\Omega$  in  $R^3$  of class  $C^2$ , and a family of holes (solid obstacles)  $\{T_{\varepsilon,k}^s\}$ , which are simply connected smooth domains satisfying

$$T_{\varepsilon,k}^s \subset \overline{T}_{\varepsilon,k}^s \subset B_{\varepsilon,k} \subset \overline{B}_{\varepsilon,k} \subset \varepsilon C_k \quad (1.1)$$

with

$$C_k := (0,1)^3 + k, \quad k \in Z^3, \quad B_{\varepsilon,k} := B(x_k, b_0 \varepsilon^\alpha) \text{ for some } x_k \in T_{\varepsilon,k}^s, \quad b_0 > 0. \quad (1.2)$$

For the sake of simplicity, we suppose that all holes  $T_{\varepsilon,k}^s = \varepsilon^\alpha T^s$  are similar to the same set  $T^s$  - a simply connected domain of class  $C^2$ .

The corresponding family of  $\varepsilon$ -dependent perforated domains is defined as

$$\Omega_\varepsilon := \Omega \setminus \bigcup_{k \in K_\varepsilon} \overline{T}_{\varepsilon,k}^s, \quad K_\varepsilon := \{k \mid \varepsilon \overline{C}_k \subset \Omega\}. \quad (1.3)$$

It is easy to check that the number of holes contained in  $\Omega$  satisfies

$$|K_\varepsilon| = \frac{|\Omega|}{\varepsilon^3} (1 + o(1)), \quad \text{as } \varepsilon \rightarrow 0. \quad (1.4)$$

We consider the following stationary Navier-Stokes equations in  $\Omega_\varepsilon$ :

$$\operatorname{div}_x(\varrho \mathbf{u}) = 0, \quad (1.5)$$

$$\operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p(\varrho) = \operatorname{div}_x \mathbb{S}(\nabla_x \mathbf{u}) + \varrho \mathbf{f} + \mathbf{g}, \quad (1.6)$$

$$\mathbb{S}(\nabla_x \mathbf{u}) = \mu \left( \nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{3} \operatorname{div}_x \mathbf{u} \mathbb{I} \right) + \eta \operatorname{div}_x \mathbf{u} \mathbb{I}, \quad \mu > 0, \quad \eta \geq 0. \quad (1.7)$$

Here,  $\varrho$  is the fluid mass density,  $\mathbf{u}$  is the velocity field,  $p = p(\varrho)$  denotes the pressure,  $\mathbb{S} = \mathbb{S}(\nabla_x \mathbf{u})$  stands for the Newtonian viscous stress tensor, and  $\mu, \lambda$  are the viscosity coefficients.

The density is nonnegative and the total mass of the fluid is fixed to be

$$\int_{\Omega_\varepsilon} \varrho \, dx = M > 0. \quad (1.8)$$

In addition, we impose the no-slip boundary condition

$$\mathbf{u} = 0 \quad \text{on } \partial\Omega_\varepsilon. \quad (1.9)$$

The fluid is driven by external forces here represented by the functions  $\mathbf{f}$  and  $\mathbf{g}$  satisfying

$$\|\mathbf{f}\|_{L^\infty(R^3; R^3)} + \|\mathbf{g}\|_{L^\infty(R^3; R^3)} \leq c. \quad (1.10)$$

We consider the pressure law of a typical form

$$p(\varrho) = a\varrho^\gamma, \quad a > 0, \quad (1.11)$$

with the adiabatic exponent  $\gamma \geq 1$ . The range of  $\gamma$  we can handle by the homogenization technique will be specified below.

For a function  $f$  defined in  $\Omega_\varepsilon$ , we use the symbol  $\tilde{f}$  to denote the zero-extension of  $f$  in  $R^3$ , that is

$$\tilde{f} = f \quad \text{in } \Omega_\varepsilon, \quad \tilde{f} = 0 \quad \text{in } R^3 \setminus \Omega_\varepsilon. \quad (1.12)$$

We also use  $c$  to a generic positive constant independent of the parameter  $\varepsilon$ . However, the specific value of  $c$  could change from line to line.

## 1.1 Weak solutions

We start by introducing the standard concept of *weak solution* to the compressible Navier-Stokes system.

**Definition 1.1** *We say that  $[\varrho, \mathbf{u}]$  is a finite energy weak solution of the Navier-Stokes equations (1.5 - 1.7) supplemented with the conditions (1.8 - 1.9) in the domain  $\Omega_\varepsilon$  if:*

$$\varrho \geq 0 \quad \text{a.e. in } \Omega_\varepsilon, \quad \int_{\Omega_\varepsilon} \varrho \, dx = M, \quad \varrho \in L^{\beta(\gamma)}(\Omega_\varepsilon), \quad \text{for some } \gamma \leq \beta(\gamma), \quad \mathbf{u} \in W_0^{1,2}(\Omega_\varepsilon; R^3); \quad (1.13)$$

$$\int_{\Omega_\varepsilon} \varrho \mathbf{u} \cdot \nabla_x \psi \, dx = 0, \quad (1.14)$$

$$\int_{\Omega_\varepsilon} \varrho \mathbf{u} \otimes \mathbf{u} : \nabla_x \varphi + p(\varrho) \operatorname{div}_x \varphi - \mathbb{S}(\nabla_x \mathbf{u}) : \nabla_x \varphi + (\varrho \mathbf{f} + \mathbf{g}) \cdot \varphi \, dx = 0, \quad (1.15)$$

for any test function  $\psi \in C_c^\infty(\Omega_\varepsilon)$  and any test function  $\varphi \in C_c^\infty(\Omega_\varepsilon; R^3)$ . Moreover, there holds the energy inequality:

$$\int_{\Omega_\varepsilon} \mathbb{S}(\nabla_x \mathbf{u}) : \nabla_x \mathbf{u} \, dx \leq \int_{\Omega_\varepsilon} (\varrho \mathbf{f} + \mathbf{g}) \cdot \mathbf{u} \, dx. \quad (1.16)$$

Furthermore, we introduce *renormalized weak solutions*.

**Definition 1.2** We say a finite energy weak solution  $[\varrho, \mathbf{u}]$  is a renormalized weak solution if its zero extension  $[\tilde{\varrho}, \tilde{\mathbf{u}}]$  (see the notation (1.12)) satisfies

$$\operatorname{div}_x(\tilde{\varrho} \tilde{\mathbf{u}}) = 0, \quad \operatorname{div}_x(b(\tilde{\varrho}) \tilde{\mathbf{u}}) + (\tilde{\varrho} b'(\tilde{\varrho}) - b(\tilde{\varrho})) \operatorname{div}_x \tilde{\mathbf{u}} = 0, \quad \text{in } \mathcal{D}'(R^3), \quad (1.17)$$

for any  $b \in C^0([0, \infty)) \cap C^1((0, \infty))$  such that

$$b'(s) \leq c s^{-\lambda_0} \text{ for } s \in (0, 1], \quad b'(s) \leq c s^{\lambda_1} \text{ for } s \in [1, \infty), \quad (1.18)$$

with

$$c > 0, \quad \lambda_0 < 1, \quad -1 < \lambda_1 \leq \frac{\beta(\gamma)}{2} - 1. \quad (1.19)$$

**Remark 1.1** The existence of finite energy renormalized weak solutions to the Navier-Stokes equations (1.5–1.7) for fixed  $\varepsilon > 0$  is known for certain range of the adiabatic exponent  $\gamma$ . The first global result has been obtained by Lions [12] for the case  $\gamma > 5/3$ . Extensions to lower values of  $\gamma$  were obtained by Březina, Novotný [4], Frehse et al. [10], and Plotnikov, Sokolowski [16]. Moreover, as shown in Theorem 4.3 in the monograph of Novotný, Straškraba [15], any finite energy weak solution  $[\varrho, \mathbf{u}]$  satisfies

$$\varrho \in L^{\beta(\gamma)}, \quad \beta(\gamma) = 3(\gamma - 1) \text{ if } 3/2 < \gamma < 3, \quad \beta(\gamma) = 2\gamma \text{ if } \gamma \geq 3. \quad (1.20)$$

**Remark 1.2** In view of DiPerna-Lions's transport theory (see Section II.3 in [5] and the improvement in Lemma 3.3 in [15]), for any  $r \in L^\beta(\Omega)$ ,  $\beta \geq 2$ , and any  $\mathbf{v} \in W_0^{1,2}(\Omega)$ , where  $\Omega \subset R^3$  is a bounded  $C^2$  domain, a couple of functions  $[r, \mathbf{v}]$  satisfying

$$\operatorname{div}_x(r \mathbf{v}) = 0 \quad \text{in } \mathcal{D}'(\Omega),$$

satisfies also the renormalized equations

$$\operatorname{div}_x(r \mathbf{v}) = 0, \quad \operatorname{div}_x(b(r) \mathbf{v}) + (r b'(r) - b(r)) \operatorname{div}_x \mathbf{v} = 0, \quad \text{in } \mathcal{D}'(R^3),$$

for any  $b$  satisfying (1.18 - 1.19) provided  $r$  and  $\mathbf{v}$  have been extended to be zero outside  $\Omega$ .

Hence, if  $\gamma \geq 5/3$ , any finite energy weak solution in the sense of Definition 1.1 is also a renormalized weak solution in the sense of Definition 1.2. The condition  $\gamma \geq 5/3$  ensures that  $\varrho \in L^2(\Omega_\varepsilon)$  by (1.20).

## 1.2 Main result

In this paper, we consider the case, where the adiabatic exponent  $\gamma$  in the pressure law (1.11) satisfies

$$\gamma \geq 3. \quad (1.21)$$

Note that the same restriction has been imposed by Masmoudi [13] to avoid certain so far unsurmountable difficulties connected with the compressible Navier-Stokes system.

Our main result is the following:

**Theorem 1.1** *Let*

$$M > 0, \gamma \geq 3$$

*be given. Let  $[\varrho_\varepsilon, \mathbf{u}_\varepsilon]_{0 < \varepsilon < 1}$  be a family of finite energy renormalized weak solutions to the compressible Navier-Stokes system (1.5–1.9) in  $\Omega_\varepsilon$ , where  $\mathbf{f}, \mathbf{g}$  obey (1.10). Suppose that the size parameter  $\alpha$  of the holes in (1.2) satisfies  $\alpha > 3$ .*

*Then*

$$\sup_{0 < \varepsilon < 1} \left\{ \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)} + \|\mathbf{u}_\varepsilon\|_{W_0^{1,2}(\Omega_\varepsilon)} \right\} < c, \quad (1.22)$$

*and, up to a subsequence, the extensions  $[\tilde{\varrho}_\varepsilon, \tilde{\mathbf{u}}_\varepsilon]$  satisfy*

$$\tilde{\varrho}_\varepsilon \rightarrow \varrho \text{ weakly in } L^{\beta(\gamma)}(\Omega), \quad \tilde{\mathbf{u}}_\varepsilon \rightarrow \mathbf{u} \text{ weakly in } W_0^{1,2}(\Omega), \quad (1.23)$$

*where the limit  $[\varrho, \mathbf{u}]$  is a finite energy renormalized weak solution to the same problem (1.5–1.9) in the limit domain  $\Omega$ .*

We find that the limit equations are the same as the original ones. This means that the homogenization process does not change the motion of the fluid when the holes (obstacles) are very small. The rest of the paper is devoted to the proof of Theorem 1.1. Although the result is formally the same as for the incompressible case studied by Allaire [1], [2], the technique for the compressible system is rather different. Moreover, in contrast with the critical case studied by Masmoudi [13], we do not perform any extra solution scaling and must establish the necessary bounds by means of the so-called Bogovskii's operator, the norm of which must be *independent* of the parameter  $\varepsilon$ . The construction of such an operator carried over in Section 2 represents the main step of the proof. The paper is finished by a short discussion in Section 3.

## 2 Proof of Theorem 1.1

This section is devoted to the proof of Theorem 1.1.

### 2.1 Bogovskii's operator

We introduce a suitable inverse of the divergence, commonly known as Bogovskii's operator, see Bogovskii [3], Galdi [11].

**Lemma 2.1** *Let  $\Omega_\varepsilon$  be defined through (1.1)–(1.3) with  $\alpha \geq 3$ .*

*Then there exists a linear operator  $\mathcal{B}_\varepsilon : L^2(\Omega_\varepsilon)/R \rightarrow W_0^{1,2}(\Omega_\varepsilon; R^3)$  such that for any  $f \in L^2(\Omega_\varepsilon)$  and  $\int_{\Omega_\varepsilon} f \, dx = 0$ , there holds*

$$\operatorname{div}_x(\mathcal{B}_\varepsilon(f)) = f \text{ in } \Omega_\varepsilon, \quad \|\mathcal{B}_\varepsilon(f)\|_{W_0^{1,2}(\Omega_\varepsilon; R^3)} \leq c \|f\|_{L^2(\Omega_\varepsilon)} \quad (2.1)$$

*for some  $c$  independent of  $\varepsilon$ .*

The existence of such a linear operator is nowadays standard, see e.g. Novotný, Straškraba [15, Chapter 3]. The key point here is to show the uniform estimate (2.1) with the constant  $c$  independent of  $\varepsilon$ . To this end, we adapt the construction of the *restriction operator*  $R_\varepsilon$  by Allaire [1]. We start with the following result [1, Section 2.2]:

**Lemma 2.2** *For  $\Omega_\varepsilon$  as in (1.1)–(1.3) with  $\alpha \geq 3$ , there exists a linear bounded operator  $R_\varepsilon$  mapping  $W_0^{1,2}(\Omega; R^3)$  to  $W_0^{1,2}(\Omega_\varepsilon; R^3)$  such that*

$$\mathbf{u} \in W_0^{1,2}(\Omega_\varepsilon; R^3) \implies R_\varepsilon(\tilde{\mathbf{u}}) = \mathbf{u} \text{ in } \Omega_\varepsilon, \quad (2.2)$$

$$\operatorname{div}_x \mathbf{u} = 0 \text{ in } \Omega \implies \operatorname{div}_x R_\varepsilon(\mathbf{u}) = 0 \text{ in } \Omega_\varepsilon, \quad (2.3)$$

$$\|R_\varepsilon(\mathbf{u})\|_{W_0^{1,2}(\Omega_\varepsilon; R^3)} \leq c \|\mathbf{u}\|_{W_0^{1,2}(\Omega; R^3)}, \quad c \text{ independent of } \varepsilon. \quad (2.4)$$

Inspecting the proof of Lemma 2.2 in Section 2.2 of [1], we observe that the restriction operator may be constructed in the following way:

Let  $b_1 > 0$  be chosen such that

$$B(x_k, b_1 \varepsilon) \subset \varepsilon C_k, \quad \overline{B}_{\varepsilon,k} = \overline{B(x_k, b_0 \varepsilon^\alpha)} \subset B(x_k, b_1 \varepsilon).$$

Consider the following decomposition of each cube  $\varepsilon C_k$  with  $k \in K_\varepsilon$ :

$$\varepsilon \overline{C}_k = T_{\varepsilon,k}^s \cup \overline{E}_{\varepsilon,k} \cup \overline{F}_{\varepsilon,k} \text{ with } E_{\varepsilon,k} := B(x_k, b_1 \varepsilon) \setminus T_{\varepsilon,k}^s, \quad F_{\varepsilon,k} := (\varepsilon C_k) \setminus B(x_k, b_1 \varepsilon).$$

For any  $\mathbf{u} \in W_0^{1,2}(\Omega; R^3)$ , we define  $R_\varepsilon$  through

$$R_\varepsilon(\mathbf{u}) = \mathbf{u} \text{ on } \varepsilon C_k \cap \Omega, \quad \text{for } k \notin K_\varepsilon, \quad (2.5)$$

and for  $k \in K_\varepsilon$ :

$$R_\varepsilon(\mathbf{u}) = \mathbf{u} \text{ on } F_{\varepsilon,k}, \quad R_\varepsilon(\mathbf{u}) = 0 \text{ on } T_{\varepsilon,k}^s, \quad R_\varepsilon(\mathbf{u}) = \mathbf{v}_{\varepsilon,k} \text{ in } E_{\varepsilon,k}, \quad (2.6)$$

where  $\mathbf{v}_{\varepsilon,k} \in W^{1,2}(\varepsilon C_k; R^3)$  solves the following Stokes problem

$$\nabla p_{\varepsilon,k} - \Delta \mathbf{v}_{\varepsilon,k} = -\Delta \mathbf{u} \quad \text{in } E_{\varepsilon,k}, \quad (2.7)$$

$$\operatorname{div}_x \mathbf{v}_{\varepsilon,k} = \operatorname{div}_x \mathbf{u} + \frac{1}{|E_{\varepsilon,k}|} \int_{T_{\varepsilon,k}^s} \operatorname{div}_x \mathbf{u} \, dx \quad \text{in } E_{\varepsilon,k}, \quad (2.8)$$

$$\mathbf{v}_{\varepsilon,k} = \mathbf{u} \text{ on } \partial E_{\varepsilon,k} - \partial T_{\varepsilon,k}^s, \quad \mathbf{v}_{\varepsilon,k} = 0 \text{ on } \partial T_{\varepsilon,k}^s. \quad (2.9)$$

The operator  $R_\varepsilon$  defined by (2.5–2.9) is a restriction operator satisfying (2.2–2.4).

Now we use Lemma 2.2, along with the properties of the restriction operator stated above, to prove Lemma 2.1.

**Proof of Lemma 2.1.** For  $f \in L^2(\Omega_\varepsilon)$  with  $\int_{\Omega_\varepsilon} f \, dx = 0$ , we consider the extension

$$\tilde{f} = f \text{ in } \Omega_\varepsilon, \quad \tilde{f} = 0 \text{ on } \Omega \setminus \Omega_\varepsilon = \bigcup_{k \in K_\varepsilon} T_{\varepsilon,k}^s. \quad (2.10)$$

Then, by employing the classical Bogovskii's operator defined on the domain  $\Omega$ , we obtain  $\mathbf{u} := \mathcal{B}(f) \in W_0^{1,2}(\Omega; R^3)$  such that

$$\operatorname{div}_x \mathbf{u} = \tilde{f} \text{ in } \Omega \quad \text{and} \quad \|\mathbf{u}\|_{W_0^{1,2}(\Omega; R^3)} \leq c \|\tilde{f}\|_{L^2(\Omega)} = c \|f\|_{L^2(\Omega_\varepsilon)} \quad (2.11)$$

for some  $c$  that only depends on  $\Omega$ . Moreover, by virtue of (2.10), we have

$$\operatorname{div}_x \mathbf{u} = \tilde{f} = 0 \quad \text{on } T_{\varepsilon,k}^s.$$

Let  $R_\varepsilon$  be the restriction operator constructed through (2.5 – 2.9).

In particular, it is easy to check that equation (2.8) gives rise to

$$\operatorname{div}_x \mathbf{v}_{\varepsilon,k} = \operatorname{div}_x \mathbf{u} = f \quad \text{in } E_{\varepsilon,k}, \quad (2.12)$$

whenever  $\mathbf{u}$  satisfies (2.11). In addition, one has  $R_\varepsilon(\mathbf{u}) = \mathbf{u}$  in  $\Omega_\varepsilon \setminus (\bigcup_{k \in K_\varepsilon} E_{\varepsilon,k})$ . Together with (2.6) and (2.12), we therefore conclude that

$$\operatorname{div}_x R_\varepsilon(u) = f \quad \text{in } \Omega_\varepsilon.$$

Thus to prove Lemma 2.1, it is enough to define

$$\mathcal{B}_\varepsilon(f) := R_\varepsilon(\mathbf{u}) = R_\varepsilon(\mathcal{B}(\tilde{f})),$$

where  $\mathcal{B}$  is the classical Bogovskii's operator on  $\Omega$ . The operator norm estimate (2.1) follows from the estimate for the restriction operator in (2.4).

## 2.2 Uniform bounds

Our goal in this section is to show the uniform estimate (1.22) under the assumption  $\gamma \geq 3$ . Note that, in accordance with what we said in Remark 1.1, we already know that

$$\varrho_\varepsilon \in L^{2\gamma}(\Omega_\varepsilon), \quad \mathbf{u}_\varepsilon \in W_0^{1,2}(\Omega_\varepsilon) \quad (2.13)$$

for any *fixed*  $\varepsilon$ . However, we have to establish *uniform bounds* on the norms  $\|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}$  and  $\|\mathbf{u}_\varepsilon\|_{W_0^{1,2}(\Omega_\varepsilon)}$  independent of  $\varepsilon \in (0, 1)$ .

By the energy inequality (1.16), together with Korn's inequality and Hölder's inequality, we have

$$\mu \|\nabla_x \mathbf{u}_\varepsilon\|_{L^2(\Omega_\varepsilon)}^2 \leq \|\mathbf{f}\|_{L^\infty(\Omega_\varepsilon)} \|\varrho_\varepsilon\|_{L^{\frac{6}{5}}(\Omega_\varepsilon)} \|\mathbf{u}_\varepsilon\|_{L^6(\Omega_\varepsilon)} + \|\mathbf{g}\|_{L^\infty(\Omega_\varepsilon)} \|\mathbf{u}_\varepsilon\|_{L^1(\Omega_\varepsilon)}. \quad (2.14)$$

Since  $\mathbf{u}_\varepsilon \in W_0^{1,2}(\Omega_\varepsilon)$  has zero trace on the boundary, we may use Poincaré's inequality and the Sobolev embedding to obtain

$$\|\mathbf{u}_\varepsilon\|_{L^6(\Omega_\varepsilon)} \leq c \|\nabla_x \mathbf{u}_\varepsilon\|_{L^2(\Omega_\varepsilon)} \quad (2.15)$$

for some constant  $c$  independent of the domain  $\Omega_\varepsilon$ . Thus we obtain

$$\|\nabla_x \mathbf{u}_\varepsilon\|_{L^2(\Omega_\varepsilon)} + \|\mathbf{u}_\varepsilon\|_{L^6(\Omega_\varepsilon)} \leq c \left( \|\mathbf{f}\|_{L^\infty(\Omega_\varepsilon)} \|\varrho_\varepsilon\|_{L^{\frac{6}{5}}(\Omega_\varepsilon)} + \|\mathbf{g}\|_{L^\infty(\Omega_\varepsilon)} \right) \leq c \left( \|\varrho_\varepsilon\|_{L^{\frac{6}{5}}(\Omega_\varepsilon)} + 1 \right). \quad (2.16)$$

Next, we introduce a test function

$$\varphi(x) := \mathcal{B}_\varepsilon \left( \varrho_\varepsilon^\gamma(x) - \frac{1}{|\Omega_\varepsilon|} \int_{\Omega_\varepsilon} \varrho_\varepsilon^\gamma \, dx \right), \quad (2.17)$$

where  $\mathcal{B}_\varepsilon$  is the Bogovskii's operator introduced in Lemma 2.1. We remark that such  $\varphi$  is well defined since  $\varrho_\varepsilon^\gamma \in L^2(\Omega_\varepsilon)$  due to (2.13). Then by Lemma 2.1, we have

$$\operatorname{div}_x \varphi(x) = \varrho_\varepsilon^\gamma(x) - \frac{1}{|\Omega_\varepsilon|} \int_{\Omega_\varepsilon} \varrho_\varepsilon^\gamma \, dx \quad \text{in } \Omega_\varepsilon, \quad (2.18)$$

$$\|\varphi\|_{W_0^{1,2}(\Omega_\varepsilon)} \leq c \left( \|\varrho_\varepsilon^\gamma\|_{L^2(\Omega_\varepsilon)} + \|\varrho_\varepsilon^\gamma\|_{L^1(\Omega_\varepsilon)} \right) \leq c \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^\gamma. \quad (2.19)$$

We plug this test function  $\varphi$  into (1.15) to obtain

$$\int_{\Omega_\varepsilon} p(\varrho_\varepsilon) \varrho_\varepsilon^\gamma \, dx = \sum_{j=1}^4 I_j \quad (2.20)$$

with

$$I_1 := \int_{\Omega_\varepsilon} p(\varrho_\varepsilon) \, dx \frac{1}{|\Omega_\varepsilon|} \int_{\Omega_\varepsilon} \varrho_\varepsilon^\gamma \, dx, \quad I_2 := \int_{\Omega_\varepsilon} \mu \nabla_x \mathbf{u}_\varepsilon : \nabla_x \varphi \, dx + \int_{\Omega_\varepsilon} \left( \frac{\mu}{3} + \eta \right) \operatorname{div}_x \mathbf{u}_\varepsilon : \operatorname{div}_x \varphi \, dx, \quad (2.21)$$

$$I_3 := - \int_{\Omega_\varepsilon} \varrho_\varepsilon \mathbf{u}_\varepsilon \otimes \mathbf{u}_\varepsilon : \nabla_x \varphi \, dx, \quad I_4 := - \int_{\Omega_\varepsilon} (\varrho_\varepsilon \mathbf{f} + \mathbf{g}) \varphi \, dx.$$

Now we estimate  $I_j$  one by one. By the interpolation between  $L^q$  spaces,

$$I_1 = \frac{a}{|\Omega_\varepsilon|} \|\varrho_\varepsilon\|_{L^\gamma(\Omega_\varepsilon)}^{2\gamma} \leq \frac{a}{|\Omega_\varepsilon|} \left( \|\varrho_\varepsilon\|_{L^1(\Omega_\varepsilon)}^{\theta_1} \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{1-\theta_1} \right)^{2\gamma} = \frac{aM^{2\gamma\theta_1}}{|\Omega_\varepsilon|} \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{2\gamma-2\gamma\theta_1},$$

where  $M$  is the total mass and  $\theta_1 \in (0, 1)$  is determined by

$$\frac{1}{\gamma} = \theta_1 + \frac{1-\theta_1}{2\gamma}.$$

By (2.16) and (2.19), we have for  $I_2$ :

$$I_2 \leq c \|\nabla_x \mathbf{u}_\varepsilon\|_{L^2(\Omega_\varepsilon)} \|\nabla_x \varphi\|_{L^2(\Omega_\varepsilon)} \leq c \left( \|\varrho_\varepsilon\|_{L^{\frac{6}{5}}(\Omega_\varepsilon)} + 1 \right) \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^\gamma \leq c \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^\gamma \left( \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)} + 1 \right).$$

Again by (2.16) and (2.19), we have for  $I_3$ :

$$\begin{aligned} I_3 &\leq \|\varrho_\varepsilon\|_{L^6(\Omega_\varepsilon)} \|\mathbf{u}_\varepsilon\|_{L^6(\Omega_\varepsilon)}^2 \|\nabla_x \varphi\|_{L^2(\Omega_\varepsilon)} \leq c \|\varrho_\varepsilon\|_{L^6(\Omega_\varepsilon)} \left( \|\varrho_\varepsilon\|_{L^{\frac{6}{5}}(\Omega_\varepsilon)}^2 + 1 \right) \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^\gamma \\ &\leq c \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{\gamma+1} \left( \|\varrho_\varepsilon\|_{L^1(\Omega_\varepsilon)}^{2\theta_2} \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{2(1-\theta_2)} + 1 \right) \\ &= c M^{2\theta_2} \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{\gamma+3-2\theta_2} + c \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{\gamma+1}, \end{aligned}$$

where  $\theta_2 \in (0, 1)$  is determined by

$$\frac{5}{6} = \theta_2 + \frac{1-\theta_2}{2\gamma}.$$

For  $I_4$ , we have

$$I_4 \leq c \left( \|\varrho_\varepsilon\|_{L^2} + 1 \right) \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^\gamma \leq c \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^\gamma + c \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{\gamma+1}.$$

We sum up the estimates for  $I_j$  and obtain

$$a \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{2\gamma} = \int_{\Omega_\varepsilon} p(\varrho_\varepsilon) \varrho_\varepsilon^\gamma \, dx = \sum_{j=1}^4 I_j \leq c \left( \|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)}^{2\gamma-\beta_1} + 1 \right) \quad (2.22)$$

for some  $\beta_1 > 0$ . Precisely, we can choose  $\beta_1$  as

$$\beta_1 = \min\{2\gamma\theta_1, \gamma-1, \gamma-3+2\theta_2\}.$$

From (2.22), we finally obtain

$$\|\varrho_\varepsilon\|_{L^{2\gamma}(\Omega_\varepsilon)} \leq c, \quad \text{for some } c \text{ independent of } \varepsilon. \quad (2.23)$$

We then go back (2.16) to derive

$$\|\mathbf{u}_\varepsilon\|_{W_0^{1,2}(\Omega_\varepsilon)} \leq c, \quad \text{for some } c \text{ independent of } \varepsilon. \quad (2.24)$$

This completes the proof of the uniform estimate claimed in (1.22).

## 2.3 Equations in a fixed domain

In this section, we derive the equations for the extended functions  $[\tilde{\varrho}_\varepsilon, \tilde{\mathbf{u}}_\varepsilon]$  in  $\Omega$ , where  $[\varrho_\varepsilon, \mathbf{u}_\varepsilon]$  is the finite energy renormalized weak solution in Theorem 1.1. In accordance with the uniform estimates (2.23 – 2.24), we have

$$\|\tilde{\varrho}_\varepsilon\|_{L^{2\gamma}(\Omega)} + \|\tilde{\mathbf{u}}_\varepsilon\|_{W_0^{1,2}(\Omega)} \leq c, \quad \text{for some } c \text{ independent of } \varepsilon. \quad (2.25)$$

### 2.3.1 Continuity equation

First we claim:

**Proposition 2.1** *Under the assumptions in Theorem 1.1, the extended functions  $\tilde{\varrho}_\varepsilon, \tilde{\mathbf{u}}_\varepsilon$  satisfy*

$$\operatorname{div}_x(\tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon) = 0, \quad \operatorname{div}_x(b(\tilde{\varrho}_\varepsilon) \tilde{\mathbf{u}}_\varepsilon) + (\tilde{\varrho}_\varepsilon b'(\tilde{\varrho}_\varepsilon) - b(\tilde{\varrho}_\varepsilon)) \operatorname{div}_x \tilde{\mathbf{u}}_\varepsilon = 0 \quad \text{in } \mathcal{D}'(R^3) \quad (2.26)$$

for any  $b \in C^0([0, \infty)) \cap C^1((0, \infty))$  satisfying (1.18 - 1.19).

This is a direct conclusion from the fact that  $[\varrho_\varepsilon, \mathbf{u}_\varepsilon]$  is a renormalized weak solution.

### 2.3.2 Momentum equation

The following proposition is crucial in the proof of convergence.

**Proposition 2.2** *Under the assumptions in Theorem 1.1, we have*

$$\operatorname{div}_x(\tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon \otimes \tilde{\mathbf{u}}_\varepsilon) + \nabla_x p(\tilde{\varrho}_\varepsilon) = \operatorname{div}_x \mathbb{S}(\nabla_x \tilde{\mathbf{u}}_\varepsilon) + \tilde{\varrho}_\varepsilon \tilde{\mathbf{f}} + \tilde{\mathbf{g}} + F_\varepsilon, \quad \text{in } \mathcal{D}'(\Omega; R^3), \quad (2.27)$$

where  $F_\varepsilon$  is a distribution satisfying

$$| \langle F_\varepsilon, \varphi \rangle_{\mathcal{D}' \times \mathcal{D}} | \leq c \varepsilon^\sigma \|\varphi\|_{L^r(\Omega; R^3)} + c \varepsilon^{\frac{3(\alpha-1)\sigma_0}{2(2+\sigma_0)}} \|\nabla_x \varphi\|_{L^{2+\sigma_0}(\Omega; R^{3 \times 3})}, \quad (2.28)$$

for any  $\varphi \in C_c^\infty((0, T) \times \Omega; R^3)$  and some constants  $c > 0$ ,  $\sigma > 0$ ,  $\sigma_0 > 0$  and  $1 < r < \infty$  independent of  $\varepsilon$ . In particular, we can choose

$$\sigma := \frac{\alpha - 3}{4}, \quad r := \frac{12(\alpha - 1)}{\alpha - 3}, \quad \sigma_0 \in (0, \infty). \quad (2.29)$$

**Proof of Proposition 2.2.** Let  $\varphi \in C_c^\infty((0, T) \times \Omega; R^3)$  be any test function. It is sufficient to show

$$\begin{aligned} I^\varepsilon &:= \int_{\Omega} \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon \otimes \tilde{\mathbf{u}}_\varepsilon : \nabla_x \varphi + p(\tilde{\varrho}_\varepsilon) \operatorname{div}_x \varphi - \mathbb{S}(\nabla_x \tilde{\mathbf{u}}_\varepsilon) : \nabla_x \varphi + \tilde{\varrho}_\varepsilon \tilde{\mathbf{f}} \varphi + \tilde{\mathbf{g}} \varphi \, dx \\ &\leq c \varepsilon^\sigma \|\varphi\|_{L^r(\Omega; R^3)} + c \varepsilon^{\frac{3(\alpha-1)\sigma_0}{2(2+\sigma_0)}} \|\nabla_x \varphi\|_{L^{2+\sigma_0}(\Omega; R^{3 \times 3})}, \end{aligned} \quad (2.30)$$

where  $c$ ,  $\sigma$ ,  $r$  and  $\sigma_0$  are the constants enjoying the properties claimed in Proposition 2.2.

Using (1.1 - 1.3) we can find cut-off functions  $g_\varepsilon \in C_c^\infty(\Omega)$  satisfying

$$0 \leq g_\varepsilon \leq 1, \quad g_\varepsilon = 0 \text{ on } \bigcup_{k \in K_\varepsilon} T_{\varepsilon, k}^s, \quad g_\varepsilon = 1 \text{ in } \Omega \setminus \bigcup_{k \in K_\varepsilon} B_{\varepsilon, k}, \quad \|\nabla_x g_\varepsilon\|_{L^\infty(\Omega)} \leq c \varepsilon^{-\alpha}. \quad (2.31)$$

Together with (1.4), we have for any  $1 \leq q \leq \infty$ :

$$\|1 - g_\varepsilon\|_{L^q(\Omega)} \leq c \varepsilon^{\frac{3(\alpha-1)}{q}}, \quad \|\nabla_x g_\varepsilon\|_{L^q(\Omega)} \leq c \varepsilon^{\frac{3(\alpha-1)}{q} - \alpha}. \quad (2.32)$$

Then direct calculation gives

$$I^\varepsilon = \int_{\Omega_\varepsilon} \varrho_\varepsilon \mathbf{u}_\varepsilon \otimes \mathbf{u}_\varepsilon : \nabla_x (g_\varepsilon \varphi) + p(\varrho_\varepsilon) \operatorname{div}_x (g_\varepsilon \varphi) - \mathbb{S}(\nabla_x \mathbf{u}_\varepsilon) : \nabla_x (g_\varepsilon \varphi) + (\varrho_\varepsilon \mathbf{f} + \mathbf{g})(g_\varepsilon \varphi) \, dx + \sum_{j=1}^3 I_j^\varepsilon, \quad (2.33)$$

where

$$\begin{aligned} I_1^\varepsilon &:= \int_{\Omega} \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon \otimes \tilde{\mathbf{u}}_\varepsilon : (1 - g_\varepsilon) \nabla_x \varphi - \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon \otimes \tilde{\mathbf{u}}_\varepsilon : (\nabla_x g_\varepsilon \otimes \varphi) \, dx, \\ I_2^\varepsilon &:= \int_{\Omega} p(\tilde{\varrho}_\varepsilon) (1 - g_\varepsilon) \operatorname{div}_x \varphi - p(\tilde{\varrho}_\varepsilon) \nabla_x g_\varepsilon \cdot \varphi \, dx, \\ I_3^\varepsilon &:= \int_{\Omega} \mathbb{S}(\nabla_x \tilde{\mathbf{u}}_\varepsilon) : (1 - g_\varepsilon) \nabla_x \varphi + \mathbb{S}(\nabla_x \tilde{\mathbf{u}}_\varepsilon) : (\nabla_x g_\varepsilon \otimes \varphi) \, dx. \end{aligned}$$

Since  $g_\varepsilon \varphi \in C_c^\infty(\Omega_\varepsilon)$  is a test function for the stationary Navier-Stokes equations in  $\Omega_\varepsilon$ , we have

$$I^\varepsilon = \sum_{j=1}^3 I_j^\varepsilon.$$

For  $I_1$ , using hypothesis  $\gamma \geq 3$  and Hölder's inequality, we get

$$I_1 \leq c \|\tilde{\varrho}_\varepsilon\|_{L^{2\gamma}(\Omega)} \|\tilde{\mathbf{u}}_\varepsilon\|_{L^6(\Omega)}^2 (\|(1 - g_\varepsilon) \nabla_x \varphi\|_{L^2(\Omega)} + \|\nabla_x g_\varepsilon \otimes \varphi\|_{L^2(\Omega)}). \quad (2.34)$$

By the uniform estimates (2.23 - 2.24 - 2.25) and Hölder's inequality, we have

$$I_1 \leq c \left( \|(1 - g_\varepsilon)\|_{L^{\frac{2(2+\sigma_0)}{\sigma_0}}(\Omega)} \|\nabla_x \varphi\|_{L^{2+\sigma_0}(\Omega)} + \|\nabla_x g_\varepsilon\|_{L^{r_1}(\Omega)} \|\varphi\|_{L^{r_2}(\Omega)} \right), \quad (2.35)$$

where

$$\sigma_0 \in (0, \infty), \quad r_j \in (2, \infty), \quad \frac{1}{r_1} + \frac{1}{r_2} = \frac{1}{2}.$$

Then by (2.32), we deduce

$$I_1 \leq c \varepsilon^{\frac{3(\alpha-1)\sigma_0}{2(2+\sigma_0)}} \|\nabla_x \varphi\|_{L^{2+\sigma_0}(\Omega)} + c \varepsilon^{\frac{3(\alpha-1)}{r_1} - \alpha} \|\varphi\|_{L^{r_2}(\Omega)},$$

where the number

$$\sigma := \frac{3(\alpha-1)}{r_1} - \alpha = \frac{3(\alpha-1)}{2} - \alpha - \frac{3(\alpha-1)}{r_2} = \frac{\alpha-3}{2} - \frac{3(\alpha-1)}{r_2}$$

is strictly positive as long as

$$r_2 > \frac{6(\alpha-1)}{\alpha-3}.$$

In particular, we can choose

$$r_2 = \frac{12(\alpha-1)}{\alpha-3} \text{ such that } \sigma := \frac{3(\alpha-1)}{r_1} - \alpha = \frac{\alpha-3}{4},$$

which is exactly (2.29).

Seeing that  $I_2^\varepsilon$  and  $I_3^\varepsilon$  can be handled similarly to  $I_1^\varepsilon$ , we have completed the proof of Proposition 2.2.

## 2.4 Passing to the limit

By the uniform estimates (2.25), up to a subsequence, we have

$$\tilde{\varrho}_\varepsilon \rightarrow \varrho \text{ weakly in } L^{2\gamma}(\Omega), \quad \tilde{\mathbf{u}}_\varepsilon \rightarrow \mathbf{u} \text{ weakly in } W_0^{1,2}(\Omega; \mathbb{R}^3). \quad (2.36)$$

It is left to show the limit  $[\varrho, \mathbf{u}]$  represents a finite energy renormalized weak solution to (1.5 – 1.9) in  $\Omega$ .

### 2.4.1 Strong convergence of velocity

By compact Sobolev embedding, we have the strong convergence

$$\tilde{\mathbf{u}}_\varepsilon \rightarrow \mathbf{u} \quad \text{strongly in } L^q(\Omega; R^3) \quad \text{for any } 1 \leq q < 6. \quad (2.37)$$

Together with the weak convergence of the density, we have the weak convergence of nonlinear terms:

$$\tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon \rightarrow \varrho \mathbf{u} \quad \text{weakly in } L^q(\Omega; R^3) \quad \text{for any } 1 < q < \frac{6\gamma}{3+\gamma}, \quad (2.38)$$

and

$$\tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon \otimes \tilde{\mathbf{u}} \rightarrow \varrho \mathbf{u} \otimes \mathbf{u} \quad \text{weakly in } L^q(\Omega; R^{3 \times 3}) \quad \text{for any } 1 < q < \frac{6\gamma}{3+2\gamma}. \quad (2.39)$$

Then we let  $\varepsilon \rightarrow 0$  in the first equation of (2.26) and in equation (2.27) to deduce the following two equations in  $\mathcal{D}'(\Omega)$ :

$$\operatorname{div}_x(\varrho \mathbf{u}) = 0, \quad (2.40)$$

$$\operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x \overline{p(\varrho)} = \operatorname{div}_x \mathbb{S}(\nabla_x \mathbf{u}) + \varrho \mathbf{f} + \mathbf{g}. \quad (2.41)$$

Here  $\overline{p(\varrho)}$  is the weak limit of  $p(\tilde{\varrho}_\varepsilon)$  in  $L^2(\Omega)$ . Moreover, in accordance with Remark 1.2, the pair of functions  $[\varrho, \mathbf{u}]$  satisfies the renormalized equation

$$\operatorname{div}_x(\varrho \mathbf{u}) = 0, \quad \operatorname{div}_x(b(\varrho) \mathbf{u}) + (\varrho b'(\varrho) - b(\varrho)) \operatorname{div}_x \mathbf{u} = 0, \quad \text{in } \mathcal{D}'(R^3), \quad (2.42)$$

for any  $b$  satisfying (1.18 - 1.19).

Consequently, to finish the proof of Theorem 1.1, we have to show  $\overline{p(\varrho)} = p(\varrho)$ . This is done in the next section.

### 2.4.2 Convergence of the pressure

First of all, we introduce the so-called effective viscous flux which is the quantity  $p(\varrho) - (\frac{4\mu}{3} + \eta) \operatorname{div}_x \mathbf{u}$ . We shall show that this quantity enjoys some weak compactness property specified in the following lemma. This property, identified first by Lions [12], plays a crucial role in the existence theory in the framework of weak solutions for the compressible Navier-Stokes system.

**Lemma 2.3** *For any  $\psi \in C_c^\infty(\Omega)$ , there holds up to a subsequence:*

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} \psi \left( p(\tilde{\varrho}_\varepsilon) - \left( \frac{4\mu}{3} + \eta \right) \operatorname{div}_x \tilde{\mathbf{u}}_\varepsilon \right) \tilde{\varrho}_\varepsilon \, dx = \int_{\Omega} \psi \left( \overline{p(\varrho)} - \left( \frac{4\mu}{3} + \eta \right) \operatorname{div}_x \mathbf{u} \right) \varrho \, dx. \quad (2.43)$$

**Proof of Lemma 2.3.** The proof of Lemma 2.3 is quite technical but nowadays well understood. The idea is to take test functions of the form

$$\psi \nabla \Delta^{-1}(1_\Omega \tilde{\varrho}_\varepsilon), \quad \psi \nabla \Delta^{-1}(1_\Omega \varrho),$$

where  $\psi \in C_c^\infty(\Omega)$  and  $\Delta^{-1}$  is the Fourier multiplier on  $R^3$  with the symbol  $1/|\xi|^2$ . It is straightforward to check that

$$\nabla \nabla \Delta^{-1} = (\mathcal{R}_{i,j})_{1 \leq i,j \leq 3}$$

are the Riesz operators. In particular, using the well known properties of the singular integral operators of Carderón-Zygmund type, we have

$$\|\nabla \nabla \Delta^{-1}(r)\|_{L^q(R^3)} \leq c \|r\|_{L^q(R^3)}.$$

for any  $r \in L^q(R^3)$ ,  $1 < q < \infty$ .

We take  $\psi \nabla \Delta^{-1}(1_\Omega \tilde{\varrho}_\varepsilon)$  as a test functions in the weak formulation of (2.27) and let  $\varepsilon \rightarrow 0$ . Then we take  $\psi \nabla \Delta^{-1}(1_\Omega \varrho)$  as a test functions in the weak formulation of (2.41), and compare the results of these operations. By using the convergence results (2.36 - 2.39), compact Sobolev embedding, the fact  $\gamma \geq 3$ , and the property  $\mathcal{R}_{i,j} = \mathcal{R}_{j,i}$ , we obtain, through long but straightforward calculations, that

$$\begin{aligned} I &:= \lim_{\varepsilon \rightarrow 0} \int_\Omega \psi \left( p(\tilde{\varrho}_\varepsilon) - \left( \frac{4\mu}{3} + \eta \right) \operatorname{div}_x \tilde{\mathbf{u}}_\varepsilon \right) \tilde{\varrho}_\varepsilon \, dx - \int_\Omega \psi \left( \overline{p(\varrho)} - \left( \frac{4\mu}{3} + \eta \right) \operatorname{div}_x \mathbf{u} \right) \varrho \, dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_\Omega \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon^i \tilde{\mathbf{u}}_\varepsilon^j \psi \mathcal{R}_{i,j}(1_\Omega \tilde{\varrho}_\varepsilon) \, dx - \int_\Omega \varrho \mathbf{u}^i \mathbf{u}^j \psi \mathcal{R}_{i,j}(1_\Omega \varrho) \, dx. \end{aligned} \quad (2.44)$$

On the other hand, by taking  $1_\Omega \nabla \Delta^{-1}(\psi \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon)$  as a test function for the first equation of (2.26) and taking  $1_\Omega \nabla \Delta^{-1}(\psi \varrho \mathbf{u})$  as a test function for (2.40), we obtain

$$\int_\Omega 1_\Omega \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon^i \mathcal{R}_{i,j}(\psi \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon) \, dx = 0, \quad \int_\Omega 1_\Omega \varrho \mathbf{u}^i \mathcal{R}_{i,j}(\psi \varrho \mathbf{u}) \, dx = 0. \quad (2.45)$$

Plugging (2.45) into (2.44) yields

$$I = \lim_{\varepsilon \rightarrow 0} \int_\Omega \tilde{\mathbf{u}}_\varepsilon^i \left( \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon^j \psi \mathcal{R}_{i,j}(1_\Omega \tilde{\varrho}_\varepsilon) - 1_\Omega \tilde{\varrho}_\varepsilon \mathcal{R}_{i,j}(\psi \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon) \right) \, dx - \int_\Omega \mathbf{u}^i \left( \varrho \mathbf{u}^j \psi \mathcal{R}_{i,j}(1_\Omega \varrho) - 1_\Omega \varrho \mathcal{R}_{i,j}(\psi \varrho \mathbf{u}) \right) \, dx. \quad (2.46)$$

The proof of the strong convergence of the densities is finished by means of the strong convergence of the velocity in (2.37) and the following property of commutators of the type appearing in (2.46):

**Lemma 2.4** *Suppose*

$$u_\varepsilon \rightarrow u \quad \text{weakly in } L^p(R^3), \quad v_\varepsilon \rightarrow v \quad \text{weakly in } L^q(R^3), \quad \frac{1}{p} + \frac{1}{q} = \frac{1}{r} < 1.$$

Then for any  $1 \leq i, j \leq 3$ ,

$$u_\varepsilon \mathcal{R}_{i,j}(v_\varepsilon) - v_\varepsilon \mathcal{R}_{i,j}(u_\varepsilon) \rightarrow u \mathcal{R}_{i,j}(v) - v \mathcal{R}_{i,j}(u) \quad \text{weakly in } L^r(R^3).$$

See [8, Lemma 3.4] for the proof.

Now, we are ready to prove the following result:

**Proposition 2.3** *Let  $\overline{p(\varrho)\varrho}$  be a weak limit of  $p(\tilde{\varrho}_\varepsilon)\tilde{\varrho}_\varepsilon$  in  $L^{\frac{2\gamma}{\gamma+1}}(\Omega)$ , we then have*

$$\overline{p(\varrho)\varrho} = \overline{p(\varrho)}\varrho \quad \text{in } L^{\frac{2\gamma}{\gamma+1}}(\Omega).$$

**Proof of Proposition 2.3.** Using the uniform bound for  $\tilde{\varrho}_\varepsilon$  in (2.25) and the specific form of  $p(\varrho)$  in (1.11), we have

$$p(\tilde{\varrho}_\varepsilon)\tilde{\varrho}_\varepsilon \rightarrow \overline{p(\varrho)\varrho} \quad \text{weakly in } L^{\frac{2\gamma}{\gamma+1}}(\Omega).$$

Taking  $b(s) = s \log s$  in the renormalized equations (2.26) and (2.42) we obtain

$$\operatorname{div}_x(\tilde{\varrho}_\varepsilon \log \tilde{\varrho}_\varepsilon \tilde{\mathbf{u}}_\varepsilon) + \tilde{\varrho}_\varepsilon \operatorname{div}_x \tilde{\mathbf{u}}_\varepsilon = 0, \quad \operatorname{div}_x(\varrho \log \varrho \mathbf{u}) + \varrho \operatorname{div}_x \mathbf{u} = 0. \quad (2.47)$$

Letting  $\varepsilon \rightarrow 0$  in (2.47) yields

$$\operatorname{div}_x(\overline{\varrho \log \varrho} \mathbf{u}) + \overline{\varrho} \operatorname{div}_x \mathbf{u} = 0.$$

Then for any  $\psi \in C_c^\infty(\Omega)$ , we have

$$\int_\Omega \psi \overline{\varrho \operatorname{div}_x \mathbf{u}} \, dx = \int_\Omega \nabla_x \psi \cdot (\overline{\varrho \log \varrho} \mathbf{u}) \, dx, \quad \int_\Omega \psi \varrho \operatorname{div}_x \mathbf{u} \, dx = \int_\Omega \nabla_x \psi \cdot (\varrho \log \varrho) \mathbf{u} \, dx. \quad (2.48)$$

Passing to the limit for  $\varepsilon \rightarrow 0$  in (2.43) and using (2.48) we get

$$\int_\Omega \psi \overline{p(\varrho)\varrho} - \left(\frac{4\mu}{3} + \eta\right) \nabla_x \psi \cdot (\overline{\varrho \log \varrho} \mathbf{u}) \, dx = \int_\Omega \psi \overline{p(\varrho)\varrho} - \left(\frac{4\mu}{3} + \eta\right) \nabla_x \psi \cdot (\varrho \log \varrho) \mathbf{u} \, dx. \quad (2.49)$$

Now we choose test functions  $\psi_n \in C_c^\infty(\Omega)$  such that

$$0 \leq \psi_n \leq 1, \quad \psi_n(x) = 1 \text{ if } d(x, \partial\Omega) > \frac{2}{n}, \quad \psi_n(x) = 0 \text{ if } d(x, \partial\Omega) < \frac{1}{n}, \quad \|\nabla_x \psi_n\|_{L^\infty(\Omega)} \leq 2n$$

and for any  $q \in [1, \infty]$ :

$$\|1 - \psi_n\|_{L^q} \leq c n^{-\frac{1}{q}}, \quad \|\nabla_x \psi_n\|_{L^q} \leq c n^{1-\frac{1}{q}},$$

and consequently

$$\|d(x, \partial\Omega) \nabla_x \psi_n\|_{L^q} \leq c n^{-\frac{1}{q}}$$

The velocity  $\mathbf{u} \in W_0^{1,2}(\Omega; R^3)$ , then we have

$$[d(x, \partial\Omega)]^{-1} \mathbf{u} \in L^2(\Omega; R^3).$$

Therefore,

$$\int_{\Omega} \nabla_x \psi_n \cdot (\overline{\varrho \log \varrho}) \mathbf{u} \, dx \leq \|d(x, \partial\Omega) \nabla_x \psi_n\|_{L^{10}} \|\overline{\varrho \log \varrho}\|_{L^{5/2}} \|[d(x, \partial\Omega)]^{-1} \mathbf{u}\|_{L^2} \leq c n^{-1/10}.$$

Letting  $\psi = \psi_n$  in (2.49) and passing to the limit  $n \rightarrow \infty$  we may infer that

$$\int_{\Omega} \overline{p(\varrho) \varrho} - \overline{p(\varrho)} \varrho \, dx = 0. \quad (2.50)$$

Since the mapping  $\varrho \mapsto p(\varrho)$  is strictly increasing, there holds

$$\overline{p(\varrho) \varrho} \geq \overline{p(\varrho)} \varrho, \quad \text{a.e. in } \Omega,$$

which, together with (2.50), gives rise to the desired conclusion

$$\overline{p(\varrho) \varrho} = \overline{p(\varrho)} \varrho, \quad \text{a.e. in } \Omega.$$

We have completed the proof of Proposition 2.3.

Our desired result  $\overline{p(\varrho)} = p(\varrho)$  is a direct corollary of Proposition 2.3, due to the monotonicity and convexity of  $p(\cdot)$ . Accordingly, we have finished the proof of Theorem 1.1.

### 3 Concluding remarks

The hypothesis concerning the shape of the holes as well as their spatial distribution may be considerably relaxed. As a matter of fact, it is enough to impose the following restriction on the model hole:

There exists a constant  $\omega > 0$  such that at each point  $x \in \partial T^s$  there exists a closed cone  $C_x$  with vertex at  $x$  and of aperture  $\omega$  such that

$$C_x \cap T^s = \{x\}.$$

Moreover, the holes need not be periodically distributed, it is enough that their mutual distance is proportional to  $\varepsilon$ , see [7] for the relevant homogenization problem in the context of *incompressible* fluids.

## References

- [1] G. Allaire, Homogenization of the Navier-Stokes equations in open sets perforated with tiny holes. I. Abstract framework, a volume distribution of holes. *Arch. Ration. Mech. Anal.*, **113** (3) (1990) 209-259.
- [2] G. Allaire, Homogenization of the Navier-Stokes equations in open sets perforated with tiny holes. II. Noncritical sizes of the holes for a volume distribution and a surface distribution of holes. *Arch. Ration. Mech. Anal.*, **113** (3) (1990) 261-298.
- [3] M. E. Bogovskii. Solution of some vector analysis problems connected with operators div and grad (in Russian). *Trudy Sem. S.L. Sobolev*, **80**(1):5-40, 1980.
- [4] J. Březina and A. Novotný. On weak solutions of steady Navier-Stokes equations for monatomic gas. *Comment. Math. Univ. Carolin.*, **49**(4):611-632, 2008.
- [5] R.j. DiPerna, P.-L. Lions, Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.*, **98** (1989), 511-547.
- [6] E. Feireisl. *Dynamics of Viscous Compressible Fluids*. Oxford University Press, Oxford, 2003.
- [7] E. Feireisl, Y. Namlyeyeva, and Š. Nečasová. Homogenization of the evolutionary Navier-Stokes system. *Manusc. Math.*, 2015. Submitted.
- [8] E. Feireisl, A. Novotný, and H. Petzeltová. On the existence of globally defined weak solutions to the Navier-Stokes equations of compressible isentropic fluids. *J. Math. Fluid Mech.*, **3**, 358-392, 2001.
- [9] E. Feireisl, A. Novotný and T. Takahashi. Homogenization and singular limits for the complete Navier-Stokes-Fourier system. *J. Math. Pures Appl.*, **94** (2010), no. 1, 33-57.
- [10] J. Frehse, M. Steinhauer, and W. Weigant. The Dirichlet problem for steady viscous compressible flow in three dimensions. *J. Math. Pures Appl. (9)*, **97**(2):85-97, 2012.
- [11] G. P. Galdi. *An introduction to the mathematical theory of the Navier - Stokes equations, I*. Springer-Verlag, New York, 1994.
- [12] P.-L. Lions. *Mathematical topics in fluid dynamics, Vol.2, Compressible models*. Oxford Science Publication, Oxford, 1998.
- [13] N. Masmoudi. Homogenization of the compressible Navier-Stokes equations in a porous medium. *ESAIM Control Optim. Calc. Var.*, **8** (2002) 885-906.

- [14] A. Mikelič. Homogenization of nonstationary Navier-Stokes equations in a domain with a grained boundary. *Ann. Mat. Pura Appl.*, **158** (1991) 167-179.
- [15] A. Novotný, I. Straškraba. *Introduction to the mathematical theory of compressible flow*. Oxford University Press, Oxford, 2004.
- [16] P. Plotnikov and J. Sokołowski. *Compressible Navier-Stokes equations*, volume 73 of *Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series) [Mathematics Institute of the Polish Academy of Sciences. Mathematical Monographs (New Series)]*. Birkhäuser/Springer Basel AG, Basel, 2012. Theory and shape optimization.