

- [12] W.T. Tutte, A geometrical version of the four color problem, *Combinatorial Mathematics and its Applications*, (eds. R. C. Bose and T. A. Dowling), University of North Carolina Press, Chapel Hill, 1967.
- [13] W.T. Tutte, Hamiltonian circuits, *Colloquio Internazionale sulle Teorie Combinatorie*, *Atti dei Convegni Lincei* 17, Accad. Naz. Lincei, Roma I (1976) 193-199.
- [14] C.-Q. Zhang, Hamiltonian weights and unique edge-3-colorings of cubic graphs, *J. Graph Theory* 20, No. 1 (1995), 91-99.

## Completely Strong Path-Connectivity of Local Tournaments\*

Bu Yue Hua

Department of Mathematics  
Zhejiang Normal University  
Jinhua 321004  
China

Zhang Ke Min

Department of Mathematics  
Nanjing University  
Nanjing 210008  
China

ABSTRACT. Let  $T = (V, A)$  be an oriented graph with  $n$  vertices.  $T$  is completely strong path-connected if for each arc  $(a, b) \in A$  and  $k$  ( $k = 2, \dots, n - 1$ ), there is a path from  $b$  to  $a$  of length  $k$  (denoted by  $P_k(a, b)$ ) and a path from  $a$  to  $b$  of length  $k$  (denoted by  $P'_k(a, b)$ ) in  $T$ . In this paper, we prove that a connected local tournament  $T$  is completely strong path-connected iff for each arc  $(a, b) \in A$ , there exist  $P_2(a, b)$  and  $P'_2(a, b)$  in  $T$ , and  $T \not\cong T_0 - D'_8$ -type digraph and  $D_8$ .

### 1 Introduction

Let  $T = (V, A)$  be an oriented graph with  $n$  vertices. If an arc  $(x, y) \in A$ , then we say that  $x$  dominates  $y$ , denoted by  $x \rightarrow y$ . If  $S_1$  and  $S_2$  are disjoint subsets of  $V$  such that there is a complete connection between them and all arcs between them are directed toward  $S_2$ , we say that  $S_1$  dominates  $S_2$ , denoted by  $S_1 \rightarrow S_2$ . We write  $x \rightarrow S_2$  (resp.,  $S_2 \rightarrow x$ ) instead of  $\{x\} \rightarrow S_2$  (resp.,  $S_2 \rightarrow \{x\}$ ). For  $x \in V$ , we define  $O(x) = \{y \mid y \in V, (x, y) \in A\}$ ,  $I(x) = \{y \mid y \in V, (y, x) \in A\}$ .

\*The Project Supported by NSFC No. 19471037 and SFJS.

\* (6)  
MR# 00a: a

$T$  is arc- $k$ -cyclic if each arc  $(a, b) \in A$ , there is a path from  $b$  to  $a$  of length  $k - 1$  in  $T$ .  $T$  is arc-pancyclic (resp., arc-antipancyclic) if for each arc  $(a, b) \in A$ , there is a path from  $b$  to  $a$  (resp., from  $a$  to  $b$ ) of length  $k$  ( $k = 2, 3, \dots, n - 1$ ) in  $T$ , denoted by  $P_k(a, b)$ , or briefly  $P_k$  (resp.,  $P'_k(a, b)$ ,  $P'_k$ ). An oriented graph  $T$  is completely strong path-connected if  $T$  is arc-pancyclic and arc-antipancyclic. Other notations and terminologies not defined in this paper can be found in [3].

A local tournament  $T$  is an oriented graph such that  $T[O(x)]$  and  $T[I(x)]$  are tournaments for every vertex  $x$  in  $T$ . Local tournaments were first introduced by J. Bang-Jensen [1], [2]. Clearly, tournaments is a special class of local tournaments. In [1], [2], it was shown that every connected local tournament has a Hamiltonian path, and every strong local tournament has a Hamiltonian cycle. Many other results for tournaments are also shown for local tournaments. In this paper, Zhang and Wu's results in [5] and [6] are extended. We get the following main result.

**Theorem.** Let  $T = (V, A)$  be a connected local tournament with  $n$  vertices ( $n \geq 3$ ). If for each arc  $(a, b) \in A$ , there exist  $P_2(a, b)$  and  $P'_2(a, b)$  in  $T$ . Then  $T$  is completely strong path-connected, except  $T \simeq T_0$ - or  $D'_8$ -type digraph or  $D_8$ . (see Figures 1, 2 and 3).




Figure 1.  $D_8$




Figure 2.  $T_0$ -type digraph.  
(Here  $T'_0, T''_0$  are tournaments)




Figure 3.  $D'_8$ -type digraph.  
(Here  $T_1$  is a tournament)

Immediately we have,

**Corollary.** ([5], Theorem 1) A tournament  $T = (V, A)$  with  $n$  vertices is completely strong path-connected if and only if for each arc  $(a, b) \in A$ , there exist  $P_2(a, b)$ , and  $P'_2(a, b)$  in  $T$ , and  $T \not\cong T_0$ -type digraph.

## 2 The Proof of the Theorem

In order to prove the Theorem, we need the following lemmas.

**Lemma 1.** Let  $T = (V, A)$  be a connected local tournament. For each arc  $(a, b) \in A$ , there exist  $P_2(a, b)$  and  $P'_2(a, b)$  in  $T$ , then there exists a cycle in the induced subgraph  $T[O(x_0)]$  (resp.,  $T[I(x_0)]$ ) for any  $x_0 \in V$ . Furthermore,  $|O(x_0)| \geq 3$ , (resp.,  $|I(x_0)| \geq 3$ ).

**Lemma 2.** Let  $T = (V, A)$  be a connected local tournament. For each arc  $(a, b) \in A$ , there exist  $P_2(a, b)$  and  $P'_2(a, b)$  in  $T$ , then there always exists a  $P'_k(a, b)$  in  $T$  for each arc  $(a, b) \in A$  ( $k = 2, 3, \dots, 6$ ).


By the definition of a local tournament, the proof of Lemma 1 and Lemma 2 is an analogous to the proof of Lemma 1 and Lemma 3 in [7].

**Lemma 3.** ([4] Theorem 1) Except for  $T_6$ -,  $T_8$ -type digraphs and  $D_8$  (see Figures 1 and 5), every arc-3-cyclic connected local tournament is arc-pancyclic.

The proof of the Theorem.

Let  $T = (V, A)$  be a connected local tournament of order  $n$  ( $n \geq 3$ ) such that for each arc  $(a, b) \in A$ , there exist  $P_2(a, b)$  and  $P'_2(a, b)$  in  $T$ . For  $T_6$ - or  $T_8$ -type digraph, it is easy to find that there exists a vertex  $x$  such that  $|O(x)| = 2$ . So  $T$  is not a  $T_6$ - or a  $T_8$ -type digraph by Lemma 1. Hence by Lemma 3  $T$  is an arc-pancyclic local tournament except  $T$  is isomorphic

to  $D_8$ . And by Lemma 2 there always exists a  $P'_k(a, b)$  in  $T$  for  $k \leq 6$ . Therefore it is enough to prove the following.



$T_6$ -type digraph  
(where  $T_1, T_1'$  are tournaments)

$T_8$ -type digraph

Figure 5.

The directions of the edges without arrow can be chosen arbitrary.

**Proposition.** Suppose  $T$  is not isomorphic to a  $T_0$ - or  $D'_8$ -type digraph or  $D_8$ . If for each arc  $(a, b) \in A$  and  $k$  ( $7 \leq k \leq n-1$ ), there exists a  $P'_{k-1}(a, b)$  in  $T$ . Then there exists a  $P'_k(a, b)$  in  $T$ .

From now on, we shall assume that there is a  $P'_{k-1}(a, b)$  in  $T$ , and denote it by  $(1, 2, \dots, k)$ , where  $a = 1$  and  $b = k$ . The set of vertices  $\{1, 2, \dots, k\}$  of  $P'_{k-1}(a, b)$  is also denoted by  $P'_{k-1}$ . Let  $W = V - P'_{k-1}$ . Hence  $|W| \geq 1$ . For any  $w \in W$  we define

$$O'(w) \equiv O(w) \cap P'_{k-1}, \quad I'(w) \equiv I(w) \cap P'_{k-1}.$$

When  $O'(w) \neq \emptyset$  and  $I'(w) \neq \emptyset$  for  $w \in W$ , set

$$a(w) = \max\{i \mid i \in O'(w)\}, \quad b(w) = \min\{i \mid i \in I'(w)\}.$$

If the condition of the proposition were false, we should assume that

$$\text{There does not exist any } P'_k(a, b) \text{ in } T. \quad (*)$$

By the assumption above, we may obtain the following claims.

(1)  $O'(w) = \{1, 2, \dots, a(w)\}$  and  $a(w) < k$  as  $O'(w) \neq \emptyset$ . Similarly,  $I'(w) = \{b(w), \dots, k\}$  and  $b(w) > 1$  as  $I'(w) \neq \emptyset$ .

Suppose  $O(w) \neq \emptyset$ . If there exists an  $i \in O'(w)$  with  $i-1 \notin O'(w)$ , by the definition of a local tournament and  $\{w, i-1\} \subseteq I(i)$ , then  $i-1$  and  $w$  are adjacent in  $T$ . Thus  $i-1 \rightarrow w$  by the definition of  $i$ . Hence

there is a  $P'_k(a, b) = (1, \dots, i-1, w, i, \dots, k)$  in  $T$ . This contradicts  $(*)$ . So  $O'(w) = \{1, 2, \dots, a(w)\}$ . And if  $a(w) = k$ , then  $w \rightarrow P'_{k-1}$ . Note that there exists a  $P_2(w, 1) = (1, x, w)$ . Clearly  $x \notin P'_{k-1}$ . Hence  $x \in W$ . Thus  $T$  contains a  $P'_k(a, b) = (1, x, w, 3, \dots, k)$ . This contradicts  $(*)$ . So  $a(w) < k$ .

(2) For any  $w \in W$ ,  $O'(w) \neq \emptyset$  if and only if  $I'(w) \neq \emptyset$ .

If  $O'(w) \neq \emptyset$ , there is a  $P_2(w, 1) = (1, x, w)$ . If  $x \in W$ , then  $1 \in I'(x)$  and  $b(x) = 1$ . This contradicts  $b(w) > 1$  by (1). Hence  $x \in I'(w)$  and  $I'(w) \neq \emptyset$ . Similarly, if  $I'(w) \neq \emptyset$ , then  $O'(w) \neq \emptyset$ .

(3) Let  $W_1 = \{w \mid w \in W, O'(w) \neq \emptyset\}$  and  $W_2 = W - W_1$  then  $W_2 = \emptyset$ . Furthermore,  $T[W]$  is a tournament and  $O'(w) \neq \emptyset$ ,  $I'(w) \neq \emptyset$  for every  $w \in W$ .

Since  $T$  is connected,  $W_1 \neq \emptyset$ . Suppose  $W_2 \neq \emptyset$ . Let  $w_1 \in W_1$  and  $w_2 \in W_2$  such that  $w_1$  and  $w_2$  are adjacent. Without loss of generality, we assume  $w_2 \rightarrow w_1$ . Since  $k \rightarrow w_1$  by (1) and (2),  $w_2$  and  $k$  are adjacent. Then  $k \rightarrow w_2$  and  $O'(w_2) \neq \emptyset$  by (1) and (2). This is a contradiction. Hence  $W_2 = \emptyset$ . i.e.  $W = W_1$ .

From (1) and (2), we have  $W \subseteq I(1)$  and  $O'(w) \neq \emptyset$ ,  $I'(w) \neq \emptyset$  for every  $w \in W$ . Thus  $T[W]$  is a tournament by the definition of a local tournament.

(4)  $b(w) = b(w')$  and  $a(w) = a(w')$  for any  $w, w' \in W$ .

Suppose there are  $w, w' \in W$  such that  $b(w) \neq b(w')$ . Set  $b(w_0) = \min\{b(w) \mid w \in W\}$ . Let  $W_3 = \{w \mid w \in W, b(w) > b(w_0)\}$  and  $W_4 = W - W_3$ . Then  $W_3 \neq \emptyset$ ,  $W_4 \neq \emptyset$  and  $b(w_0) = b(w) \rightarrow w$  for any  $w \in W_4$ .

Case 1. There exist  $w_3 \in W_3$  and  $w_4 \in W_4$  such that  $w_3 \rightarrow w_4$ .

Since  $b(w_4) = b(w_0) < b(w_3)$  and  $b(w_3) - 1 \rightarrow w_4$ ,  $w_3$  and  $b(w_3) - 1$  are adjacent by  $w_3 \rightarrow w_4$  and the definition of a local tournament. From (1) we have  $w_3 \rightarrow b(w_3) - 1$ . Thus  $a(w_3) = b(w_3) - 1$ . Similarly, since  $b(w_4) - 1 < b(w_3) - 1 = a(w_3)$  and  $w_3 \rightarrow w_4$ , we have  $a(w_4) = b(w_4) - 1$ .

Now we need the following three Lemmas

**Lemma 4.** There are no  $u, v, n$  and  $m$  in  $P'_{k-1}$  such that  $u < n \leq b(w_4) - 1 < b(w_3) \leq v < m$  and  $(u, v), (n, m) \in A$ .

**Proof:** Otherwise, it will contradict  $(*)$ .  $\square$

Now,  $(n, m), (u, v) \in A$  are called cis-crosswise arcs with respect to the  $P'_{k-1}$  (briefly cis-crosswise arcs) if  $n, m, u$  and  $v$  are on  $P'_{k-1}$  such that  $u < n < v < m$ .

**Lemma 5.** (a) For each  $i \in \{3, 4, \dots, b(w_4) - 1\}$ , we have  $(i, 1) \in A$ . (b) For each  $j \in \{b(w_3), \dots, k - 2\}$ , we have  $(k, j) \in A$ .

**Proof:** (a) Since  $\{1, 2, \dots, b(w_4) - 1 = a(w_4)\} \subseteq O(w_4)$ ,  $T[\{3, 4, \dots, b(w_4) - 1, 1\}]$  is a tournament. If there is an  $i_0 \in \{3, 4, \dots, b(w_4) - 1\}$  such that  $1 \rightarrow i_0$ , then  $w_3 \rightarrow i_0 - 1$  by  $i_0 - 1 < a(w_3)$ . There is a  $P_2(w_3, i_0 - 1) =$

$(i_0-1, u, w_3)$ . By the definition of  $b(w_4)$ , we have  $u \notin W$ . Hence  $u \in I'(w_3)$ . Thus there is a  $P'_k(a, b) = (1, i_0, \dots, u-1, w_4, 2, \dots, i_0-1, u, \dots, k)$ . This contradicts  $(*)$ . So (a) is valid.

An analogous proof of (a), we have that (b) is true.  $\square$

**Lemma 6.** *If  $(b(w_4)-1, b(w_3)) \in A$  and  $(b(w_4)-1, b(w_3)) \neq (a, b) = (1, k)$ , then there is an arc  $(u, v) \in A$  such that  $(u, v)$  and  $(b(w_4)-1, b(w_3))$  are cis-crosswise arcs.*

**Proof:** By Lemmas 1, 4 and 5, using an analogous proof of Lemma 3 in [5], Lemma 6 follows.  $\square$

Now, let's back to discuss case 1.

There are  $P_2(w_3, a(w_4) = b(w_4)-1) = (b(w_4)-1, m, w_3)$  and  $P_2(w_3, w_4) = (w_4, w_5, w_3)$ , where  $m \notin W$  and  $w_5 \notin P'_{k-1}$ , by the choice of  $w_4$  and  $b(w_3) > b(w_4)$ . Hence we have that  $b(w_3) \leq m \leq k$  and  $w_5 \in W$ .

If  $b(w_3) - b(w_4) \geq 4$ , then  $a(w_3) \geq b(w_4) + 3$ . There is a  $P'_k(a, b) = (1, \dots, b(w_4), w_4, w_5, w_3, b(w_4) + 3, \dots, a(w_3), \dots, k)$ . This contradicts  $(*)$ . Hence  $b(w_3) - b(w_4) \leq 3$ .

**Subcase 1.1.**  $(b(w_4) - 1, b(w_3)) \notin A$ .

First, we have  $m > b(w_3)$ . Let  $P_2(b(w_3), w_4) = (w_4, y, b(w_3))$ . If  $y \in W$ , then  $a(y) \geq b(w_3)$ . If  $a(y) \geq b(w_4) + 2$ , then there is a  $P'_k(a, b) = (1, \dots, b(w_4), w_4, y, b(w_4) + 2, \dots, a(y), \dots, k)$ . This contradicts  $(*)$ . Hence  $a(y) \leq b(w_4) + 1$ . Since  $a(y) \geq b(w_3) \geq b(w_4) + 1$ , we have  $a(y) = b(w_3) = b(w_4) + 1$ . By Lemma 1, there exists an  $x \in O(1) - \{2, k\}$ . Obviously,  $x \notin W$ . And  $x \notin \{3, \dots, b(w_4) - 1\}$  by Lemma 5. So  $x \geq b(w_4)$ .

(a)  $x = b(w_4)$ . Since  $x \geq 3$  and  $a(w_4) = b(w_4) - 1 \geq 2$ , there is a  $P'_k(a, b) = (1, x = b(w_4), \dots, m-1, w_4, 2, \dots, b(w_4) - 1, m, \dots, k)$ . This contradicts  $(*)$ .

(b)  $x = b(w_4) + 1$ . Note that  $m-1 > b(w_3) - 1 = b(w_4)$  and  $a(y) = b(w_4) + 1 = a(w_4) + 2 \geq 3$ . If  $a(w_4) > 1$ , then there is a  $P'_k(a, b) = (1, x = b(w_3), \dots, m-1, w_4, y, 2, \dots, a(w_4) = b(w_4) - 1, m, \dots, k)$ . This contradicts  $(*)$ . Hence  $a(w_4) = 1$ . Thus we have  $I(k) \subseteq \{1, b(w_3) - 1 = 2, k - 1\}$  by Lemma 5 (b). And then  $2 \rightarrow k$  by Lemma 1. Hence there is a  $P'_k(a, b) = (1, x = b(w_3), \dots, k-1, w_3, b(w_4) = 2, k)$ . This contradicts  $(*)$  too.

(c)  $x > b(w_3) = b(w_4) + 1$ . Since  $x < k$  and  $1 \rightarrow x$ , there is no  $j_0 \in \{2, \dots, b(w_4) - 1\}$  such that  $j_0 \rightarrow k$  by Lemma 4. Then  $I(k) \subseteq \{1, b(w_4), k-1\}$  by Lemma 5 (b). Hence  $I(k) = \{1, b(w_4), k-1\}$  by Lemma 1. That is,  $b(w_4) \rightarrow k$ . If  $k = 7$ , there is a  $P'_k(a, b) = (1, x, w_4, y, b(w_3), w_3, b(w_4), k)$ . This contradicts  $(*)$ . Hence  $k > 7$ . There are two distinct vertices  $i, j \in P'_{k-1} - \{1, a(w_4), b(w_4) = a(w_4) + 1, b(w_3) = a(w_4) + 2, x, k\}$ . Using two arcs  $(1, x)$ ,  $(b(w_4), k)$  and  $w_3, w_4, y$ , then there is always a  $P'_k(a, b)$  in  $T$ . e.g.,  $1 < i < a(w_4)$  and  $b(w_3) < j < x$ , then there is a  $P'_k(a, b) = (1, x, \dots, k-1, w_4, y, b(w_3), \dots, x-2, w_3, 3, \dots, a(w_4), b(w_4), k)$ . These contradict  $(*)$ .

Hence  $y \notin W$  and  $1 \leq y \leq a(w_4) = b(w_4) - 1$ . Since  $(b(w_4) - 1, b(w_3)) \notin A$ , we have  $y < b(w_4) - 1$ . Now, there are two arcs  $(y, b(w_3))$  and  $(b(w_4) - 1, m)$  in  $T$  with  $y < b(w_4) - 1 < b(w_3) < m$ . This contradicts  $(*)$  by Lemma 4.

**Subcase 1.2**  $(b(w_4) - 1, b(w_3)) \in A$ .

Since  $b(w_3) - b(w_4) \leq 3$  and  $k \geq 7$ , we have  $(b(w_4) - 1, b(w_3)) \neq (a, b)$ . There exists an arc  $(u, v)$  such that  $(u, v)$  and  $(b(w_4) - 1, b(w_3))$  are cis-crosswise arcs by Lemma 6.

Suppose  $u < b(w_4) - 1 < v < b(w_3)$ . For  $v = b(w_4)$  or  $b(w_4) + 1$  or  $b(w_4) + 2$ , there exists a  $P'_k(a, b)$  in  $T$  respectively. e.g., we assume  $v = b(w_4) + 1$ . If  $b(w_3) = b(w_4) + 3$ , then there is a  $P'_k(a, b) = (1, \dots, u, v = b(w_4) + 1, w_4, w_5, w_3, u+1, \dots, b(w_4) - 1, b(w_3), \dots, k)$ . If  $b(w_3) = b(w_4) + 2$ , then  $v = b(w_3) - 1 = a(w_3)$ . Since  $w_4 \rightarrow w_5$  and  $w_4 \rightarrow a(w_4), w_5$  and  $a(w_4)$  are adjacent. By the definition of  $b(w_4) = b(w_0)$ , we have  $w_5 \rightarrow a(w_4)$ . Hence  $a(w_5) \geq a(w_4) > u$  and  $w_5 \rightarrow u+1$ . Thus there is a  $P'_k(a, b) = (1, \dots, u, v = b(w_4) + 1, w_4, w_5, u+1, \dots, b(w_4) - 1, b(w_3) = b(w_4) + 2, \dots, k)$ . These contradict  $(*)$ .

Using an analogous method, if  $b(w_4) - 1 < u < b(w_3) < v$ , then there is also a  $P'_k(a, b)$  in  $T$ . This contradicts  $(*)$ .

Therefore no vertex of  $W_3$  dominates any vertex of  $W_4$ . We have that  $W_4 \rightarrow W_3$  since  $T[W]$  is a tournament.

**Case 2.**  $W_4 \rightarrow W_3$ .

We choose  $w_3 \in W_3, w_4 \in W$ , such that  $b(w_3) = \max\{b(w) \mid w \in W_3\}$ . Thus  $w_4 \rightarrow w_3$ . Since  $b(w_3) > b(w_4)$ , there exists a  $P'_2(w_4, w_3) = (w_4, w_6, w_3)$  with  $w_6 \in W$ . Now, we have the following claims.

(4.1)  $b(w_4) \leq a(w_3) \leq b(w_4) + 1$ .

Let  $P_2(w_4, w_3) = (w_3, y, w_4)$ . Since  $W_4 \rightarrow W_3$ , we have  $y \notin W$  and  $y \in P'_{k-1}$ . Thus  $b(w_4) \leq y \leq a(w_3)$ . If  $a(w_3) - 2 \geq b(w_4)$ , then there is a  $P'_k(a, b) = (1, 2, \dots, a(w_3) - 2, w_4, w_3, a(w_3), \dots, k)$ . This contradicts  $(*)$ . Hence  $a(w_3) \leq b(w_4) + 1$ .

(4.2)  $(a(w_4), b(w_3)) \in A$ .

Let  $P_2(b(w_3), w_4) = (w_4, u, b(w_3))$  and  $P_2(w_3, a(w_4)) = (a(w_4), m, w_3)$ . By the choice of  $w_3$  and  $w_4$ , we have  $u, m \notin W$ . Then  $u \leq a(w_4)$  and  $b(w_3) \leq m$ . If  $u < a(w_4)$  and  $b(w_3) < m$ , then

(a)  $a(w_3) \geq a(w_4) + 2$ . Since  $b(w_3) - 1 \geq b(w_4)$ , there is a  $P'_k(a, b) = (1, \dots, u, b(w_3), \dots, m-1, w_3, a(w_4) + 2, \dots, a(w_3), \dots, b(w_3) - 1, w_4, u + 1, \dots, a(w_4), m, \dots, k)$ ;

(b)  $b(w_4) \leq b(w_3) - 2$ . There is a  $P'_k(a, b) = (1, \dots, u, b(w_3), \dots, m-1, w_3, a(w_4) + 1, \dots, b(w_4), \dots, b(w_3) - 2, w_4, u + 1, \dots, a(w_4), m, \dots, k)$ ;

(c)  $a(w_3) \leq a(w_4) + 1$  and  $b(w_4) \geq b(w_3) - 1$ . Since  $a(w_3) \geq b(w_4) \geq a(w_4) + 1$  and  $b(w_4) \leq b(w_3) - 1$ , we have  $a(w_3) = b(w_4) = a(w_4) + 1 = b(w_3) - 1$ . Thus there is a  $P'_k(a, b) = (1, \dots, u, b(w_3), \dots, m-1, w_4, w_3, u +$

$1, \dots, b(w_3) - 2 = a(w_4), m, \dots, k$ .

These contradict  $(\star)$ . So  $u = a(w_4)$  or  $m = b(w_3)$ . Thus (4. 2) is valid.

(4.3)  $a(w_3) = b(w_3) - 1$  and  $a(w_4) = b(w_4) - 1$ .

If  $a(w_3) < b(w_3) - 1$ , then  $i$  and  $w_3$  are nonadjacent for each  $i \in \{a(w_3) + 1, \dots, b(w_3) - 1\}$ . Since  $\{a(w_3) + 1, \dots, b(w_3) - 1, k\} \subseteq I(w_4)$ ,  $k$  and  $j$  are adjacent for each  $j \in \{a(w_3) + 1, \dots, b(w_3) - 1\}$ . Since the definition of local tournaments and  $k \rightarrow w_3$ , we have  $\{a(w_3) + 1, \dots, b(w_3) - 1\} \rightarrow k$ . If  $b(w_3) < k$ , then there is a  $P'_k(a, b) = (1, \dots, a(w_4), b(w_3), \dots, k - 1, w_3, a(w_4) + 1, \dots, b(w_3) - 1, k)$  by (4.2) and  $a(w_4) + 1 \leq b(w_4) \leq a(w_3)$ . This contradicts  $(\star)$ . So  $b(w_3) = k$ . Since  $1, k - 1 \in I(k)$ ,  $1$  and  $k - 1 = b(w_3) - 1$  are adjacent, and then  $1 \rightarrow b(w_3) - 1$  by  $w_3 \rightarrow 1$ . Now, we consider the following two subcases:

(a)  $a(w_3) < b(w_3) - 2 = k - 2$ .

If  $a(w_3) \geq 3$ , then there is a  $P'_k(a, b) = (1, b(w_3) - 1, w_4, w_3, 3, \dots, b(w_3) - 2, k)$ . This contradicts  $(\star)$ . Hence  $a(w_3) \leq 2$ . Since  $a(w_3) \geq b(w_4) > a(w_4) \geq 1$ , we have  $a(w_3) = 2$ . Then  $b(w_3) - 3 > a(w_3)$  by  $k \geq 7$ . Hence  $b(w_3) - 3 \rightarrow k$  and there is a  $P'_k(a, b) = (1, b(w_3) - 1 = k - 1, w_4, w_3, a(w_3) = 2, \dots, k - 3 = b(w_3) - 3, k)$ . This also contradicts  $(\star)$ .

(b)  $a(w_3) = b(w_3) - 2 = k - 2$ .

Since  $k \geq 7$ , we have  $a(w_3) \geq 5$  and  $b(w_4) \geq a(w_3) - 1 \geq b(w_4) - 1 \geq 4$  by (4.1). If  $b(w_4) = a(w_4) + 1$ , then  $a(w_4) \geq 3$ . When  $a(w_3) = b(w_4) + 1$ , there is a  $P'_k(a, b) = (1, b(w_3) - 1 = k - 1, w_4, w_3, 2, \dots, k - 4 = a(w_4), b(w_3) = k)$  by (4.2). When  $a(w_3) = b(w_4)$ , there is a  $P'_k(a, b) = (1, b(w_3) - 1 = k - 1, w_4, w_3, 2, \dots, k - 3 = a(w_4), b(w_3) = k)$ . These contradict  $(\star)$ . So  $b(w_4) \geq a(w_4) + 2$ . Since  $a(w_4) \rightarrow b(w_3) = k$  and  $a(w_4) \rightarrow a(w_4) + 1, k$  and  $a(w_4) + 1$  are adjacent, and then  $a(w_4) + 1 \rightarrow k$  by  $k \rightarrow w_4$  and  $w_4$  and  $a(w_4) + 1$  are nonadjacent. Similarly, we can get that  $\{a(w_4) + 1, \dots, b(w_4) - 1\} \rightarrow k$  and  $1 \rightarrow \{a(w_4) + 1, \dots, b(w_4) - 1\}$  since  $\{1, a(w_4) + 1, \dots, b(w_4) - 1\} \subseteq O'(w_3)$ . If  $b(w_4) - 2 \geq a(w_4) + 1$ , then there is a  $P'_k(a, b) = (1, b(w_4) - 1, \dots, a(w_3) = k - 2, w_4, w_3, 2, \dots, b(w_4) - 2, k)$  by (4.1) and  $b(w_4) \geq 4$ . If  $b(w_4) = a(w_4) + 2$ , then there is a  $P'_k(a, b) = (1, a(w_4) + 1, \dots, k - 1, w_4, 2, \dots, a(w_4), b(w_3) = k)$  by (4.2) and  $a(w_4) \geq 2$ . These contradict  $(\star)$ . So  $a(w_3) = b(w_3) - 1$ .

Similarly, we can prove that  $a(w_4) = b(w_4) - 1$ . (4.3) is valid.

Now, by (4.1), (4.2) and (4.3), we have that  $b(w_3) - b(w_4) \leq 2$ ,  $a(w_3) = b(w_3) - 1$ ,  $a(w_4) = b(w_4) - 1$  and  $(b(w_4) - 1, b(w_3)) \in A$ . Using an analogous proof of subcase 1.2, there is a  $P'_k(a, b)$  in  $T$ . This contradicts  $(\star)$ .

Up to now, we prove that  $b(w) = b(w')$  for any  $w, w' \in W$ . Similarly, we can prove that  $a(w) = a(w')$  for any  $w, w' \in W$ . So (4) is valid.

We denote  $a_0 = a(w)$  and  $b_0 = b(w)$  for any  $w \in W$ . Then  $O'(w) = \{1, 2, \dots, a_0\}$  and  $I'(w) = \{b_0, \dots, k\}$  for any  $w \in W$ , and then  $T[\{1, \dots, a_0\}]$  and  $T[\{b_0, b_0 + 1, \dots, k\}]$  both are tournaments. Clearly for any  $i \in \{a_0 +$

$1, \dots, b_0 - 1\}$  and any  $w \in W$ ,  $i$  and  $w$  are nonadjacent.

Now, we shall use the following lemmas and symbols.

For  $1 \leq t \leq a_0$  and  $b_0 \leq j \leq k$ , let  $R(t) = \{i \mid (t, i) \in A, b_0 \leq i \leq k\}$  and  $L(j) = \{i \mid (i, j) \in A, 1 \leq i \leq a_0\}$ . Since there exist  $P_2(w, t), P_2(j, w)$  for any  $w \in W$  and  $1 \leq t \leq a_0, b_0 \leq j \leq k$ , it is easy to check  $R(t) \neq \emptyset, L(j) \neq \emptyset$ . Hence we can define,

$\psi(t) = \max\{R(t)\}, \psi_1(t) = \min\{R(t)\}, \varphi_1(j) = \max\{L(j)\}$  and  $\varphi(j) = \min\{L(j)\}$ .

Then  $b_0 \leq \psi_1(t) \leq \psi(t) \leq k, 1 \leq \varphi(j) \leq \varphi_1(j) \leq a_0$ , and  $(t, \psi(t)), (t, \psi_1(t)), (\varphi(j), j), (\varphi_1(j), j) \in A$  for any  $1 \leq t \leq a_0$  and  $b_0 \leq j \leq k$ .

**Lemma 7.** *If there are  $\alpha < \gamma < \delta$  in  $P'_{k-1}$  such that  $1 \leq \alpha \leq a_0 - 1, \alpha + 1 < \gamma, b_0 + 1 \leq \delta$  and  $(\alpha, \gamma), (\gamma - 1, \delta) \in A$ , then  $T$  contains a  $P'_k(a, b)$  in  $T$ .*

**Proof:** Let  $\alpha, \gamma$  and  $\delta$  satisfy the condition of Lemma 7. Then there is a  $P'_k(a, b) = (1, \dots, \alpha, \gamma, \dots, \delta - 1, \alpha + 1, \dots, \gamma - 1, \delta, \dots, k)$ .  $\square$

**Lemma 8.** ([2], Corollary 3.13) *Let  $P_1 = (x_1, \dots, x_m)$  and  $P_2 = (y_1, \dots, y_t)$  with  $m \geq 2$  and  $t \geq 3$  be paths in a connected local tournament  $T$ . If there exist  $i, j$  with  $1 \leq i < j \leq m$  such that  $x_i = y_1, x_j = y_t$  and  $V(P_1) \cap (v(P_2) - \{y_1, y_t\}) = \emptyset$ . Then  $T$  has an  $(x_1, x_m)$ -path  $P$  such that  $V(P) = V(P_1) \cup V(P_2)$ .*

(5)  $b_0 = a_0 + 1$

Suppose  $b_0 > a_0 + 1$ . If  $b_0 = k$ , then  $\psi(i) = k$  for each  $i \in \{1, 2, \dots, a_0\}$ . That is,  $\{1, 2, \dots, a_0\} \rightarrow k$ . Let  $P_2(a_0, a_0 + 1) = (a_0 + 1, x, a_0)$ . Obviously,  $x \notin W$ . If  $x \in \{1, 2, \dots, a_0 - 1\}$ , then  $a_0 + 1$  and  $w$  are adjacent by  $w \rightarrow x$ . This is a contradiction. So  $x \notin \{1, 2, \dots, a_0 - 1\}$ . Similarly,  $x \notin \{a_0 + 1, \dots, b_0 - 1\}$ . Thus  $x = b_0 = k$ . i.e.,  $k = x \rightarrow a_0$ . This contradicts  $a_0 \rightarrow k$ . Hence  $b_0 \leq k - 1$ . Similarly, we have  $a_0 \geq 2$ .

Let  $P_2(a_0, a_0 + 1) = (a_0 + 1, t, a_0)$ . Using an analogous proof as above, we have  $t \notin W \cup \{1, 2, \dots, a_0 - 1, a_0 + 1, \dots, b_0 - 1\}$ . That is,  $b_0 \leq t \leq k$ . If  $t = b_0$ , then we have  $b_0 \rightarrow a_0, \varphi(b_0) < a_0$  and  $\psi(a_0) > b_0$ .  $\varphi(b_0)$  and  $b_0 - 1$  are adjacent by  $\varphi(b_0) \rightarrow b_0$  and  $b_0 - 1 \rightarrow b_0$ . Since  $b_0 - 1$  and  $w$  are nonadjacent and  $w \rightarrow \varphi(b_0)$ , we have  $\varphi(b_0) \rightarrow b_0 - 1$ . Similarly, we can obtain  $\varphi(b_0) \rightarrow \{a_0 + 1, \dots, b_0 - 1\}$ . Let  $\alpha = \varphi(b_0), \gamma = a_0 + 1$  and  $\delta = \psi(a_0)$ . Then there is a  $P'_k(a, b)$  in  $T$  by Lemma 7. This contradicts  $(\star)$ . Hence  $t > b_0$ . Similarly, letting  $P_2(b_0 = 1, b_0) = (b_0, y, b_0 - 1)$ , we have  $1 \leq y < a_0$ .

If  $b_0 > a_0 + 2$ , then  $t$  and  $a_0 + 2$  are adjacent by  $a_0 + 1 \rightarrow t$  and  $a_0 + 1 \rightarrow a_0 + 2$ . If  $t \rightarrow a_0 + 2$ , then it will deduce that  $a_0 + 2$  and  $w$  are adjacent by  $t \rightarrow w$ , a contradiction. Hence  $a_0 + 2 \rightarrow t$ . Similarly, we have  $\{a_0 + 1, \dots, b_0 - 1\} \rightarrow t$ . Let  $\alpha = y (< a_0), \gamma = b_0 - 1$  and  $\delta = t (> b_0)$ . There is a  $P'_k(a, b)$  in  $T$  by Lemma 7. This contradicts  $(\star)$ . Hence  $b_0 = a_0 + 2$ .

$a_0 + 1$  and  $t - 1$  are adjacent since  $a_0 + 1 \rightarrow t$  and  $t - 1 \rightarrow t$ . Thus  $a_0 + 1 \rightarrow t - 1$  by  $t - 1 \rightarrow w$  and  $w$  and  $a_0 + 1$  are nonadjacent. Similarly, we have

$$a_0 + 1 \rightarrow \{b_0 + 1, \dots, t - 1, t\} \quad (**)$$

Now, we consider the following four cases.

**Case 1.**  $a_0 > 2$  and  $k \geq b_0 + 2$ .

If  $\varphi(b_0) < a_0$ , letting  $\alpha = \varphi(b_0)$ ,  $\gamma = b_0$  and  $\delta = t$ , then there is a  $P'_k(a, b)$  in  $T$  by Lemma 7 and  $(b_0 - 1, t) = (a_0 + 1, t) \in A$ . This contradicts  $(*)$ . Hence  $\varphi(b_0) = a_0$ . That is,  $a_0 \rightarrow b_0$ . Since  $1, a_0 \in O(w)$ ,  $1$  and  $a_0$  are adjacent. Suppose  $(1, a_0) \in A$ . If  $\psi(a_0 - 1) > b_0$ , letting  $\alpha = 1$ ,  $\gamma = a_0$  and  $\delta = \psi(a_0 - 1)$ , then there is a  $P'_k(a, b)$  in  $T$  by Lemma 7. This contradicts  $(*)$ . So  $\psi(a_0 - 1) = b_0$ . i.e.,  $a_0 - 1 \rightarrow \psi(a_0 - 1) = b_0$ . Now, letting  $\alpha = a_0 - 1$ ,  $\gamma = b_0$  and  $\delta = t$ , there is a  $P'_k(a, b)$  in  $T$  by  $(**)$  and Lemma 7. This contradicts  $(*)$  too. Hence in the following we always assume that  $(a_0, 1) \in A$ .

$$(5.1) \{1, 2, \dots, a_0 - 1\} \rightarrow a_0 + 1 \text{ and } a_0 + 1 \rightarrow k.$$

$1 \rightarrow a_0 + 1$  since  $a_0 + 1$  and  $w$  are nonadjacent and  $1, a_0 + 1 \in O(a_0)$ . Furthermore,  $2 \rightarrow a_0 + 1$  by  $1 \rightarrow 2$ . Similarly, we have that  $\{1, 2, \dots, a_0 - 1\} \rightarrow a_0 + 1$  and  $a_0 + 1 \rightarrow k$  by  $1 \rightarrow a_0 + 1$  and  $1 \rightarrow k$ .

$$(5.2) b_0 \rightarrow 1 \text{ and } \{b_0 + 2, \dots, k\} \rightarrow b_0.$$

Since  $a_0 \rightarrow 1$  and  $a_0 \rightarrow b_0$ ,  $1$  and  $b_0$  are adjacent. If  $1 \rightarrow b_0$ , then, letting  $\alpha = 1$ ,  $\gamma = b_0$  and  $\delta = t$ , there is a  $P'_k(a, b)$  in  $T$  by  $(**)$  and Lemma 7. This contradicts  $(*)$ . Hence  $b_0 \rightarrow 1$ .

If there exists a  $j \in \{b_0 + 2, \dots, k\}$  such that  $b_0 \rightarrow j$ , then  $T$  contains a  $P'_k(a, b) = (1, \dots, a_0 - 1, a_0 + 1 = b_0 - 1, b_0 + 1, \dots, j - 1, w, a_0, b_0, j, \dots, k)$  by (5.1) and  $(**)$ . This contradicts  $(*)$ . Hence  $\{b_0 + 2, \dots, k\} \rightarrow b_0$ .

$$(5.3) a_0 = 3, b_0 = 5 \text{ and } (a_0 - 1, b_0) \notin A.$$

If  $\psi(a_0 - 1) = b_0$ , then there is a  $P'_k(a, b) = (1, \dots, a_0 - 1, \psi(a_0 - 1) = b_0, \dots, k - 1, w, a_0, a_0 + 1, k)$  by (5.1). This contradicts  $(*)$ . So  $\psi(a_0 - 1) > b_0$  and  $(a_0 - 1, b_0) \notin A$ .

By  $a_0 - 1 \rightarrow \psi(a_0 - 1) > b_0$ , Lemma 7 and  $(*)$ , we have  $a_0 \rightarrow 2$ . If  $a_0 \geq 4$ , then there is a  $P'_k(a, b) = (1, a_0 + 1, \dots, \psi(a_0 - 1) - 1, w, a_0, 2, \dots, a_0 - 1, \psi(a_0 - 1), \dots, k)$  by (5.1). This contradicts  $(*)$ . Hence  $a_0 \leq 3$ , and then  $a_0 = 3$  by  $a_0 > 2$ . Thus  $b_0 = a_0 + 2 = 5$ .

$$(5.4) k = b_0 + 2 = 7$$

Suppose  $k > b_0 + 2$ . When  $\varphi(b_0 + 1) \in \{1, 2\}$ , there is a  $P'_k(a, b) = (1, \dots, \varphi(b_0 + 1), b_0 + 1, \dots, k - 1, b_0, w, \varphi(b_0 + 1) + 1, \dots, a_0 + 1, k)$  by (5.1) and (5.2). When  $\varphi(b_0 + 1) = a_0 = 3$ , letting  $\alpha = 1$ ,  $\gamma = a_0 + 1$  and  $\delta = b_0 + 1$ , there is a  $P'_k(a, b)$  in  $T$  by Lemma 7 and (5.1). These contradict  $(*)$ . Hence  $k = b_0 + 2 = 7$  by  $k \geq b_0 + 2$ .

$$(5.5) k \rightarrow a_0$$

$a_0$  and  $k$  are adjacent since  $a_0 \rightarrow b_0$  and  $k \rightarrow b_0$  by (5.2). If  $a_0 \rightarrow k$ , letting  $\alpha = 1$ ,  $\gamma = a_0 + 1$  and  $\delta = k$ , then there is a  $P'_k(a, b)$  in  $T$  by Lemma 7. This contradicts  $(*)$ . So  $k \rightarrow a_0$ .

Now,  $1$  and  $b_0 + 1$  are adjacent since  $1, b_0 + 1 = k - 1 \in I(k)$ . We consider the following two cases.

$$(a) b_0 + 1 \rightarrow 1.$$

$a_0$  and  $b_0 + 1$  are adjacent by  $a_0 \rightarrow 1$ . If  $a_0 \rightarrow b_0 + 1$ , then there is a  $P'_k(a, b) = (1, a_0 + 1, b_0, w, 2, a_0 = 3, b_0 + 1, k)$ . This contradicts  $(*)$ . Hence  $b_0 + 1 \rightarrow a_0$ .

Let  $P'_2(b_0, b_0 + 1) = (b_0 u, b_0 + 1)$ . Obviously,  $u \notin W$ . Since  $b_0 + 1 \rightarrow 1$ ,  $a_0 \rightarrow b_0$ ,  $a_0 + 1 \rightarrow b_0$  and  $b_0 + 1 \rightarrow k$ , we have  $u = 2$ . i.e.,  $b_0 \rightarrow u = 2 \rightarrow b_0 + 1$ . Let  $P'_2(1, 2) = (1, z, 2)$ . Obviously,  $z \notin W$ . Since  $a_0 \rightarrow 1$ ,  $2 \rightarrow a_0 + 1$  by (5.1),  $b_0 \rightarrow 1$  by (5.2) and  $b_0 + 1 \rightarrow 1$ , we have  $z = k$ . i.e.,  $k = z \rightarrow 2$ . Suppose  $|W| > 1$ . Clearly  $T$  contains a  $P'_k(a, b)$ . This contradicts  $(*)$ . Hence  $|W| = 1$ .

Now, by  $a_0 + 1 \rightarrow b_0 + 1$ ,  $a_0 \rightarrow b_0$ ,  $a_0 \rightarrow 1$ , (5.1)~(5.5) and (a), we have that  $T \simeq D'_8$ . This contradicts the assumption of the Theorem.

$$(b) 1 \rightarrow b_0 + 1.$$

Since  $1 \rightarrow 2$  and  $1 \rightarrow b_0 + 1$ , we have  $2$  and  $b_0 + 1$  are adjacent. If  $2 \rightarrow b_0 + 1$ , then  $2$  and  $b_0$  are adjacent and  $b_0 \rightarrow 2 = a_0 - 1$  by (5.3). Thus there is a  $P'_k(a, b) = (1, b_0 + 1, w, a_0, b_0, 2 = a_0 - 1, a_0 + 1, k)$  by (5.1) and (5.2). This contradicts  $(*)$ . So  $b_0 + 1 \rightarrow 2$ .

Let  $P'_2(b_0 + 1, k) = (b_0 + 1, y, k)$ . Obviously,  $y \notin W$ . Note that  $1 \rightarrow b_0 + 1$ ,  $k \rightarrow a_0$  by (5.5),  $a_0 + 1 \rightarrow b_0 + 1$  by  $(**)$  and  $b_0 \rightarrow b_0 + 1$ . We have  $y = 2$ . i.e.,  $2 \rightarrow k$ .

We easily check that  $|W| = 1$  and  $3$  and  $6$ ,  $2$  and  $5$  are nonadjacent. Otherwise,  $T$  contains a  $P'_k(a, b)$ . e.g.,  $(3, 6) \in A$ , there is a  $P'_k(a, b) = (1, a_0 + 1, b_0, w, 2, 3, 6 = b_0 + 1, k)$  by (5.1). These contradict  $(*)$ .

Now, by  $a_0 + 1 \rightarrow b_0 + 1$ ,  $a_0 \rightarrow b_0$ ,  $a_0 \rightarrow 1$ , (5.1)~(5.5) and (b), we have that  $T \simeq D_8$ . This contradicts the assumption of the Theorem.

**Case 2.**  $a_0 > 2$  and  $k = b_0 + 1$ .

Since  $b_0 < t \leq k$ , we have  $t = k = b_0 + 1$ . i.e.,  $b_0 - 1 = a_0 + 1 \rightarrow t = k$ . Since  $k = b_0 + 1 = a_0 + 3 \geq 7$ , we have  $a_0 \geq 4$ . If there exists a  $j_0 \in \{1, 2, \dots, a_0 - 1\}$  such that  $j_0 \rightarrow b_0$ , letting  $\alpha = j_0$ ,  $\gamma = b_0$  and  $\delta = k$ , then there is a  $P'_k(a, b)$  by Lemma 7. This contradicts  $(*)$ . Hence  $(j, b_0) \notin A$  for each  $j \in \{1, 2, \dots, a_0 - 1\}$ . Then  $I(b_0) \subseteq \{a_0, a_0 + 1 = b_0 - 1\}$  by  $b_0 \rightarrow W$ . This contradicts Lemma 1.

**Case 3.**  $a_0 = 2$  and  $k \geq b_0 + 2$ .

Consider the converse  $\overleftarrow{T}$  of  $T$ , thus we change case 3 in  $T$  for case 2 in  $T$ . So this case is impossible.

Case 4.  $a_0 = 2$  and  $k = b_0 + 1$ .

In this case  $k = b_0 + 1 = a_0 + 2 + 1 = 5$ , this contradicts  $k \geq 7$ .

Up to now, we have proved that  $b_0 = a_0 + 1$ . (5) is valid.

(6) Under the condition  $b_0 = a_0 + 1$ , we can obtain the following claims.

For convenience, let  $s = a_0 + 1 = b_0$ . Then  $a_0 = s - 1$ ,  $1 < s \leq k$  and  $T[\{1, 2, \dots, s-1\}]$  and  $T[\{s, \dots, k\}]$  both are tournaments.

(6.1)  $3 \leq s \leq k - 1$

$T$  is a  $T_0$ -type digraph when  $s = 2$  or  $k$ . e.g.,  $s = 2$ , then  $O'(w) = \{1\}$  and  $I'(w) = \{s = 2, \dots, k\}$  for any  $w \in W$ . Hence  $\varphi(i) = 1$  for each  $i \in I'(W)$ . i.e.,  $1 \rightarrow \{2, \dots, k\}$ . Let  $T'_0 = T[\{2, \dots, k\}]$ ,  $T''_0 = T[W]$  and  $v_0 = 1$ . Thus  $T'_0 \rightarrow T''_0 \rightarrow v_0 \rightarrow T'_0$ . Since  $T'_0$  and  $T''_0$  both are tournaments,  $T \simeq T_0$ -type digraph. This contradicts the assumption of the Theorem. So  $3 \leq s \leq k - 1$ .

(6.2)  $\varphi(s) < s - 1$ ,  $\psi(s - 1) > s$  can not hold simultaneously.

Suppose  $\varphi(s) < s - 1$ ,  $\psi(s - 1) > s$ . We may choose  $\alpha = \varphi(s)$ ,  $\gamma = s$  and  $\delta = \psi(s - 1)$ . Then there is a  $P'_k(a, b)$  in  $T$  by Lemma 7. This contradicts  $(*)$ . Hence we have  $\varphi(s) = s - 1$  or  $\psi(s - 1) = s$ . We may assume, without loss of generality,  $\psi(s - 1) = s$ . Otherwise, we consider the converse  $\bar{T}$  of  $T$ . Then

$$1 \leq \varphi(j) \leq \varphi_1(j) \leq s - 2 \quad (***)$$

for each  $j \in \{s + 1, \dots, k\}$  by the definition of  $\psi(s - 1)$ . We may define  $\tilde{n} = \max\{\varphi_1(j) \mid s + 1 \leq j \leq k\}$ ,  $\tilde{m} = \min\{j \mid \varphi_1(j) = \tilde{n}, s + 1 \leq j \leq k\}$ . Then  $\tilde{n} \leq s - 2$ ,  $s < \tilde{m}$ ,  $(\tilde{n}, \tilde{m}) = (\varphi_1(\tilde{m}), \tilde{m}) \in A$  and  $(\tilde{n}, \tilde{m}) \neq (1, k)$ . In fact, if  $(\tilde{n}, \tilde{m}) = (1, k)$  then every vertex in  $\{1, 2, \dots, s - 2\}$  does not dominate every vertex in  $\{s + 1, \dots, k\}$  except for an arc  $(1, k)$ . If  $k > s + 1$ , then  $\varphi(s + 1) \leq s - 2$  by  $(***)$ . i.e.,  $\varphi(s + 1) \rightarrow s + 1$ , a contradiction. So  $k = s + 1$ . Since  $k \geq 7$ , we have  $s \geq 6$ . Since  $(i, k) \notin A$  for each  $i \in \{2, \dots, s - 2\}$  and  $\psi(s - 1) = s$ , we have  $I(k) \subseteq \{1, k - 1 = s\}$ . This contradicts Lemma 1. Hence  $(\tilde{n}, \tilde{m}) \neq (1, k) = (a, b)$ .

(6.3) (1°) For each  $j \in \{\tilde{n} + 1, \dots, s - 1\}$  and  $i \in \{s + 1, \dots, k\}$ , we have  $(j, i) \notin A$ ;

(2°)  $\{\tilde{n} + 1, \dots, s - 1\} \rightarrow s$ ;

(3°)  $\tilde{n} + 1 \rightarrow \{1, 2, \dots, \tilde{n} - 1\}$  as  $\tilde{n} \geq 2$ .

By the definition of  $\tilde{n}$  and  $\psi(s - 1) = s$ , we easily check that (1°) and (2°) are valid. By  $\tilde{n} \rightarrow \tilde{m}$ , Lemma 7 and  $(*)$ , we have  $\tilde{n} + 1 \rightarrow \{1, 2, \dots, \tilde{n} - 1\}$ .

(6.4)  $k \rightarrow \{s - 1, s, \dots, k - 2\}$  as  $k \geq s + 2$ .

If there exists a  $j_0 \in \{s, \dots, k - 2\}$  such that  $j_0 \rightarrow k$ , letting  $\alpha = \varphi(j_0 + 1)$ ,  $\gamma = j_0 + 1$  and  $\delta = k$ , then there is a  $P'_k(a, b)$  in  $T$  by Lemma 7 and  $\varphi(j_0 + 1) \leq s - 2$ . This contradicts  $(*)$ . Hence  $k \rightarrow \{s, \dots, k - 2\}$ . Since

$k \rightarrow s$  and  $s - 1 \rightarrow s$ , we have  $k$  and  $s - 1$  are adjacent and  $k \rightarrow s - 1$  by (6.3).

(6.5)  $\{s + 2, \dots, k - 1, k\} \rightarrow \{\tilde{n} + 1, \dots, s\}$  if  $k \geq s + 2$ ;  $s + 1 \rightarrow \{\tilde{n} + 1, \dots, s - 1\}$  if  $k \geq s + 3$ .

$k$  and  $s - 2$  are adjacent since  $k \rightarrow s - 1$  by (6.4). If  $s - 2 \geq \tilde{n} + 1$ , then  $k \rightarrow s - 2$  by (6.3) (1°). Similarly, we have  $k \rightarrow s - 3, \dots, k \rightarrow \tilde{n} + 1$ . Thus  $k \rightarrow \{\tilde{n} + 1, \dots, s - 1, s, \dots, k - 2\}$  by (6.4). Hence  $T[\{\tilde{n} + 1, \dots, s - 1, s, \dots, k - 2, k\}]$  is a tournament. Then by (6.3) we have  $\{s + 2, \dots, k - 2, k\} \rightarrow \{\tilde{n} + 1, \dots, s - 1\}$  when  $k \geq s + 2$ , and  $s + 1 \rightarrow \{\tilde{n} + 1, \dots, s - 1\}$  when  $k \geq s + 3$ . Since  $\varphi(s + 1) \leq s - 2$ , we have  $\{s + 2, \dots, k - 1, k\} \rightarrow s$  by Lemma 7 and  $(*)$ . Since  $s - 1 \rightarrow s$  and  $k - 1 \rightarrow s$ , we have  $k - 1 \rightarrow s - 1$  by (6.3). Similarly,  $k - 1 \rightarrow \{\tilde{n} + 1, \dots, s - 1, s\}$ . Hence (6.5) is valid.

(6.6)  $1 < \varphi_1(k) \leq \tilde{n}$

Note that  $I(k) \subseteq \{1, 2, \dots, \tilde{n}, k - 1\}$  by (6.3) and (6.4). Then there exists an  $i_0 \in I(k) - \{1, k - 1\}$  with  $1 < i_0 \leq \tilde{n}$  by Lemma 1. Hence  $1 < i_0 \leq \varphi_1(k) \leq \tilde{n}$ .

In the following we consider two cases.

Case 1.  $O(1) \cap \{s, s + 1, \dots, k - 1\} \neq \emptyset$ .

Let  $p = \max\{j \mid O(1) \cap \{s, s + 1, \dots, k - 1\}\}$ . Then  $s \leq p \leq k - 1$ .

Subcase 1.1  $|W| \geq 2$ . (let  $w, w' \in W$  and  $w \neq w'$ ).

(6.7)  $p = s$ . That is,  $1 \rightarrow s$  and  $(1, j) \notin A$  for each  $j \in \{s + 1, \dots, k - 1\}$ .

Suppose  $p > s$ . Since  $k \geq 7$ , there exists an  $i \in P'_{k-1} - \{1, \varphi_1(k), s - 1, s, p, k\}$ . If  $1 < i < \varphi_1(k)$ , then there is a  $P'_k(a, b) = (1, p, \dots, k - 1, w, \varphi_1(k) + 1, \dots, p - 1, w', 3, \dots, \varphi_1(k), k)$ . Similarly,  $T$  contains a  $P'_k(a, b)$  when  $\varphi_1(k) < i < s - 1$  or  $s < i < p$  or  $p < i < k$ . These contradict  $(*)$ . Hence (6.7) is valid.

(6.8)  $T[V_1]$  is a strong tournament, where  $V_1 = \{\tilde{n} + 1, \dots, s - 1\}$ .

Since  $V_1 \subseteq O(w)$ ,  $T[V_1]$  is a tournament. If  $T[V_1]$  is not strong, then  $|V_1| \geq 2$  and  $\tilde{n} + 1 \rightarrow s - 1$ . Let  $P_2(\tilde{n} + 1, s - 1) = (s - 1, q, \tilde{n} + 1)$  in  $T$ . Obviously,  $q \notin W$ . Since  $T[V_1]$  is not strong, we have  $q \notin V_1$ .  $q \notin \{s, s + 1, \dots, k\}$  and  $q \notin \{1, 2, \dots, \tilde{n} - 1\}$  by (6.3). Hence  $q = \tilde{n}$  and  $s - 1 \rightarrow q = \tilde{n}$ . Let  $P_1 = (1, s, \dots, \tilde{m} - 1, w, 2, \dots, \tilde{n}, \tilde{m}, \dots, k)$  and  $P_2 = (w, \tilde{n} + 1, \dots, s - 1, \tilde{n})$ . Then by Lemma 8 there is a  $P'_k(a, b)$  in  $T$ . This contradicts  $(*)$ . So  $T[V_1]$  is a strong tournament.

(6.9)  $\tilde{n} = 2$  and  $2 \rightarrow k$ .

If  $\tilde{n} > 2$ , then  $\tilde{n} + 1 \rightarrow 2$  by (6.3). We may assume that  $(\tilde{n} + 1, h, \dots, \tilde{n} + 1)$  is a Hamiltonian cycle in  $T[V_1]$  by (6.8). Thus there is a  $P'_k(a, b) = (1, s, \dots, \tilde{m} - 1, w, h, \dots, \tilde{n} + 1, 2, \dots, \tilde{n}, \tilde{m}, \dots, k)$ . This contradicts  $(*)$ . So  $\tilde{n} \leq 2$ . Thus  $\tilde{n} = 2$  and  $\varphi_1(k) = 2$  by (6.6). So  $2 \rightarrow k$ .

(6.10)  $k = s + 1$ .

Suppose  $k-1 \geq s+1$ . If there exists a  $j_0 \in \{s, \dots, k-3\}$  such that  $j_0 \rightarrow k-1$ , then, letting  $\alpha = \varphi(j_0+1)$ ,  $\gamma = j_0+1$  and  $\delta = k-1$ ,  $T$  contains a  $P'_k(a, b)$  in  $T$  by (6.2) and Lemma 7. This contradicts  $(\star)$ . Hence  $k-1 \rightarrow \{s, \dots, k-3\}$ . By (6.3), (6.7) and (6.9),  $I(k-1) \subseteq \{\tilde{n} = 2, k-2\}$ . This contradicts Lemma 1. So  $k = s+1$ .

$$(6.11) \quad O(1) = \{2, s = k-1, k\}$$

Otherwise, there exists a  $y \in O(1) - \{2, s = k-1, k\}$ . Then  $y \in V_1$ . Let  $(y, \dots, h, y)$  be a Hamiltonian cycle in  $T[V_1]$ . Then there is a  $P'_k(a, b) = (1, y, \dots, h, s, w, 2 = \varphi_1(k), k)$  by (6.3). This contradicts  $(\star)$ . Hence  $O(1) = \{2, s = k-1, k\}$ .

$$(6.12) \quad 2 \rightarrow s.$$

Let  $P'_2(1, s) = (1, z, s)$ , then  $z \in O(1) = \{2, s = k-1, k\}$ . Since  $s = k-1 \rightarrow k$ , we have  $z = 2$ . Hence  $2 = z \rightarrow s$ .

So far, since  $O(1) = \{2, s = k-1, k\}$ ,  $2 \rightarrow s$  and  $2 \rightarrow k$  by (6.9), there exists no  $P'_2(1, 2)$  in  $T$ . This contradicts the assumption of the Theorem.

**Subcase 1.2**  $|W| = 1$ .

Using an analogous method of subcase 1.1, we can get  $p \neq s$ . Hence we may assume that  $s+1 \leq p \leq k-1$ .

$$(6.13) \quad k = s+2 \text{ and } p = s+1.$$

If  $k \geq s+3$ , then  $k-1 \rightarrow \{\tilde{n}+1, \dots, s-2, s-1, s\}$  by (6.5). Since  $\tilde{n}+2 \leq s$  by the definition of  $\tilde{n}$ , we have  $k-1 \rightarrow \tilde{n}+2$ . If  $\tilde{n} \geq 3$ , then  $\tilde{n}+1 \rightarrow 2$  by (6.3). Thus there is a  $P'_k(a, b) = (1, p, \dots, k-1, \tilde{n}+2, \dots, p-1, w, \varphi_1(k)+1, \dots, \tilde{n}+1, 2, \dots, \varphi_1(k), k)$ . This contradicts  $(\star)$ . Hence  $\tilde{n} = 2$ . And then  $2 = \varphi_1(k) \rightarrow k$  since  $2 \leq \varphi_1(k) \leq \tilde{n}$ . Thus there is a  $P'_k(a, b) = (1, p, \dots, k-1, \tilde{n}+1 = 3, \dots, p-1, w, 2 = \varphi_1(k), k)$ . This contradicts  $(\star)$ . So  $k = s+2$  and  $p = s+1$  since  $s+1 \leq p \leq k-1$ .

$$(6.14) \quad 2 \rightarrow s \text{ and } \tilde{n} \rightarrow s-1 \text{ as } \tilde{n} \geq 3.$$

Since  $\tilde{n}+1 \rightarrow s$  and  $\tilde{n}+1 \rightarrow 2$  by (6.3),  $2$  and  $s$  are adjacent. If  $s \rightarrow 2$ , then there is a  $P'_k(a, b) = (1, p = s+1, w, \varphi_1(k)+1, \dots, s, 2, \dots, \varphi_1(k), k)$ . This contradicts  $(\star)$ . So  $2 \rightarrow s$ .

If  $s-1 \rightarrow \tilde{n}$ , letting  $P_1 = (1, 2, s, \dots, \tilde{m}-1, w, 3, \dots, \tilde{n}, \tilde{m}, \dots, k)$  (Note that  $s+1 \leq \tilde{m} \leq k = s+2$ ) and  $P_2 = (w, \tilde{n}+1, \dots, s-1, \tilde{n})$ , than by Lemma 8 there is a  $P'_k(a, b)$  in  $T$ . This contradicts  $(\star)$ . Hence  $\tilde{n} \rightarrow s-1$ .

(6.15)  $T[V_1]$  is a strong tournament, where  $V_1 = \{\tilde{n}+1, \dots, s-1\}$  and  $\tilde{n} \geq 3$ .

If not, then  $|V_1| \geq 2$  and  $\tilde{n}+1 \rightarrow s-1$ . Let  $P_2(\tilde{n}+1, s-1) = (s-1, q, \tilde{n}+1)$ . Obviously,  $q \notin W$ . By (6.3),  $q \notin \{1, 2, \dots, \tilde{n}-1\} \cup \{s, s+1, s+2 = k\}$ . Since  $T[V_1]$  is not strong, we have  $q \notin V_1$ . Then  $q = \tilde{n}$ . i.e.,  $s-1 \rightarrow q = \tilde{n}$ . This contradicts (6.14).

$$(6.16) \quad V_1 \rightarrow \{1, 2\} \text{ and } s \rightarrow 1 \text{ as } \tilde{n} \geq 3.$$

Let  $(y, \dots, h, y)$  be a Hamiltonian cycle in  $T[V_1]$ . If there exists a  $y \in V_1$  such that  $1 \rightarrow y$ , then there is a  $P'_k(a, b) = (1, y, \dots, h, s, \dots, \tilde{m}-1, w, 2, \dots, \tilde{n}, \tilde{m}, \dots, k)$  by (6.3). This contradicts  $(\star)$ . So  $V_1 \rightarrow 1$ . Similarly, we have  $V_1 \rightarrow 2$ .

Since  $1 \rightarrow p = s+1$  and  $s \rightarrow s+1$ ,  $1$  and  $s$  are adjacent. If  $1 \rightarrow s$ , then there is a  $P'_k(a, b) = (1, s, \dots, \tilde{m}-1, w, \tilde{n}+1, \dots, s-1, 2, \dots, \tilde{n}, \tilde{m}, \dots, k)$ . This contradicts  $(\star)$ . So  $s \rightarrow 1$ .

$$(6.17) \quad \tilde{n} = 3.$$

In fact, if  $\tilde{n} \geq 4$ , then  $\tilde{n}+1 \rightarrow 3$  by (6.3). Let  $(h, \dots, \tilde{n}+1, h)$  be a Hamiltonian cycle in  $T[V_1]$ . There is a  $P'_k(a, b) = (1, 2, s, \dots, \tilde{m}-1, w, h, \dots, \tilde{n}+1, 3, \dots, \tilde{n}, \tilde{m}, \dots, k)$  by (6.14). This contradicts  $(\star)$ .

If  $\tilde{n} = 2$ , then  $\varphi_1(k) = 2$  and  $2 = \varphi_1(k) \rightarrow k$ . Since  $2, s+1 = p \in O(1)$ ,  $2$  and  $s+1$  are adjacent. If  $2 \rightarrow s+1$ , then by  $\tilde{n}+1 = 3$ ,  $s+1 \in O(2)$ ,  $\tilde{n}+1$  and  $s+1$  are adjacent. By (6.3) we have  $s+1 \rightarrow \tilde{n}+1$ . There is a  $P'_k(a, b) = (1, p = s+1, \tilde{n}+1, \dots, s, w, 2 = \varphi_1(k), k)$ . This contradicts  $(\star)$ . Hence  $s+1 \rightarrow 2$ . By (6.3) we have that  $I(s+1) \subseteq \{1, s\}$ . This contradicts Lemma 1. So  $\tilde{n} = 3$ .

$$(6.18) \quad 3 \rightarrow V_1.$$

Suppose there exists a  $y \in V_1$  such that  $y \rightarrow 3$ . Let  $(h, \dots, x, h)$  be a Hamiltonian cycle in  $T[V_1]$ . Then there is a  $P'_k(a, b) = (1, 2, s, \dots, \tilde{m}-1, w, h, \dots, y, 3 = \tilde{n}, \tilde{m}, \dots, k)$  by (6.14). This contradicts  $(\star)$ . So  $3 \rightarrow V_1$ .

For  $2 \leq \varphi_1(k) \leq n = 3$ , we consider the following two cases

(a)  $\varphi_1(k) = 3$ . (that is  $3 = \tilde{n} \rightarrow k$ )

Since  $\tilde{n}, s+1 = k-1 \in I(k)$ ,  $\tilde{n}$  and  $s+1$  are adjacent. If  $\tilde{n} \rightarrow s+1$ , then, by  $s+1, \tilde{n}+1 \in O(\tilde{n})$ ,  $s+1$  and  $\tilde{n}+1$  are adjacent. By (6.3) we have  $s+1 \rightarrow \tilde{n}+1$ . There is a  $P'_k(a, b) = (1, p = s+1, \tilde{n}+1, \dots, s, w, 2, 3 = \tilde{n}, k)$ . This contradicts  $(\star)$ . So  $s+1 \rightarrow \tilde{n}$ .

By (6.3) and Lemma 1, we have  $I(s+1) = \{1, 2, s\}$ . Then  $2 \rightarrow s+1$ .

Since  $s, \tilde{n} = 3 \in O(2)$  by (6.14),  $s$  and  $\tilde{n}$  are adjacent. If  $s \rightarrow \tilde{n}$ , then there is a  $P'_k(a, b) = (1, 2, s+1, w, \tilde{n}+1, \dots, s, \tilde{n}, k)$ . This contradicts  $(\star)$ . So  $3 = \tilde{n} \rightarrow s$ .

By  $(\star)$  and  $2 \rightarrow s+1$ , we have  $3 \rightarrow 1$ .

Since  $2, k \in O(1)$ ,  $2$  and  $k$  are adjacent. If  $2 \rightarrow k$ , then there is a  $P'_k(a, b) = (1, s+1, \tilde{n}, \dots, s, w, 2, k)$ . This contradicts  $(\star)$ . So  $k \rightarrow 2$ .

If there exists a  $y \in V_1$  such that  $s+1$  and  $y$  are adjacent, then  $s+1 \rightarrow y$  by (6.3). Let  $(y, \dots, h, y)$  be a Hamiltonian cycle in  $T[V_1]$ . There is a  $P'_k(a, b) = (1, 2, s+1, y, \dots, h, s, w, 3 = \tilde{n}, k)$  by (6.3). This contradicts  $(\star)$ . Hence  $y$  and  $s+1$  are nonadjacent for each  $y \in V_1$ .

So far, by (6.3)~(6.5), (6.13)~(6.18) and (a), we have that  $T \simeq D_8^s$ -type digraph. This contradicts the assumption of the Theorem.

(b)  $\varphi_1(k) < \tilde{n} = 3$ , then  $2 = \varphi_1(k) \rightarrow k$ .



By the definition of  $\varphi_1(k)$  and  $\tilde{n}$ , we have  $\tilde{m} = s+1$ . Hence  $\tilde{n} \rightarrow \tilde{m} = s+1$ . Since  $3 = \tilde{n}, k \in O(2)$ ,  $k$  and  $\tilde{n}$  are adjacent. By the definition of  $\varphi_1(k)$ , we have  $k \rightarrow \tilde{n}$ . Let  $P'_2(2, k) = (2, z, k)$ . By  $k \rightarrow \tilde{n} = 3, 1 \rightarrow 2$  and (6.5), we have  $z = s+1$ . i.e.,  $2 \rightarrow s+1$ . By (6.16),  $2 \rightarrow s+1, 2 \rightarrow k$  and  $2 \rightarrow 3$ . Hence there is no  $P'_2(1, 2)$  in  $T$ , a contradiction.

Case 2.  $O(1) \cap \{s, s+1, \dots, k-1\} = \emptyset$ .

(6.19)  $s+1 \rightarrow \tilde{n}+1$  if  $k = s+2$ .

Since  $s+1 = k-1, l \in I(k)$ ,  $1$  and  $s+1$  are adjacent. Then it must be  $s+1 \rightarrow 1$ . By (6.3) we have  $\tilde{n}+1 \rightarrow 1$ . Hence  $s+1$  and  $\tilde{n}+1$  are adjacent. Thus  $s+1 \rightarrow \tilde{n}+1$  by (6.3).

(6.20)  $\tilde{m} = s+1$

If  $\tilde{m} \geq s+2$ , then we have  $\varphi(s+1) < \tilde{n}$  by the definition of  $\tilde{m}$ .

(a) If  $k \geq s+3$ , then  $\tilde{m}-1 \rightarrow \tilde{n}+1$  by (6.5). There is a  $P'_k(a, b) = (1, \dots, \varphi(s+1), s+1, \dots, \tilde{m}-1, \tilde{n}+1, \dots, s, w, \varphi(s+1)+1, \dots, \tilde{n}, \tilde{m}, \dots, k)$ . This contradicts  $(\star)$ .

(b) If  $k = s+2$ , then  $\tilde{m} = s+2 = k$ . There is a  $P'_k(a, b) = (1, \dots, \varphi(s+1), s+1, \tilde{n}+1, \dots, s, w, \varphi(s+1)+1, \dots, \tilde{n}, \tilde{m} = k = s+2)$  by (6.19). This contradicts  $(\star)$ .

So (6.20) is valid.

(6.21) There exists an arc  $(u', v')$  in  $A$  such that  $u' < \tilde{n} < v' < \tilde{m}$ .

If there does not exist any arc  $(u', v')$  as mentioned above, then  $(1, i) \notin A$  for each  $i \in \{\tilde{n}+1, \dots, s = \tilde{m}-1\}$  and  $\psi(j) \geq \tilde{m} = s+1$  for each  $j \in \{2, \dots, \tilde{n}-1\}$ . By  $j \rightarrow \psi(j) \geq s+1$  and  $(\star)$ , we have  $(1, j+1) \notin A$  for each  $j \in \{2, \dots, \tilde{n}-1\}$ . That is,  $\{3, \dots, \tilde{n}\} \rightarrow 1$ . Thus we have that  $O(1) = \{2, k\}$  by the assumption of case 2. This contradicts Lemma 1. Hence (6.21) is valid.

Let  $A' = \{(u', v') \in A \mid u' < \tilde{n} < v' < \tilde{m}\}$ . Let  $\tilde{v} = \min\{v' \mid (u', v') \in A'\}$  and  $\tilde{u} = \max\{u' \mid (u', v') \in A'\}$ . Obviously,  $(\tilde{u}, \tilde{v}) \in A' \subset A$  and  $\tilde{u} < \tilde{n} < \tilde{v} < \tilde{m}$ . By (6.3) we have  $\tilde{v} > \tilde{n}+1$ .

(6.22)  $\{\tilde{n}+1, \dots, \tilde{v}-1\} \rightarrow \{1, 2, \dots, \tilde{n}-1\}$ .

By the definition of  $\tilde{v}$  and  $\{1, 2, \dots, \tilde{n}-1, \tilde{n}, \dots, \tilde{v}-1\} \subseteq O'(w)$ , (6.22) is valid.

(6.23)  $\tilde{u} = \tilde{n}-1$ .

If  $\tilde{u} < \tilde{n}-1$ , then  $\tilde{v}-1 \rightarrow \tilde{u}+1$  by (6.22). There is a  $P'_k(a, b) = (1, \dots, \tilde{u}, \tilde{v}, \dots, s, w, \tilde{n}+1, \dots, \tilde{v}-1, \tilde{u}+1, \dots, \tilde{n}, \tilde{m} = s+1, \dots, k)$ . This contradicts  $(\star)$ .

(6.21)  $\tilde{v} \notin \{\tilde{n}+1, \dots, s-2\}$ .

We assume that  $\tilde{v} \in \{\tilde{n}+1, \tilde{n}+2, \dots, s-2, s-1\} = V_1$ . Note that  $T[V_1]$  is a tournament. Suppose  $T[V_1]$  is strong. Let  $(h, \tilde{v}, \dots, h)$  be a Hamiltonian cycle in  $T[V_1]$ . Then there is a  $P'_k(a, b) = (1, \dots, \tilde{n}-1 = \tilde{u}, \tilde{v}, \dots, h, s, w, \tilde{n}, \tilde{m} = s+1, \dots, k)$  by (6.3). This contradicts  $(\star)$ . So  $T[V_1]$

is not strong. Let  $\hat{T}_1$  be a condensation of  $T[V_1]$ . Then  $\hat{T}_1$  is a transitive tournament (see [3]. 10.1.9). Let  $\hat{v}$  denote the dicomponent including  $\tilde{v}$  in  $T[V_1]$  and denote it in  $\hat{T}_1$  too. And let  $L$  (resp.,  $R$ ) be the set of vertices corresponding to  $I_{\hat{T}_1}(\hat{v})$  (resp.,  $O_{\hat{T}_1}(\hat{v})$ ) in  $T$ . Obviously,  $L, R$  and  $\hat{v}$  have Hamiltonian paths, denoted by  $\mu_1, \mu_2$  and  $\mu$  respectively. Since  $\hat{v}$  is strong, we may assume that  $\tilde{v}$  is a initial vertex of  $\mu$ . For  $L, R$ , and  $\mu$ , we have

(6.24.1)  $L \rightarrow R$  and  $L \rightarrow \hat{v} \rightarrow R$ . That is,  $\mu_1 \rightarrow \mu_2$  and  $\mu_1 \rightarrow \mu \rightarrow \mu_2$ .

(6.24.2) For any  $i \in L$ , we have  $i < \tilde{v}$ . Also for any  $j \in R$ , we have  $\tilde{v} < j$ .

If there exists an  $i \in L$  such that  $i > \tilde{v}$ , then we have  $(i, \tilde{v}, \dots, (i-1), i)$  and  $i \in \hat{v}$ . This is a contradiction. Similarly, for any  $j \in R$ , we have  $\tilde{v} < j$ .

(6.24.3)  $L \neq \emptyset$ .

If  $L = \emptyset$ , then there is a  $P'_k(a, b) = (1, \dots, \tilde{n}-1 = \tilde{u}, \mu, \mu_2, s, w, \tilde{n}, \tilde{m}, \dots, k)$  by (6.3). This contradicts  $(\star)$ .

(6.24.4)  $R = \emptyset$ .

In fact, if  $R \neq \emptyset$ , we have  $L \rightarrow R$ . By (6.3) and (6.22),  $P_2(L, R)$  must be  $R \rightarrow \tilde{n} \rightarrow L$ , i.e.,  $(\mu_1, \mu_2, \tilde{n})$  is a path. Hence there is a  $P'_k(a, b) = (1, \dots, \tilde{n}-1 = \tilde{u}, \mu, s, w, \mu_1, \tilde{m}, \dots, k)$  by (6.3) and (6.20). This contradicts  $(\star)$ .

(6.24.5)  $\hat{v} = \{\tilde{v}\}$ .

Suppose  $\hat{v} \neq \{\tilde{v}\}$ . Let  $\mu'$  be a Hamiltonian path in  $\hat{v} - \{\tilde{v}\}$ . By (6.3), (6.22) and (6.24.1),  $P_2(L, \hat{v})$  must be  $\hat{v} \rightarrow \tilde{n} \rightarrow L$ , i.e.,  $(\mu_1, \mu', \tilde{n})$  is a path. There is a  $P'_k(a, b) = (1, \dots, \tilde{n}-1 = \tilde{u}, \tilde{v}, s, w, \mu_1, \mu', \tilde{n}, \tilde{m}, \dots, k)$ . This contradicts  $(\star)$ .

So far, by (6.24.2), (6.24.4) and (6.24.5), we have  $\tilde{v} = s-1$ . So (6.24) is valid.

(6.25)  $\tilde{v} \notin \{s-1, s\}$ .

If  $\tilde{v} \in \{s-1, s\}$ , then  $T$  has the following properties

(6.25.1)  $\tilde{v} \rightarrow \{1, 2, \dots, \tilde{n}-2\}$  and  $s \rightarrow \{1, 2, \dots, \tilde{n}-3\}$  as  $\tilde{v} = s-1$ .

Since  $\tilde{v}-1 \rightarrow \tilde{v}$  and  $\tilde{v}-1 \rightarrow \{1, 2, \dots, \tilde{n}-1\}$  by (6.22).  $\tilde{v}$  and  $i$  are adjacent for any  $i \in \{1, 2, \dots, \tilde{n}-1\}$ . If there exists an  $i_0 \in \{1, 2, \dots, \tilde{n}-2\}$  such that  $i_0 \rightarrow \tilde{v}$ , then there is a  $P'_k(a, b) = (1, \dots, i_0, \tilde{v}, \dots, s, w, \tilde{n}+1, \dots, \tilde{v}-1, i_0+1, \dots, \tilde{n}, \tilde{m} = s+1, \dots, k)$ . This contradicts  $(\star)$ . So  $\tilde{v} \rightarrow \{1, 2, \dots, \tilde{n}-2\}$ . When  $\tilde{v} = s-1$ , by  $s-1 = \tilde{v} \rightarrow \{1, 2, \dots, \tilde{n}-2\}$  and  $s-1 \rightarrow s$ , we have that  $s$  and  $j$  are adjacent for any  $j \in \{1, 2, \dots, \tilde{n}-2\}$ . If there exists a  $j \in \{1, 2, \dots, \tilde{n}-3\}$  such that  $j \rightarrow s$ , then there is a  $P'_k(a, b) = (1, \dots, j, s, w, \tilde{n}+1, \dots, s-1 = \tilde{v}, j+1, \dots, \tilde{n}, \tilde{m} = s+1, \dots, k)$ . This contradicts  $(\star)$ . Hence  $s \rightarrow \{1, 2, \dots, \tilde{n}-3\}$  as  $\tilde{v} = s-1$ .

(6.25.2) For each  $i, j \in \{1, 2, \dots, \tilde{n}-1\}$  and  $i > j+1$ , we have  $(i, j) \in A$ , except the case of  $\tilde{v} = s-1, (\tilde{n}-2, s) \in A$  and  $(i, j) = (\tilde{n}-1, \tilde{n}-3)$ .

Suppose  $j + 1 < i < \tilde{n} - 1$  and  $j \rightarrow i$ . By  $i - 1 \leq \tilde{n} - 3$  and (6.25.1), we have  $s \rightarrow i - 1$  and  $\psi(i - 1) > s$ . Let  $\alpha = j$ ,  $\gamma = i$  and  $\delta = \psi(i - 1)$ . There is a  $P'_k(a, b)$  in  $T$  by Lemma 7. This contradicts  $(\star)$ . Hence  $(i, j) \in A$ .

Suppose  $j + 1 < i = \tilde{n} - 1$  and  $j \rightarrow i$ . (a) If  $\tilde{v} = s$ , then  $s = \tilde{v} \rightarrow \tilde{n} - 2 = i - 1$ . Hence  $\psi(i - 1) > s$ . Using an analogous proof as above,  $T$  contains a  $P'_k(a, b)$ . (b) If  $\tilde{v} = s - 1$  and  $(\tilde{n} - 2, s) \notin A$ , then  $\psi(i - 1) = \psi(\tilde{n} - 2) > s$  and  $T$  contains a  $P'_k(a, b)$ . (c) If  $\tilde{v} = s - 1$ ,  $(\tilde{n} - 2, s) \in A$  and  $j < \tilde{n} - 3$ , then  $\psi(i - 2) = \psi(\tilde{n} - 3) > s$  by (6.25.1). Furthermore  $T$  contains a  $P'_k(a, b) = (1, \dots, j, i = \tilde{n} - 1, \dots, s - 1 = \tilde{v}, \tilde{n} - 2, s, \dots, \psi(\tilde{n} - 3) - 1, w, j + 1, \dots, \tilde{n} - 3, \psi(\tilde{n} - 3), \dots, k)$  by (6.25.1). These contradict  $(\star)$ . So (6.25.2) is valid.

(6.25.3) For each  $i, j \in \{s, s + 1, \dots, k\}$  and  $i > j + 1$ , we have  $i \rightarrow j$ .

If there exist  $i, j \in \{s, s + 1, \dots, k\}$  and  $i > j + 1$  such that  $j \rightarrow i$ . By  $(\star\star\star)$ ,  $\varphi(j + 1) < s - 1$ . Let  $\alpha = \varphi(j + 1)$ ,  $\gamma = j + 1$  and  $\delta = i$ . There is a  $P'_k(a, b)$  in  $T$  by Lemma 7. This contradicts  $(\star)$ .

(6.25.4) If  $s < k - 1$ , then  $k \rightarrow \tilde{n}$ .

Since  $\tilde{n} \rightarrow \tilde{n} + 1$  and  $k \rightarrow \tilde{n} + 1$  by (6.5),  $k$  and  $\tilde{n}$  are adjacent. Suppose  $\tilde{n} \rightarrow k$ . If  $(i, k - 1) \in A$  for each  $i \in \{1, 2, \dots, \tilde{n} - 1\}$ , then there is a  $P'_k(a, b) = (1, \dots, i, k - 1, \tilde{n} + 1, \dots, k - 2, w, i + 1, \dots, \tilde{n}, k)$  by (6.5) and (6.19). This contradicts  $(\star)$ . Hence  $(i, k - 1) \notin A$ . So far, by (6.3), (6.25.3) and  $k - 1 \rightarrow W$ , we have  $I(k - 1) \subseteq \{k - 2, \tilde{n}\}$ . This contradicts Lemma 1. So  $k \rightarrow \tilde{n}$ .

(6.25.5)  $\tilde{n} = 4$ .

By (6.6) we have  $\tilde{n} \geq 2$ . Thus it is enough to consider the following three cases.

(a)  $\tilde{n} = 2$ .

$\tilde{u} = \tilde{n} - 1 = 1$  by (6.23). If  $\tilde{v} = s$ , then  $1 = \tilde{u} \rightarrow \tilde{v} = s$ . This contradicts the assumption of case 2. So  $\tilde{v} = s - 1$ . Hence  $L = \{\tilde{n} + 1, \dots, \tilde{v} - 1\} \rightarrow \tilde{v}$  by the proof of (6.24). Thus  $O(\tilde{v}) \subseteq \{s, \tilde{n} = 2\}$  by (6.3). This contradicts Lemma 1.

(b)  $\tilde{n} = 3$ .

By the assumption of case 2, (6.22) and (6.25.1), we have  $O(1) \subseteq \{2, 3 = \tilde{n}, k\}$ . Thus  $O(1) = \{2, 3 = \tilde{n}, k\}$  by Lemma 1. We have  $k \rightarrow 2$  by Lemma 7 and  $(\star)$ . Hence  $\varphi_1(k) = 3$ . i.e.,  $\tilde{n} = 3 = \varphi_1(k) \rightarrow k$ . And then  $k = s + 1$  by (6.25.4).  $O(2) \subseteq \{3 = \tilde{n}, \tilde{v}, s\}$  by  $s + 1 = k \rightarrow 2$  and (6.22). Then  $O(2) = \{3 = \tilde{n}, \tilde{v}, s\}$  and  $\tilde{v} = s - 1$  by Lemma 1. There exists a  $P'_2(\tilde{u}, \tilde{v}) = P'_2(2, \tilde{v}) = (2, z, \tilde{v})$  in  $T$ , then  $z \in O(2) = \{3, \tilde{v}, s\}$ . So  $z = 3$  and  $3 \rightarrow \tilde{v}$ .

Since  $\tilde{v} = s - 1$ , we have  $L = \{\tilde{n} + 1 = 4, \dots, \tilde{v} - 1 = s - 2\} \rightarrow \tilde{v}$  by the proof of (6.24). Then  $O(\tilde{v}) \subseteq \{1, s\}$  by (6.3). This contradicts Lemma 1.

(c)  $\tilde{n} > 4$ .

We have, by (6.22), (6.25.1), (6.25.2) and the assumption of the case

2, that  $O(1) \subseteq \{2, \tilde{n}, k\}$  and  $O(1) = \{2, \tilde{n}, k\}$  by Lemma 1. Thus  $1 \rightarrow \tilde{n}$ . Furthermore, by  $\tilde{n} - 3 > 1$  and (6.25.1) we have  $s \rightarrow \tilde{n} - 3$ . So  $\psi(\tilde{n} - 3) > s$ . There is a  $P'_k(a, b) = (1, \tilde{n}, \dots, \tilde{v} - 1, \tilde{n} - 2, \tilde{n} - 1 = \tilde{u}, \tilde{v}, \dots, \psi(\tilde{n} - 3) - 1, w, 2, \dots, \tilde{n} - 3, \psi(\tilde{n} - 3), \dots, k)$  by (6.22). This contradicts  $(\star)$ .

By (a), (b) and (c), (6.25.5) is valid.

(6.25.6)  $1 \rightarrow \tilde{n}$ .

If  $\tilde{n} \rightarrow 1$ , we have, by the assumption of case 2, (6.22), (6.25.1) and Lemma 1, that  $O(1) = \{2, 3, k\}$ . Thus by  $1 \rightarrow 3$ ,  $(\star)$  and Lemma 7, we have  $k \rightarrow 2$  and  $(2, i) \notin A$  for each  $i \in \{s + 1, \dots, k - 1\}$ . Hence  $P'_2(1, k)$  must be  $(1, 3, k)$ . And then we have  $3 \rightarrow k$ . We also have  $4 = \tilde{n} \rightarrow 2$  by Lemma 7 and  $(\star)$ . Thus  $O(2) \subseteq \{3, s\}$  by (6.22) and (6.25.1). This contradicts Lemma 1. So  $1 \rightarrow \tilde{n}$ .

(6.25.7)  $k \rightarrow 2$  and  $(3, k) \notin A$ .

If  $2 \rightarrow k$ , there is a  $P'_k(a, b) = (1, \tilde{n}, \dots, \tilde{v} - 1, 3 = \tilde{n} - 1 = \tilde{u}, \tilde{v}, \dots, k - 1, w, 2, k)$  by (6.22) and (6.25.6). This contradicts  $(\star)$ . So  $k \rightarrow 2$ . By  $1 \rightarrow \tilde{n} = 4$ ,  $(\star)$  and Lemma 7, we have  $(3, k) \notin A$ . So (6.25.7) is valid.

(6.25.8)  $s \neq k - 1$ .

If  $s = k - 1$ , then  $\tilde{m} = s + 1 = k$  by (6.20). we consider the following two cases.

(a)  $\tilde{v} = s$ . Since  $k \rightarrow 2$ , we have  $\psi(2) = s$ . i.e.,  $2 \rightarrow s = k - 1$ . There is a  $P'_k(a, b) = (1, 2, s = k - 1, w, \tilde{n} + 1, \dots, s - 1, \tilde{n} - 1 = 3, \tilde{n} = 4, m = k)$  by (6.22). This contradicts  $(\star)$ .

(b)  $\tilde{v} = s - 1$ . We have, by  $k \rightarrow 2$ , (6.22), (6.25.1) and Lemma 1, that  $O(2) = \{3, 4 = \tilde{n}, s = k - 1\}$ . Then  $P'_2(2, 3)$  must be  $(2, s, 3)$ . So  $s \rightarrow 3$  and  $\psi(3) > s$ . Thus  $\psi(3) = k$ . Let  $\alpha = 1$ ,  $\gamma = 4 = \tilde{n}$  and  $\delta = \psi(3)$ .  $T$  contains a  $P'_k(a, b)$  by (6.25.6) and Lemma 7. This contradicts  $(\star)$ .

By (a) and (b), (6.25.8) is valid.

(6.25.9)  $k - 1 \leq s$ .

If  $k - 1 > s$ , then we have, by (6.4), (6.5), (6.25.4) and (6.25.7), that  $I(k) = \{1, k - 1\}$ . This contradicts Lemma 1. So (6.25.9) is valid.

Since (6.25.8) and (6.25.9) contradict (6.1), we have (6.25) is valid.

Finally, we have  $\tilde{v} \notin \{\tilde{n} + 1, \dots, s - 1, s\}$  by (6.24) and (6.25). But it contradicts (6.20) and (6.21). On the other hand, note that  $\overleftarrow{D}_8 \simeq D'_8$ . Hence, under the condition of (6)  $b_0 = a_0 + 1$ , except  $T \simeq T_0$ - or  $D'_8$ -type digraph, there always exists a  $P'_k(a, b)$  in  $T$ .

Up to now, under the condition of the Theorem, we have exhausted all possible cases of  $T$  and deduced that there always exists a  $P'_k(a, b)$  in  $T$ . Therefore the proof of the Theorem is completed.  $\square$

### 3 Remark

Using the definition to check whether a local tournament of order  $n$  is completely strong path-connected needs  $O(n!)$  steps. But using the Theorem of this paper it only needs  $O(n^3)$  steps. Therefore from the complexity point of view, it can make a polynomial-time good algorithm.

### References

- [1] J. Bang-Jensen, Locally semicomplete digraphs. A generalization of tournaments, *J. Graph Theory* **14** (1990), 371–390.
- [2] J. Bang-Jensen, On the structure of local semicomplete digraphs, *Dis. Math*, **100** (1992), 243–265.
- [3] J.A. Bondy and U.R.S. Murty, *Graph Theory with Applications*, Amer. Elsevier, New York, 1976.
- [4] Bu Yue Hua and Zhang Ke Min, On the arc-pancyclicity of local tournaments, *Ars Combinatoria* (to appear in 1997).
- [5] Zhang Ke Min, Completely strong path-connected tournaments, *J. Combin. Theory* **B33** (1982), 166–177.
- [6] Zhang Ke Min and Wu Zheng Sheng, A necessary and sufficient condition for completely strong path-connectivity of tournaments, *Chinese Ann Math.* **4A**(3) (1983), 385–392.
- [7] Zhang Ke Min, Wu Zheng Sheng and Zhou Yuan, On the existence of a class of tournaments with completely strong path connectivity, *J. of Nanjing Univerity* **3** (1981), 303–307.

## Some Results on Packing Graphs in their Complements

T. Gangopadhyay  
XLRI Jamshedpur  
Post Box 222  
Jamshedpur 831 001  
India

**ABSTRACT.** A supergraph  $H$  of a graph  $G$  is called tree-covered if  $H - E(G)$  consists of exactly  $|V(G)|$  vertex-disjoint trees, with each tree having exactly one point in common with  $G$ . In this paper, we show that if a graph  $G$  can be packed in its complement and if  $H$  is a tree-covered supergraph of  $G$  then  $H$  itself is self-packing unless  $H$  happens to be a member of a specified class of graphs. This is a generalisation of earlier results that almost all trees and unicyclic graphs can be packed in their complements.

### 1 Introduction and definitions

The notion of packing a pair of graphs is now fairly old. According to Schuster [10], this notion first appears in 1977 and 1978 (see [1], [2], [9]). We quote the definition from Schuster [10]:

**Definition.** Let  $G$  and  $\bar{G}$  be two graphs of the same order  $p$ . A packing of  $G_1$  with  $G_2$  is an isomorphic embedding of  $G_1$  in  $\bar{G}_2$ , the complement of  $G_2$ .

We say that  $G$  is self-packing if there is a packing of  $G$  with  $G$  itself. For a self-packing graph  $G$ , let  $\sigma$  denote an injection from  $G$  into  $\bar{G}$  such that if  $uv \in E(G)$  then  $\sigma(u)\sigma(v) \in E(\bar{G})$ . Such a  $\sigma$  is called an isomorphic embedding (or an isomorphism) from  $G$  into  $\bar{G}$ . The image of  $\sigma$  in  $\bar{G}$  is denoted by  $G^*$  and is called a copy of  $G$  in  $\bar{G}$  under  $\sigma$ . In what follows, letters  $G, H$  will always denote simple undirected graphs.

Let  $G$  be a self-packing graph on  $n$  points. We say that a supergraph  $H$  of  $G$  is tree-covered if  $H - E(G)$  consists of exactly  $n$  vertex-disjoint trees