

ON SOME UNIVERSAL SUMS OF GENERALIZED POLYGONAL NUMBERS

FAN GE AND ZHI-WEI SUN

ABSTRACT. For $m = 3, 4, \dots$ those $p_m(x) = (m-2)x(x-1)/2 + x$ with $x \in \mathbb{Z}$ are called generalized m -gonal numbers. Sun [13] studied for what values of positive integers a, b, c the sum $ap_5 + bp_5 + cp_5$ is universal over \mathbb{Z} (i.e., any $n \in \mathbb{N} = \{0, 1, 2, \dots\}$ has the form $ap_5(x) + bp_5(y) + cp_5(z)$ with $x, y, z \in \mathbb{Z}$). We prove that $p_5 + bp_5 + 3p_5$ ($b = 1, 2, 3, 4, 9$) and $p_5 + 2p_5 + 6p_5$ are universal over \mathbb{Z} , as conjectured by Sun. Sun also conjectured that any $n \in \mathbb{N}$ can be written as $p_3(x) + p_5(y) + p_{11}(z)$ and $3p_3(x) + p_5(y) + p_7(z)$ with $x, y, z \in \mathbb{N}$; in contrast, we show that $p_3 + p_5 + p_{11}$ and $3p_3 + p_5 + p_7$ are universal over \mathbb{Z} . Our proofs are essentially elementary and hence suitable for general readers.

1. INTRODUCTION

For $m = 3, 4, \dots$ we set

$$(1.1) \quad p_m(x) = (m-2) \frac{x(x-1)}{2} + x.$$

Those $p_m(n)$ with $n \in \mathbb{N} = \{0, 1, 2, \dots\}$ are the well-known *m-gonal numbers* (or polygonal numbers of order m). We call those $p_m(x)$ with $x \in \mathbb{Z}$ *generalized m-gonal numbers*. Note that (generalized) 3-gonal numbers are triangular numbers and (generalized) 4-gonal numbers are squares of integers.

In 1638, Fermat asserted that each $n \in \mathbb{N}$ can be written as the sum of m polygonal numbers of order m . This was proved by Lagrange, Gauss and Cauchy in the cases $m = 4$, $m = 3$ and $m \geq 5$ respectively (see Moreno and Wagstaff [10, pp. 54-57] or Nathanson [11, Chapter 1, pp. 3-34]). The generalized pentagonal numbers play a crucial role in Euler's famous recurrence for the partition function.

For $a, b, c \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$ and $i, j, k \in \{3, 4, \dots\}$, Sun [13] called the sum $ap_i + bp_j + cp_k$ universal over \mathbb{N} (resp., over \mathbb{Z}) if for any $n \in \mathbb{N}$ the equation $n = ap_i(x) + bp_j(y) + cp_k(z)$ has solutions over \mathbb{N} (resp., over \mathbb{Z}). In 1862 Liouville (cf. [4, p. 23]) determined all those universal $ap_3 + bp_3 + cp_3$. The second author [12] initiated the determination of those universal sums

2010 *Mathematics Subject Classification*. Primary 11E25; Secondary 11B75, 11D85, 11E20, 11P32.

Key words and phrases. Polygonal numbers, representations of integers, ternary quadratic forms.

$ap_i + bp_j + cp_k$ with $\{i, j, k\} = \{3, 4\}$, and this project was completed via [12, 5, 9]. For almost universal sums $ap_i + bp_j + cp_k$ with $\{i, j, k\} \subseteq \{3, 4\}$, see [8, 1, 2].

It is known that generalized hexagonal numbers are identical with triangular numbers (cf. [6] or [13, (1.3)]).

The second author recently established the following result.

Theorem 1.1. (Sun [13, Theorem 1.1]) *Suppose that $ap_k + bp_k + cp_k$ is universal over \mathbb{Z} , where $k \in \{4, 5, 7, 8, 9, \dots\}$, $a, b, c \in \mathbb{Z}^+$ and $a \leq b \leq c$. Then $k = 5$, $a = 1$ and (b, c) is among the following 20 ordered pairs:*

$$\begin{aligned} &(1, c) \ (c \in \{1, 2, 3, 4, 5, 6, 8, 9, 10\}), \\ &(2, 2), \ (2, 3), \ (2, 4), \ (2, 6), \ (2, 8), \\ &(3, 3), \ (3, 4), \ (3, 6), \ (3, 7), \ (3, 8), \ (3, 9). \end{aligned}$$

Guy [6] realized that $p_5 + p_5 + p_5$ is universal over \mathbb{Z} , and Sun [13] proved that the sums

$$\begin{aligned} &p_5 + p_5 + 2p_5, \ p_5 + p_5 + 4p_5, \ p_5 + 2p_5 + 2p_5, \\ &p_5 + 2p_5 + 4p_5, \ p_5 + p_5 + 5p_5, \ p_5 + 3p_5 + 6p_5 \end{aligned}$$

are universal over \mathbb{Z} . So the converse of Theorem 1.1 reduces to the following conjecture of Sun.

Conjecture 1.2. (Sun [13, Remark 1.2]) *The sum $p_5 + bp_5 + cp_5$ is universal over \mathbb{Z} if the ordered pair (b, c) is among*

$$\begin{aligned} &(1, 3), \ (1, 6), \ (1, 8), \ (1, 9), \ (1, 10), \ (2, 3), \\ &(2, 6), \ (2, 8), \ (3, 3), \ (3, 4), \ (3, 7), \ (3, 8), \ (3, 9). \end{aligned}$$

Our following result confirms this conjecture for six ordered pairs (b, c) for the first time.

Theorem 1.3. *For*

$$(b, c) = (1, 3), \ (2, 3), \ (2, 6), \ (3, 3), \ (3, 4), \ (3, 9),$$

the sum $p_5 + bp_5 + cp_5$ is universal over \mathbb{Z} .

Remark. This result appeared in the initial preprint version of this paper posted to arXiv in 2009.

Sun [13] investigated those universal sums $ap_i + bp_j + cp_k$ over \mathbb{N} . By Sun [13, Conjectures 1.10 and 1.13], $p_3 + p_5 + p_{11}$ and $3p_3 + p_5 + p_7$ should be universal over \mathbb{N} . Though we cannot prove this, we are able to show the following result.

Theorem 1.4. *The sums $p_3 + p_5 + p_{11}$ and $3p_3 + p_5 + p_7$ are universal over \mathbb{Z} .*

Theorems 1.3 and 1.4 will be shown in Sections 2 and 3 respectively. Our proofs are essentially elementary and hence suitable for general readers.

2. PROOF OF THEOREM 1.3

Lemma 2.1. (Sun [13, Lemma 3.2]) *Let $w = x^2 + 3y^2 \equiv 4 \pmod{8}$ with $x, y \in \mathbb{Z}$. Then there are odd integers u and v such that $w = u^2 + 3v^2$.*

Lemma 2.2. *Let $w = x^2 + 3y^2$ with x, y odd and $3 \nmid x$. Then there are integers u and v relatively prime to 6 such that $w = u^2 + 3v^2$.*

Proof. It suffices to consider the case $3 \mid y$. Without loss of generality, we may assume that $x \not\equiv y \pmod{4}$ (otherwise we may use $-y$ instead of y). Thus $(x - y)/2$ and $(x + 3y)/2 = (x - y)/2 + 2y$ are odd. Observe that

$$(2.1) \quad x^2 + 3y^2 = \left(\frac{x + 3y}{2}\right)^2 + 3\left(\frac{x - y}{2}\right)^2.$$

As $3 \nmid x$ and $3 \mid y$, neither $(x - y)/2$ nor $(x + 3y)/2$ is divisible by 3. Therefore $u = (x + 3y)/2$ and $v = (x - y)/2$ are relatively prime to 6. This concludes the proof. \square

Lemma 2.3. (Jacobi's identity) *We have*

$$(2.2) \quad 3(x^2 + y^2 + z^2) = (x + y + z)^2 + 2\left(\frac{x + y - 2z}{2}\right)^2 + 6\left(\frac{x - y}{2}\right)^2.$$

We need to introduce some more notation. For $a, b, c \in \mathbb{Z}^+$, we set

$$E(ax^2 + by^2 + cz^2) = \{n \in \mathbb{N} : n \neq ax^2 + by^2 + cz^2 \text{ for any } x, y, z \in \mathbb{Z}\}.$$

Proof of Theorem 1.3. Let $b, c \in \mathbb{Z}^+$. For $n \in \mathbb{N}$ we have

$$\begin{aligned} n &= p_5(x) + bp_5(y) + cp_5(z) = \frac{3x^2 - x}{2} + b\frac{3y^2 - y}{2} + c\frac{3z^2 - z}{2} \\ \iff 24n + b + c + 1 &= (6x - 1)^2 + b(6y - 1)^2 + c(6z - 1)^2. \end{aligned}$$

If $w \in \mathbb{Z}$ is relatively prime to 6, then w or $-w$ is congruent to -1 modulo 6. Thus, $p_5 + bp_5 + cp_5$ is universal over \mathbb{Z} if and only if for any $n \in \mathbb{N}$ the equation $24n + b + c + 1 = x^2 + by^2 + cz^2$ has integral solutions with x, y, z relatively prime to 6.

Below we fix a nonnegative integer n .

(i) By Dickson [3, Theorem III],

$$(2.3) \quad E(x^2 + y^2 + 3z^2) = \{9^k(9l + 6) : k, l \in \mathbb{N}\}.$$

So $24n + 5 = u^2 + v^2 + 3w^2$ for some $u, v, w \in \mathbb{Z}$. As $3w^2 \not\equiv 5 \pmod{4}$, u or v is odd. Without loss of generality we assume that $2 \nmid u$. Since $v^2 + 3w^2 \equiv 5 - u^2 \equiv 4 \pmod{8}$, by Lemma 2.1 we can rewrite $v^2 + 3w^2$ as $s^2 + 3t^2$ with s, t odd. Now we have $24n + 5 = u^2 + s^2 + 3t^2$ with u, s, t odd. By $u^2 + s^2 \equiv 5 \equiv 2 \pmod{3}$, both u and s are relatively prime to 3. Applying Lemma 2.2 we can express $s^2 + 3t^2$ as $y^2 + 3z^2$ with y, z relatively prime to 6. Thus $24n + 5 = u^2 + y^2 + 3z^2$ with u, y, z relatively prime to 6. This proves the universality of $p_5 + p_5 + 3p_5$ over \mathbb{Z} .

(ii) By Dickson [3, Theorem X],

$$(2.4) \quad E(x^2 + 2y^2 + 3z^2) = \{4^k(16l + 10) : k, l \in \mathbb{N}\}.$$

So $24n + 6 = 2u^2 + v^2 + 3w^2$ for some $u, v, w \in \mathbb{Z}$. Clearly v and w have the same parity. Thus $4 \mid v^2 + 3w^2$ and hence $2u^2 \equiv 6 \pmod{4}$. So u is odd and $v^2 + 3w^2 \equiv 6 - 2u^2 \equiv 4 \pmod{8}$. By Lemma 2.1 we can rewrite $v^2 + 3w^2$ as $s^2 + 3t^2$ with s, t odd. Now we have $24n + 6 = 2u^2 + s^2 + 3t^2$ with u, s, t odd. Note that $s^2 + 2u^2 > 0$ and $s^2 + 2u^2 \equiv 0 \pmod{3}$. By [7, p. 173] or [13, Lemma 2.1], we can rewrite $s^2 + 2u^2$ as $x^2 + 2y^2$ with x and y relatively prime to 3. As $x^2 + 2y^2 = s^2 + 2u^2 \equiv 3 \pmod{8}$, both x and y are odd. By Lemma 2.2, $x^2 + 3t^2 = r^2 + 3z^2$ for some integers $r, z \in \mathbb{Z}$ relatively prime to 6. Thus $24n + 6 = r^2 + 2y^2 + 3z^2$ with r, y, z relatively prime to 6. It follows that $p_5 + 2p_5 + 3p_5$ is universal over \mathbb{Z} .

(iii) By Dickson [3, Theorem IV],

$$(2.5) \quad E(x^2 + 3y^2 + 3z^2) = \{9^k(3l + 2) : k, l \in \mathbb{N}\}.$$

So $24n + 7 = u^2 + 3v^2 + 3w^2$ for some $u, v, w \in \mathbb{Z}$. Since $u^2 \not\equiv 7 \pmod{4}$, without loss of generality we assume that $2 \nmid w$. As $u^2 + 3v^2 \equiv 7 - 3w^2 \equiv 4 \pmod{8}$, by Lemma 2.1 there are odd integers s and t such that $u^2 + 3v^2 = s^2 + 3t^2$. Thus $24n + 7 = s^2 + 3t^2 + 3w^2$ with s, t, w odd. Clearly, s is relatively prime to 6. By Lemma 2.2, $s^2 + 3t^2 = x_0^2 + 3y^2$ for some integers x_0 and y relatively prime to 6, and $x_0^2 + 3w^2 = x^2 + 3z^2$ for some integers x and z relatively prime to 6. Therefore $24n + 7 = x^2 + 3y^2 + 3z^2$ with x, y, z relatively prime to 6. This proves the universality of $p_5 + 3p_5 + 3p_5$ over \mathbb{Z} .

(iv) By [13, Theorem 1.7(iii)], $24n + 8 = u^2 + v^2 + 3w^2$ for some $u, v, w \in \mathbb{Z}$ with $2 \nmid w$. Clearly $u \not\equiv v \pmod{2}$. Without loss of generality, we assume that $u = 2r$ with $r \in \mathbb{Z}$. Since $(2r)^2 + v^2 \equiv 8 \equiv 2 \pmod{3}$, both r and v are relatively prime to 3. As v and w are odd, $v^2 + 3w^2 \equiv 4 \pmod{8}$ and hence r is odd. By Lemma 2.2, we can rewrite $v^2 + 3w^2$ as $x^2 + 3y^2$ with x and y relatively prime to 6. Note that $24n + 8 = 4r^2 + v^2 + 3w^2 = x^2 + 3y^2 + 4r^2$

with x, y, r relatively prime to 6. It follows that $p_5 + 3p_5 + 4p_5$ is universal over \mathbb{Z} .

(v) By (2.3), $24n + 13 = u^2 + v^2 + 3w^2$ for some $u, v, w \in \mathbb{Z}$. Since $3w^2 \not\equiv 13 \equiv 1 \pmod{4}$, without loss of generality we may assume that u is odd. As $v^2 + 3w^2 \equiv 13 - u^2 \equiv 4 \pmod{8}$, by Lemma 2.1 we can rewrite $v^2 + 3w^2$ as $s^2 + 3t^2$ with s and t odd. Thus $24n + 13 = u^2 + s^2 + 3t^2$ with u, s, t odd. Since $u^2 + s^2 \equiv 13 \equiv 1 \pmod{3}$, without loss of generality we may assume that $3 \nmid u$ and $s = 3r$ with $r \in \mathbb{Z}$. By Lemma 2.2, $u^2 + 3t^2 = x^2 + 3y_0^2$ for some integers x and y_0 relatively prime to 6, also $y_0^2 + 3r^2 = y^2 + 3z^2$ for some integers y and z relatively prime to 6. Thus $24n + 13 = x^2 + 3y_0^2 + 9r^2 = x^2 + 3y^2 + 9z^2$ with x, y, z relatively prime to 6. This proves the universality of $p_5 + 3p_5 + 9p_5$ over \mathbb{Z} .

(vi) By the Gauss-Legendre theorem (cf. [11, pp. 17-23]), $8n + 3 = x^2 + y^2 + z^2$ for some odd integers x, y, z . Without loss of generality we may assume that $x \not\equiv y \pmod{4}$. By Jacobi's identity (2.2), we have $3(8n + 3) = u^2 + 2v^2 + 6w^2$, where $u = x + y + z$, $v = (x + y)/2 - z$ and $w = (x - y)/2$ are odd integers. As $u^2 + 2v^2$ is a positive integer divisible by 3, by [7, p. 173] or [13, Lemma 2.1] we can write $u^2 + 2v^2 = a^2 + 2b^2$ with a and b relatively prime to 3. Since $a^2 + 2b^2 = u^2 + 2v^2 \equiv 3 \pmod{8}$, both a and b are odd. By Lemma 2.2, $b^2 + 3w^2 = c^2 + 3d^2$ for some integers c and d relatively prime to 6. Thus $24n + 9 = a^2 + 2b^2 + 6w^2 = a^2 + 2c^2 + 6d^2$ with a, c, d relatively prime to 6. It follows that $p_5 + 2p_5 + 6p_5$ is universal over \mathbb{Z} .

In view of the above, we have completed the proof of Theorem 1.3. \square

3. PROOF OF THEOREM 1.4

Proof of Theorem 1.4. (i) Let $n \in \mathbb{N}$. By part (v) in the proof of Theorem 1.3, there are integers $u, v, w \in \mathbb{Z}$ relatively prime to 6 such that

$$72n + 61 = 24(3n + 2) + 13 = 9u^2 + 3v^2 + w^2.$$

Clearly $w^2 \equiv 61 - 3v^2 \equiv 7^2 \pmod{9}$ and hence $w \equiv \pm 7 \pmod{9}$. So there are $x, y, z \in \mathbb{Z}$ such that

$$72n + 61 = 9(2x + 1)^2 + 3(6y - 1)^2 + (18z - 7)^2$$

and hence $n = p_3(x) + p_5(y) + p_{11}(z)$. (Note that $p_{11}(x) = 9(x^2 - x)/2 + x = (9x^2 - 7x)/2$.)

(ii) Let $n \in \mathbb{N}$. It is easy to see that

$$\begin{aligned} n &= 3p_3(x) + p_5(y) + p_7(z) \\ \iff 120n + 77 &= 5(3(2x + 1))^2 + 5(6y - 1)^2 + 3(10z - 3)^2. \end{aligned}$$

Suppose $120n + 77 = 5x^2 + 5y^2 + 3z^2$ for some $x, y, z \in \mathbb{Z}$ with z odd. Then $x^2 + y^2 \equiv 77 - 3z^2 \equiv 2 \pmod{4}$ and hence x and y are odd. Note that $3z^2 \equiv 77 \equiv 12 \pmod{5}$ and hence $z \equiv \pm 3 \pmod{10}$. As $5x^2 + 5y^2 \equiv 77 \equiv 5 \pmod{3}$, exactly one of x and y is divisible by 3. Thus there are $u, v, w \in \mathbb{Z}$ such that

$$120n + 77 = 5(3(2u + 1))^2 + 5(6v - 1)^2 + 3(10w - 3)^2.$$

By the above, to prove the universality of $3p_3 + p_5 + p_7$ over \mathbb{Z} , we only need to show that $120n + 77 = 5x^2 + 5y^2 + 3z^2$ for some $x, y, z \in \mathbb{Z}$ with z odd.

By (2.3), there are $u, v, w \in \mathbb{Z}$ such that $120n + 77 = u^2 + v^2 + 3w^2$. As $3w^2 \not\equiv 77 \equiv 1 \pmod{4}$, u or v is odd, say, $2 \nmid u$. As $v^2 + 3w^2 \equiv 77 - u^2 \equiv 4 \pmod{8}$, by Lemma 2.1 we may assume that v and w are odd without loss of generality.

We claim that $120n + 77 = a^2 + b^2 + 3c^2$ for some odd integers a, b, c with $c \equiv \pm 2 \pmod{5}$. This holds if $w \equiv \pm 2 \pmod{5}$. Suppose that $w \not\equiv \pm 2 \pmod{5}$. If $w \equiv \pm 1 \pmod{5}$, then $u^2 + v^2 \equiv 77 - 3w^2 \equiv -1 \pmod{5}$ and hence u or v is divisible by 5. If $w \equiv 0 \pmod{5}$, then $u^2 + v^2 \equiv 77 \equiv 2 \pmod{5}$ and hence $u^2 \equiv v^2 \equiv 1 \pmod{5}$. Without loss of generality, we assume that one of v and w is divisible by 5 and the other one is congruent to 1 or -1 modulo 5, we may also suppose that $v \not\equiv w \pmod{4}$ (otherwise we may use $-w$ instead of w). By the identity (2.1),

$$v^2 + 3w^2 = \left(\frac{v + 3w}{2}\right)^2 + 3\left(\frac{v - w}{2}\right)^2.$$

Note that both $(v - w)/2$ and $(v + 3w)/2 = (v - w)/2 + 2w$ are odd. Also, $(v - w)/2$ is congruent to 2 or -2 modulo 5. This confirms the claim.

By the above, there are odd integers $a, b, c \in \mathbb{Z}$ with $c \equiv \pm 2 \pmod{5}$ such that $120n + 77 = a^2 + b^2 + 3c^2$. Since $3c^2 \equiv 77 \pmod{5}$, we have $5 \mid a^2 + b^2$ and hence $a^2 \equiv (2b)^2 \pmod{5}$. Without loss of generality we assume that $a \equiv 2b \pmod{5}$. Then $x = (2a + b)/5$ and $y = (a - 2b)/5$ are odd integers, and

$$a^2 + b^2 = (2x + y)^2 + (x - 2y)^2 = 5(x^2 + y^2).$$

Now we have $120n + 77 = 5(x^2 + y^2) + 3c^2$ with x, y, c odd.

This concludes our proof of Theorem 1.4. \square

Acknowledgements. The authors would like to thank Dr. Hao Pan for his helpful comments. The second author was supported by the National Natural Science Foundation of China (grant 11571162).

REFERENCES

- [1] W. K. Chan and A. Haensch, *Almost universal ternary sums of squares and triangular numbers*, in: Quadratic and Higher Degree Forms, Dev. Math., Vol. 31, Springer, New York, 2013, pp. 51–62.
- [2] W. K. Chan and B.-K. Oh, *Almost universal ternary sums of triangular numbers*, Proc. Amer. Math. Soc. **137** (2009), 3553–3562.
- [3] L. E. Dickson, *Integers represented by positive ternary quadratic forms*, Bull. Amer. Math. Soc. **33** (1927), 63–77.
- [4] L. E. Dickson, History of the Theory of Numbers, Vol. II, AMS Chelsea Publ., 1999.
- [5] S. Guo, H. Pan and Z.-W. Sun, *Mixed sums of squares and triangular numbers (II)*, Integers **7** (2007), #A56, 5pp (electronic).
- [6] R. K. Guy, *Every number is expressible as the sum of how many polygonal numbers?* Amer. Math. Monthly **101** (1994), 169–172.
- [7] B. W. Jones and G. Pall, *Regular and semi-regular positive ternary quadratic forms*, Acta Math. **70** (1939), 165–191.
- [8] B. Kane and Z.-W. Sun, *On almost universal mixed sums of squares and triangular numbers*, Trans. Amer. Math. Soc. **362** (2010), 6425–6455.
- [9] B.-K. Oh and Z.-W. Sun, *Mixed sums of squares and triangular numbers (III)*, J. Number Theory **129** (2009), 964–969.
- [10] C. J. Moreno and S. S. Wagstaff, Sums of Squares of Integers, Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [11] M. B. Nathanson, Additive Number Theory: The Classical Bases, Grad. Texts in Math., vol. 164, Springer, New York, 1996.
- [12] Z.-W. Sun, *Mixed sums of squares and triangular numbers*, Acta Arith. **127** (2007), 103–113.
- [13] Z.-W. Sun, *On universal sums of polygonal numbers*, Sci. China Math. **58** (2015), 1367–1396.

(FAN GE) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627, U.S.A.

E-mail address: fange.math@gmail.com

(ZHI-WEI SUN, CORRESPONDING AUTHOR) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S REPUBLIC OF CHINA

E-mail address: zwsun@nju.edu.cn