

Preprint (August 14, 2006)

FLECK QUOTIENTS AND BERNOULLI NUMBERS

ZHI-WEI SUN

Department of Mathematics, Nanjing University
 Nanjing 210093, People's Republic of China
 zwsun@nju.edu.cn
<http://pweb.nju.edu.cn/zwsun>

ABSTRACT. Let p be a prime, and let $n > 0$ and r be integers. In 1913 Fleck showed that

$$F_p(n, r) = (-p)^{-\lfloor (n-1)/(p-1) \rfloor} \sum_{k \equiv r \pmod{p}} \binom{n}{k} (-1)^k \in \mathbb{Z}.$$

Nowadays this result plays important roles in many aspects. Recently Sun and Wan investigated $F_p(n, r) \pmod{p}$ in [SW2]. In this paper, using p -adic methods we determine $(F_p(m, r) - F_p(n, r))/(m - n)$ modulo p in terms of Bernoulli numbers, where $m > 0$ is an integer with $m \neq n$ and $m \equiv n \pmod{p(p-1)}$. Consequently, $F_p(n, r) \pmod{p^{\text{ord}_p(n)+1}}$ is determined; for example, if $n \equiv n_* \pmod{p-1}$ with $0 < n_* < p-2$ then

$$\frac{F_p(pn, 0)}{pn} \equiv \frac{n_*!}{n_* + 1} B_{p-1-n_*} \pmod{p}.$$

This yields an application to Stirling numbers of the second kind. We also study extended Fleck quotients; in particular we prove that if $a > 0$ and $l \geq 0$ are integers with $2 \leq n - l \leq p$ then

$$\frac{1}{p^{n-l}} \sum_{l < k \leq n} \binom{p^a n - d}{p^a k - d} (-1)^{pk} \binom{k-1}{l} \equiv \frac{(-1)^{l-1} n!}{l!(n-l)} B_{p-n+l} \pmod{p}$$

for all $d = 1, \dots, \max\{p^{a-2}, 1\}$.

2000 *Mathematics Subject Classification*. Primary 11B65; Secondary 05A10, 11A07, 11B68, 11B73, 11S80, 11T24.

Supported by the National Science Fund for Distinguished Young Scholars (No. 10425103) in China.

1. INTRODUCTION

Let p be a prime. In 1819 C. Babbage observed that

$$\begin{aligned} (-1)^{p-1} \binom{2p-1}{p-1} &= \prod_{k=1}^{p-1} \left(1 - \frac{2p}{k}\right) \\ &\equiv 1 - 2p \sum_{k=1}^{p-1} \frac{1}{k} = 1 - p \sum_{k=1}^{p-1} \left(\frac{1}{k} + \frac{1}{p-k}\right) \equiv 1 \pmod{p^2}. \end{aligned}$$

In 1862 J. Wolstenholme proved further that if $p > 3$ then

$$\binom{2p-1}{p-1} = \frac{1}{2} \binom{2p}{p} \equiv 1 \pmod{p^3}.$$

This is a fundamental congruence involving binomial coefficients. When $p > 3$, we also have

$$(-1)^{r+1} \binom{2p-1}{p-r-1} \equiv -2p^2 H_r^2 + 2pH_r - 1 \pmod{p^3} \quad (1.1)$$

for each $r = 1, \dots, p-1$, where $H_r = \sum_{0 < k \leq r} 1/k$; in fact,

$$\begin{aligned} &(-1)^{r+1} \binom{2p-1}{p-r-1} + \binom{2p-1}{p-1} \\ &= \sum_{s=1}^r \left((-1)^{s+1} \binom{2p-1}{p-s-1} - (-1)^s \binom{2p-1}{p-s} \right) \\ &= \sum_{s=1}^r (-1)^{s+1} \frac{2p}{p-s} \binom{2p-1}{p-s-1} = 2p \sum_{s=1}^r \frac{s+p}{s^2-p^2} \prod_{0 < k < p-s} \left(1 - \frac{2p}{k}\right) \\ &\equiv 2p \sum_{s=1}^r \left(\frac{1}{s} + \frac{p}{s^2}\right) \left(1 - 2p \left(H_{p-1} - \sum_{t=1}^s \frac{1}{p-t}\right)\right) \\ &\equiv 2p \sum_{s=1}^r \left(\frac{1}{s} + \frac{p}{s^2}\right) (1 - 2pH_s) \equiv 2p \sum_{s=1}^r \left(\frac{1}{s} + \frac{p}{s^2} - 2p \frac{H_s}{s}\right) \pmod{p^3} \end{aligned}$$

and hence

$$\begin{aligned} (-1)^{r+1} \binom{2p-1}{p-r-1} + 1 &\equiv 2pH_r + 2p^2 \sum_{s=1}^r \frac{1}{s^2} - 4p^2 \sum_{1 \leq t \leq s \leq r} \frac{1}{st} \\ &\equiv 2pH_r - 2p^2 H_r^2 \pmod{p^3}. \end{aligned}$$

Let $n \in \mathbb{N} = \{0, 1, 2, \dots\}$ and $r \in \mathbb{Z}$. In 1913 A. Fleck (cf. [D, p. 274]) showed that

$$\text{ord}_p(C_p(n, r)) \geq \left\lfloor \frac{n-1}{p-1} \right\rfloor,$$

where $[\cdot]$ is the well-known floor function, $\text{ord}_p(\alpha)$ denotes the p -adic order of a p -adic number α (we regard $\text{ord}_p(0)$ as $+\infty$), and

$$C_p(n, r) = \sum_{k \equiv r \pmod{p}} \binom{n}{k} (-1)^k. \quad (1.2)$$

(In [S02] the author expressed certain sums like (1.2) in terms of linear recurrences.) Fleck's result plays fundamental roles in the recent investigation of the ψ -operator related to Fontaine's theory, Iwasawa's theory and p -adic Langlands correspondence (cf. [Co], [W] and [SW1]), and Davis and Sun's study of homotopy exponents of special unitary groups (cf. [DS] and [SD]). It is also related to Leopoldt's formula for p -adic L -functions (cf. [Mu, Theorem 8.5]).

Note that if $p \neq 2$ then

$$C_p(2p-1, -1) = \binom{2p-1}{p-1} (-1)^{p-1} + \binom{2p-1}{2p-1} (-1)^{2p-1} = \binom{2p-1}{p-1} - 1$$

and

$$C_p(2p, 0) = \binom{2p}{0} (-1)^0 + \binom{2p}{p} (-1)^p + \binom{2p}{2p} (-1)^{2p} = 2 - \binom{2p}{p}.$$

So, in the case $n = 2p - 1$ and $r = -1$, or the case $n = 2p$ and $r = 0$, Fleck's result yields Babbage's congruence.

For $m = 0, 1, 2, \dots$, the m th order Bernoulli polynomials $B_k^{(m)}(t)$ ($k \in \mathbb{N}$) are given by

$$\frac{x^m e^{tx}}{(e^x - 1)^m} = \sum_{k=0}^{\infty} B_k^{(m)}(t) \frac{x^k}{k!} \quad (0 < |x| < 2\pi),$$

and those $B_k^{(m)} = B_k^{(m)}(0)$ ($k \in \mathbb{N}$) are called m th order Bernoulli numbers. Clearly $B_k^{(0)}(t) = t^k$. The usual Bernoulli polynomials and numbers are $B_k(t) = B_k^{(1)}(t)$ and $B_k = B_k(0) = B_k^{(1)}$ respectively. It can be easily seen that

$$B_k^{(m)}(t) = \sum_{j=0}^k \binom{k}{j} B_j^{(m)} t^{k-j} \quad \text{and} \quad B_k^{(m)}(m-t) = (-1)^k B_k^{(m)}(t).$$

Since $B_k^{(m)}/k!$ coincides with $[x^k](x/(e^x - 1))^m$, the coefficient of x^k in the power series expansion of $(x/(e^x - 1))^m$, if $m \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$ then

$$\frac{B_k^{(m)}}{k!} = \sum_{i_1 + \dots + i_m = k} \frac{B_{i_1} \cdots B_{i_m}}{i_1! \cdots i_m!}.$$

It is well known that $B_0 = 1$, $B_1 = -1/2$ and $B_{2k+1} = 0$ for $k = 1, 2, 3, \dots$. The von Staudt-Clasusen theorem (cf. [IR, pp.233–236] or [Mu, Theorem 2.7]) states that $B_{2k} + \sum_{p-1|2k} 1/p \in \mathbb{Z}$ for any $k \in \mathbb{Z}^+$. Thus, B_0, \dots, B_{p-2} are p -adic integers and hence so are $B_0^{(m)}, \dots, B_{p-2}^{(m)}$. Therefore $B_k^{(m)}(t) \in \mathbb{Z}_p[t]$ if $0 \leq k < p-1$, where \mathbb{Z}_p denotes the ring of p -adic integers.

In terms of higher-order Bernoulli polynomials, the author and D. Wan [SW2] determined the Fleck quotient

$$F_p(n, r) := (-p)^{-\lfloor (n-1)/(p-1) \rfloor} \sum_{k \equiv r \pmod{p}} \binom{n}{k} (-1)^k + \llbracket n = 0 \rrbracket \quad (1.3)$$

modulo p . (Throughout this paper, for an assertion A we let $\llbracket A \rrbracket$ take 1 or 0 according as A holds or not.) Namely, by [SW2, Theorem 1.2], we have

$$F_p(n, r) \equiv -n_*! B_{n_*}^{(m)}(-r) \pmod{p} \quad \text{for } m \in \mathbb{N} \text{ with } m \equiv -n \pmod{p}, \quad (1.4)$$

where n_* is the smallest positive residue of n modulo $p-1$, and n^* is the least nonnegative residue $\{-n\}_{p-1}$ of $-n$ modulo $p-1$. For convenience the notations n_* and n^* will be often used, and we remind the reader of the difference. Note that $n_* + n^* = p-1$ and hence

$$n_*! n^*! = \frac{n_*!(p-1)!}{\prod_{0 < k \leq n_*} (p-k)} \equiv (-1)^{n_*-1} \equiv (-1)^{n^*-1} \equiv (-1)^{n-1} \pmod{p}.$$

We mention that $((p-1)/2)! \pmod{p}$ is related to the class number of the quadratic field $\mathbb{Q}(\sqrt{(-1)^{(p-1)/2}p})$ (cf. [Ch] and [M]).

Let ζ_p be a fixed primitive p th root of unity in the algebraic closure of the p -adic field \mathbb{Q}_p . It is easy to see that $\text{ord}_p(1 - \zeta_p) = 1/(p-1)$. The main trick in [SW2] is to determine $F_p(n, r)$ modulo $1 - \zeta_p$.

Corollary 1.5(i) of Sun and Wan [SW2] states that if $2 \leq n \leq p$ then

$$\frac{1}{p^n} \sum_{k=1}^n \binom{pn-1}{pk-1} (-1)^{pk} \equiv -(n-1)! B_{p-n} \pmod{p}. \quad (1.5)$$

This is a further extension of Wolstenholme's congruence. When $1 < n < p-1$ and $2 \mid n$, the right-hand side of the above congruence is zero since $B_{p-n} = 0$; inspired by (1.5) the author's student H. Pan used *Mathematica* to find the conjecture

$$\frac{1}{p^n} \sum_{k=1}^n \binom{pn-1}{pk-1} (-1)^k \equiv \frac{n!n}{2(n+1)} p B_{p-1-n} \pmod{p^2}. \quad (1.6)$$

For $m, n \in \mathbb{Z}^+$, the Stirling number $S(m, n)$ of the second kind is the number of ways to partition a set of cardinality m into n subsets. It is easy to show that

$$(-1)^{n-1} n! S(m\varphi(p^b), n) \equiv C_p(n, 0) = (-p)^{\lfloor (n-1)/(p-1) \rfloor} F_p(n, 0) \pmod{p^b}$$

for any $b = 1, 2, 3, \dots$ (cf. [GL, Lemma 5]), where φ is Euler's totient function. Thus, for sufficiently large $b > 0$, we have

$$\text{ord}_p(n! S(m\varphi(p^b), n)) = \left\lfloor \frac{n-1}{p-1} \right\rfloor + \text{ord}_p(F_p(n, 0)).$$

In this paper we want to reveal further connections between Fleck quotients and Bernoulli numbers, including the determination of $F_p(pn, r)$ modulo $p^{\text{ord}_p(n)+2}$ by which (1.6) holds when $1 < n < p-1$ and $2 \mid n$. The method of [SW2] does not work for this purpose; instead of $1 - \zeta_p$ we define

$$\pi := - \sum_{k=1}^{p-1} \frac{(1 - \zeta_p)^k}{k} \in \mathbb{Z}_p[\zeta_p]. \quad (1.7)$$

It can be shown that $\pi^{p-1}/p \equiv -1 \pmod{p}$ (see Section 2). The p -adic method in this paper deals with congruences modulo powers of π , and it is so powerful that we need not appeal to the Stickelberger congruence (cf. Theorems 11.2.1 and 11.2.10 of [BEW]) which is of advanced nature.

Sun and Wan [SW2, Corollary 1.7] proved the following periodical results:

$$F_p(n + p^b(p-1), r) \equiv F_p(n, r) \pmod{p^b} \quad \text{for } b = 1, 2, 3, \dots$$

Thus, if $m \in \mathbb{N}$, $m \neq n$ and $m \equiv n \pmod{p(p-1)}$, then $(F_p(m, r) - F_p(n, r))/(m - n) \in \mathbb{Z}_p$. We determine this quotient modulo p in our first theorem.

Theorem 1.1. *Let p be a prime, and let $m, n \in \mathbb{N}$, $m \neq n$ and $m \equiv n \pmod{p(p-1)}$. Then we have*

$$\begin{aligned} \frac{F_p(m, r) - F_p(n, r)}{m - n} &\equiv \frac{(-1)^{n^*}}{n^*!} \sum_{1 < k \leq n^*} \binom{n^*}{k} \frac{B_k}{k} B_{n^*-k}^{\{\{-n\}_p\}}(-r) \\ &\equiv (-1)^{n^*-1} n^*! \sum_{1 < k \leq n^*} \binom{n^* + k}{n^*} r^{n^*-k} \sum_{1 < j \leq k} \binom{k}{j} \frac{B_j}{j} B_{k-j}^{\{\{-n\}_p\}} \pmod{p}. \end{aligned} \quad (1.8)$$

Corollary 1.1. *Let $p \geq 5$ be a prime, and let $n > 0$ be an integer with $n \not\equiv 0, -1 \pmod{p-1}$. Then, there are at least $p - n^* + 2 \geq 4$ values of $r \in \{0, 1, \dots, p-1\}$ such that $\text{ord}_p(F_p(m, r) - F_p(n, r)) = \text{ord}_p(m - n)$ for all $m \in \mathbb{N}$ with $m \equiv n \pmod{p-1}$.*

Proof. Clearly $n^* = \{-n\}_{p-1} \geq 2$ and $n_* = p - 1 - n^* < p - 2$. Since the polynomial

$$P(x) = \sum_{k=2}^{n^*} \binom{n_* + k}{n_*} x^{n^* - k} \sum_{j=2}^k \binom{k}{j} \frac{B_j}{j} B_{k-j}^{\{\{-n\}_p\}} \in \mathbb{Z}_p[x]$$

has degree at most $n^* - 2$, and

$$\begin{aligned} [x^{n^* - 2}]P(x) &= \binom{n_* + 2}{n_*} \sum_{j=2}^2 \binom{2}{j} \frac{B_j}{j} B_{2-j}^{\{\{-n\}_p\}} \\ &= \binom{n_* + 2}{n_*} \frac{B_2}{2} B_0^{\{\{-n\}_p\}} = \frac{(n_* + 1)(n_* + 2)}{24} \not\equiv 0 \pmod{p}, \end{aligned}$$

there are at most $n^* - 2$ values of $r \in \{0, 1, \dots, p-1\}$ satisfying $P(r) \equiv 0 \pmod{p}$. (Recall that a polynomial of degree $d \geq 0$ over a field cannot have more than d zeroes in the field.) Combining this with Theorem 1.1 we obtain the desired result. \square

Corollary 1.2. *Let p be a prime and let $r \in \mathbb{Z}$. Suppose that $n \in \mathbb{N}$ and $n \equiv 0 \pmod{p-1}$. Then, for any $b = 2, 3, \dots$ we have*

$$F_p(n, r) \equiv F_p(\{n\}_{\varphi(p^b)}, r) \pmod{p^b} \quad (1.9)$$

and

$$F_p(n + p - 2, r) \equiv F_p(\{n\}_{\varphi(p^b)} + p - 2, r) \pmod{p^b}. \tag{1.10}$$

Consequently,

$$\frac{F_p(pn, r) + p\llbracket p \mid r \rrbracket - 1}{pn} \equiv 0 \pmod{p}. \quad (1.11)$$

Proof. (i) Let $b > 1$ be an integer. Write $n = \varphi(p^b)q + \{n\}_{\varphi(p^b)}$ with $q \in \mathbb{N}$. As $n^* = \{-n\}_{p-1} = 0$ and $(n + p - 2)^* = \llbracket p \neq 2 \rrbracket$, if $q > 0$ then by Theorem 1.1 we have

$$\frac{F_p(n, r) - F_p(\{n\}_{\varphi(p^b)}, r)}{p^{b-1}(p-1)q} \equiv 0 \pmod{p}$$

and

$$\frac{F_p(n + p - 2, r) - F_p(\{n\}_{\varphi(p^b)} + p - 2, r)}{p^{b-1}(p-1)q} \equiv 0 \pmod{p}.$$

So (1.9) and (1.10) are valid.

(ii) Clearly $b = \text{ord}_p(pn) + 1 \geq 2$ and $\varphi(p^b) \mid pn$. In view of (1.9),

$$F_p(pn, r) \equiv F_p(0, r) = -pC_p(0, r) + 1 = 1 - p\llbracket p \mid r \rrbracket \pmod{p^b}$$

and hence (1.11) holds. \square

Corollary 1.3. *Let p be a prime and let $r \in \mathbb{Z}$. Assume that $m, n \in \mathbb{N}$, $m \neq n$ and $m \equiv n \pmod{p-1}$. Then*

$$\begin{aligned} & \frac{F_p(pm + p - 1, r) - F_p(pn + p - 1, r)}{p(m - n)} \\ & \equiv \frac{(-1)^{n^*}}{n^*!} \left(\sum_{0 < k < n^*} \frac{B_k(-r)}{k} B_{n^*-k}(-r) - H_{n^*-1} B_{n^*}(-r) \right) \pmod{p}. \end{aligned} \quad (1.12)$$

Proof. In light of Theorem 1.1,

$$\frac{F_p(pm + p - 1, r) - F_p(pn + p - 1, r)}{p(m - n)} \equiv \frac{(-1)^{n^*}}{n^*!} B \pmod{p},$$

where

$$B = \sum_{1 < k \leq n^*} \binom{n^*}{k} \frac{B_k}{k} B_{n^*-k}(-r).$$

By a polynomial form of Miki's identity (cf. [PS, (2.3)]),

$$\begin{aligned} B + H_{n^*-1} B_{n^*}(-r) &= \frac{n^*}{2} \sum_{0 < k < n^*} \frac{B_k(-r) B_{n^*-k}(-r)}{k(n^* - k)} \\ &= \frac{1}{2} \sum_{0 < k < n^*} \left(\frac{1}{k} + \frac{1}{n^* - k} \right) B_k(-r) B_{n^*-k}(-r) \\ &= \sum_{0 < k < n^*} \frac{B_k(-r)}{k} B_{n^*-k}(-r). \end{aligned}$$

So we have the desired (1.12). \square

Lemma 1.1. *Let p be an odd prime, and let n be a positive integer not divisible by $p-1$. Then $F_p(n, 0)/n \in \mathbb{Z}_p$.*

With help of this lemma and Theorem 1.1, we can deduce the following theorem.

Theorem 1.2. *Let p be an odd prime, and let $n \in \mathbb{Z}^+$ with $2 \mid n$ and $p-1 \nmid n$. Then*

$$\frac{2 \sum_{k=1}^n \binom{pn-1}{pk-1} (-1)^k}{(-p)^{\lfloor (n-2)/(p-1) \rfloor} p^{n+1} n} = \frac{F_p(pn, 0)}{pn} \equiv \frac{n_*!}{n_* + 1} B_{p-1-n_*} \pmod{p}. \quad (1.13)$$

Given $b, m \in \mathbb{Z}^+$ with $b > 2 \lfloor (pn-1)/(p-1) \rfloor$, we have

$$\begin{aligned} \frac{2}{pn} \cdot \frac{(pn-1)! S(m\varphi(p^b), pn-1)}{(-p)^{\lfloor (pn-2)/(p-1) \rfloor}} &\equiv - \frac{(pn-1)! S(m\varphi(p^b), pn)}{(-p)^{\lfloor (pn-1)/(p-1) \rfloor}} \\ &\equiv \frac{n_*!}{n_* + 1} B_{p-1-n_*} \pmod{p}. \end{aligned} \quad (1.14)$$

Proof. As $n - 1$ is odd, $pn - 1 \equiv n - 1 \not\equiv 0 \pmod{p - 1}$. Thus

$$\begin{aligned}
& p^n (-p)^{\lfloor (n-2)/(p-1) \rfloor} F_p(pn, 0) \\
&= (-p)^{\lfloor (pn-1)/(p-1) \rfloor} F_p(pn, 0) = C_p(pn, 0) \\
&= \sum_{k=1}^n \binom{pn-1}{pk-1} (-1)^{pk} + \sum_{k=0}^{n-1} \binom{pn-1}{pk} (-1)^{pk} \\
&= \sum_{k=1}^n \binom{pn-1}{pk-1} (-1)^k + \sum_{k=0}^{n-1} \binom{pn-1}{p(n-k)-1} (-1)^{n-k} \\
&= 2 \sum_{k=1}^n \binom{pn-1}{pk-1} (-1)^k = 2 \sum_{k=1}^n \binom{pn-1}{p(n-k)} (-1)^{n-k}.
\end{aligned}$$

Since

$$\begin{aligned}
& p^{\lfloor (pn-1)/(p-1) \rfloor + 1} = (1 + (p-1))^{\lfloor (pn-1)/(p-1) \rfloor + 1} \\
& \geq 1 + (p-1) \left(\left\lfloor \frac{pn-1}{p-1} \right\rfloor + 1 \right) > 1 + (p-1) \frac{pn-1}{p-1} = pn,
\end{aligned}$$

we have

$$\left\lfloor \frac{pn-2}{p-1} \right\rfloor = \left\lfloor \frac{pn-1}{p-1} \right\rfloor = n + \left\lfloor \frac{n-1}{p-1} \right\rfloor \geq \text{ord}_p(pn)$$

and hence

$$b - \left\lfloor \frac{pn-1}{p-1} \right\rfloor > \left\lfloor \frac{pn-1}{p-1} \right\rfloor \geq \text{ord}_p(pn).$$

Recall that

$$(-1)^{pn-1} \frac{(pn)! S(m\varphi(p^b), pn)}{(-p)^{\lfloor (pn-1)/(p-1) \rfloor}} \equiv F_p(pn, 0) \pmod{p^{b - \lfloor (pn-1)/(p-1) \rfloor}}$$

and

$$\begin{aligned}
& (-1)^{pn-2} \frac{(pn-1)! S(m\varphi(p^b), pn-1)}{(-p)^{\lfloor (pn-2)/(p-1) \rfloor}} \\
& \equiv F_p(pn-1, 0) = \frac{F_p(pn, 0)}{2} \pmod{p^{b - \lfloor (pn-1)/(p-1) \rfloor}}.
\end{aligned}$$

By the above, it suffices to show that

$$\frac{F_p(pn, 0)}{pn} \equiv \frac{n_*!}{n_* + 1} B_{p-1-n_*} \pmod{p}.$$

Clearly $n^* \neq 0, 1$. By Theorem 1.1 in the case $r = 0$,

$$\begin{aligned} \frac{F_p(p^2n, 0) - F_p(pn, 0)}{pn(p-1)} &\equiv \frac{(-1)^{n^*}}{n^*!} \sum_{1 < k \leq n^*} \binom{n^*}{k} \frac{B_k}{k} B_{n^*-k}^{(0)} \\ &\equiv -n_*! \frac{B_{n^*}}{n^*} \equiv \frac{n_*!}{n_*+1} B_{p-1-n_*} \pmod{p}. \end{aligned}$$

As $F_p(p^2n, 0)/(pn) \equiv 0 \pmod{p}$ by Lemma 1.1, the desired result follows. \square

Remark 1.1. Let p be a prime and let $n \in \mathbb{Z}^+$.

(i) If $p \neq 2$ and $2 \nmid n$, then $F_p(pn, 0) = 0$ because

$$C_p(pn, 0) = \sum_{k=0}^{(n-1)/2} \left(\binom{pn}{pk} (-1)^{pk} + \binom{pn}{p(n-k)} (-1)^{p(n-k)} \right) = 0.$$

If $m \in \mathbb{Z}^+$, $m \equiv n \pmod{p}$ and $m \equiv n \not\equiv 0 \pmod{p-1}$, then by Theorem 1.2 and (1.4) we have the following Kummer-type congruence:

$$\frac{F_p(m, 0)}{m} \equiv \frac{F_p(n, 0)}{n} \pmod{p}.$$

If $1 < n < p-1$ and $2 \mid n$, then (1.13) yields (1.6).

(ii) When $2 \mid n$ and $p-1 \nmid n$, for any integers $m > 0$ and $b > 2 \lfloor (pn-1)/(p-1) \rfloor$ we have

$$\text{ord}_p((pn-1)!S(m\varphi(p^b), pn-1)) \geq \left\lfloor \frac{pn-2}{p-1} \right\rfloor + \text{ord}_p(pn)$$

and

$$\text{ord}_p((pn)!S(m\varphi(p^b), pn)) \geq \left\lfloor \frac{pn-1}{p-1} \right\rfloor + \text{ord}_p(pn)$$

by (1.14). In 2001 I. M. Gessel and T. Lengyel [GL, Conjecture 1] conjectured that equality always holds in our last two inequalities; this is not true since it might happen that $B_{n^*} = B_{p-1-n_*} \equiv 0 \pmod{p}$, e.g., $\text{ord}_{37}(B_{32}) = \text{ord}_{59}(B_{44}) = 1$. (p is said to be *irregular* if $B_{2k} \equiv 0 \pmod{p}$ for some $0 < k < (p-1)/2$. According to [IR, p. 241] or [Mu, Theorem 2.13], there are infinitely many irregular primes.) By the way, Conjecture 2 of [GL] is an easy consequence of the congruence (1.4) due to Sun and Wan [SW2].

Corollary 1.4. *Let $p \geq 5$ be a prime. Then*

$$\binom{2p-1}{p-1} - 1 \equiv -\frac{2}{3}p^3 B_{p-3} \pmod{p^4} \quad (1.15)$$

and

$$\binom{4p}{p} - \binom{4p-1}{2p-1} - 1 \equiv -\frac{48}{5}p^5 B_{p-5} \pmod{p^6}. \quad (1.16)$$

Proof. (1.15) follows from (1.13) in the case $n = 2$. Applying (1.13) with $n = 4$ we find that

$$\frac{2}{4p^5} \left(-\binom{4p-1}{p-1} + \binom{4p-1}{2p-1} - \binom{4p-1}{3p-1} + \binom{4p-1}{4p-1} \right) \equiv \frac{4!}{5} B_{p-5} \pmod{p}.$$

This is equivalent to (1.16) since

$$\binom{4p-1}{p-1} + \binom{4p-1}{3p-1} = \binom{4p-1}{p-1} + \binom{4p-1}{p} = \binom{4p}{p}.$$

We are done. \square

Remark 1.2. (1.15) was first discovered by J.W.L. Glaisher (cf. [G1, p. 21] and [G2, p. 323]). For a prime $p \geq 5$ and $r \in \{1, \dots, p-1\}$, we can determine $\binom{2p-1}{p-1-r} \pmod{p^4}$ in view of (1.15), and (1.1) and its proof.

In the next theorem we determine $F_p(pn, r) \pmod{p^{\text{ord}_p(n)+2}}$ in the case $p-1 \nmid n$ and $p \nmid r$.

Theorem 1.3. *Let p be an odd prime, and let $n > 0$ and r be integers with $p-1 \nmid n$ and $p \nmid r$. Then, for any $b = 1, \dots, \text{ord}_p(pn)$ we have*

$$\begin{aligned} & \frac{(-r)^n F_p(pn, r) + (-1)^{(b-1)n} \prod_{1 \leq k \leq b'n_*, p \nmid k} k}{n_*!} \\ & \equiv n_*(pB_{\varphi(p^b)} - p + 1) - p^b n_* H_{n_*} + n(r^{p-1} - 1) \\ & - pn \sum_{1 < k < p-n_*} \binom{n_* + k}{n_*} \frac{B_k}{kr^k} \pmod{p^{b+1}}, \end{aligned} \quad (1.17)$$

where $b' = (p^b - 1)/(p - 1)$.

Remark 1.3. Let p be an odd prime. A result of L. Carlitz [C] states that $(B_k + p^{-1} - 1)/k \in \mathbb{Z}_p$ for all $k \in \mathbb{Z}^+$ with $p-1 \mid k$. So we have $pB_{\varphi(p^b)} \equiv p-1 \pmod{p^b}$ for all $b \in \mathbb{Z}^+$.

Corollary 1.5. *Let p be an odd prime. Let $n \in \mathbb{Z}^+$ and $r \in \mathbb{Z}$ with $p-1 \nmid n$ and $p \nmid r$. Set $b = \text{ord}_p(pn)$ and $b' = (p^b - 1)/(p - 1)$. Then*

$$F_p(pn, r) + \frac{(-1)^{bn}}{r^n} \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k \in p^b \mathbb{Z}_p = pn \mathbb{Z}_p, \quad (1.18)$$

Proof. This is because the right-hand side of the congruence (1.17) belongs to $p^b \mathbb{Z}_p$. \square

Corollary 1.6. *Let p be an odd prime. If $n \in \mathbb{Z}^+$, $r \in \mathbb{Z}$, $p-1 \nmid n$ and $p \nmid r$, then we can determine $F_p(pn, r) \pmod{p^2}$ in the following way:*

$$\begin{aligned} & \frac{(-r)^n F_p(pn, r) + n_*!}{n_*! n_*} + pH_{n_*} - pB_{p-1} + p - 1 \\ & \equiv \frac{pn}{n_*} \left(q_p(r) - \sum_{1 < k < p-n_*} \binom{n_* + k}{n_*} \frac{B_k}{kr^k} \right) \pmod{p^2}, \end{aligned} \quad (1.19)$$

where $q_p(r)$ denotes the Fermat quotient $(r^{p-1} - 1)/p$.

Proof. Just apply (1.17) with $b = 1$. \square

Remark 1.4. If p is a prime, $n \in \mathbb{N}$ and $r \in \mathbb{Z}$, then

$$F_p(pn + s, r) = \sum_{t=0}^s \binom{s}{t} (-1)^t F_p(pn, r - t) \quad \text{for any } s = 0, \dots, n^*;$$

because $\lfloor (pn + s - 1)/(p - 1) \rfloor = (pn + n^*)/(p - 1) - 1 = \lfloor (pn - 1)/(p - 1) \rfloor$ and $C_p(pn + s, r)$ coincides with

$$\sum_{k \equiv r \pmod{p}} \sum_{t=0}^s \binom{s}{t} \binom{pn}{k-t} (-1)^k = \sum_{t=0}^s \binom{s}{t} (-1)^t C_p(pn, r - t)$$

by the Chu-Vandermonde convolution identity (cf. [GKP, (5.27)]).

In the next section we determine $(\zeta_p^a - 1)^{p^b n}$ modulo $p^{b+1} \pi^{p^b n}$ (where $a \in \mathbb{Z}$ and $b \in \mathbb{N}$) in terms of Bernoulli numbers or higher-order Bernoulli numbers. On the basis of this, we prove Theorem 1.1 and Lemma 1.1 in Section 3 by a p -adic method. In the proof of Theorem 1.3 given in Section 4, we have to employ the p -adic Γ -function and the Gross-Koblitz formula for Gauss sums. In Section 5, we study extended Fleck quotients and give an extension of (1.4) which implies the following generalization of (1.5).

Theorem 1.4. *Let p be a prime, and let $a \in \mathbb{Z}^+$ and $l, m, n \in \mathbb{N}$ with $m < p$ and $2 \leq n - l - m \leq p$. Then we have*

$$\begin{aligned} & \frac{1}{p^{n-l}} \sum_{l < k \leq n} \binom{p^a n - p^{a-1} m - d}{p^a k - p^{a-1} m - d} (-1)^{pk} \binom{k-1}{l} \\ & \equiv \frac{(-1)^{l-1} n! / l!}{\prod_{k=0}^m (n-l-k)} B_{p-n+l+m}^{(m+1)} \pmod{p} \end{aligned} \quad (1.20)$$

for all $d = 1, \dots, \max\{p^{a-2}, 1\}$.

2. A THEOREM ON ROOTS OF UNITY

In this section we establish the following auxiliary result.

Theorem 2.1. *Let p be a prime and define π as in (1.7).*

(i) *We have*

$$\pi^{p-1} \equiv -p \pmod{p^2}, \quad \text{i.e.,} \quad \frac{\pi^{p-1}}{p} \equiv -1 \pmod{p}. \quad (2.1)$$

(ii) *Let $a \in \mathbb{Z}$ and $n \in \mathbb{N}$. If $m \in \mathbb{N}$ and $m \equiv -n \pmod{p}$, then*

$$(\zeta_p^a - 1)^n \equiv \sum_{j=0}^{p-2} B_j^{(m)} \frac{(a\pi)^{n+j}}{j!} \pmod{p\pi^n}. \quad (2.2)$$

For each $b \in \mathbb{Z}^+$ we have

$$(\zeta_p^a - 1)^{p^b n} \equiv (a\pi)^{p^b n} + p^b n \sum_{1 < k < p-1} \frac{B_k}{k!k} (a\pi)^{p^b n+k} \pmod{p^{b+1} \pi^{p^b n}}. \quad (2.3)$$

Lemma 2.1. *Let p be any prime. Then $\text{ord}_p(\pi) = 1/(p-1)$ and $\pi^{p-1}/p \equiv -1 \pmod{\pi}$. Also,*

$$\zeta_p^a \equiv \sum_{k=0}^{p-1} \frac{(a\pi)^k}{k!} \pmod{p\pi} \quad \text{for all } a \in \mathbb{Z}. \quad (2.4)$$

Proof. Clearly

$$\frac{\pi}{\zeta_p - 1} = \sum_{k=1}^{p-1} \frac{(1 - \zeta_p)^{k-1}}{k} = 1 - (1 - \zeta_p)\eta$$

for some $\eta \in \mathbb{Z}_p[\zeta_p]$, hence

$$\frac{\pi}{\zeta_p - 1} \sum_{j=0}^{p-2} (1 - \zeta_p)^j \eta^j = 1 - (1 - \zeta_p)^{p-1} \eta^{p-1}.$$

Since $p/(1 - \zeta_p)^{p-1} = \prod_{a=1}^{p-1} ((1 - \zeta_p^a)/(1 - \zeta_p))$ is a unit in the ring $\mathbb{Z}_p[\zeta_p]$, by the above $\pi/(1 - \zeta_p)$ and π^{p-1}/p are also units in $\mathbb{Z}_p[\zeta_p]$ and hence $\text{ord}_p(\pi) = \text{ord}_p(1 - \zeta_p) = 1/(p-1)$. As $\pi/(1 - \zeta_p) \equiv -1 \pmod{1 - \zeta_p}$ and

$$\frac{p}{(1 - \zeta_p)^{p-1}} = \prod_{a=1}^{p-1} (1 + \zeta_p + \cdots + \zeta_p^{a-1}) \equiv \prod_{a=1}^{p-1} a \equiv -1 \pmod{\zeta_p - 1},$$

we have

$$\frac{\pi^{p-1}}{p} = \frac{(1 - \zeta_p)^{p-1}}{p} \left(\frac{\pi}{1 - \zeta_p} \right)^{p-1} \equiv -(-1)^{p-1} \equiv -1 \pmod{\pi}.$$

Write

$$\sum_{k=0}^{p-1} \frac{(-\sum_{j=1}^{p-1} x^j/j)^k}{k!} = P(x) + x^p Q(x)$$

with $P(x), Q(x) \in \mathbb{Z}_p[x]$ and $\deg P(x) < p$. If $-1 < x < 1$ then

$$1 - x = e^{\log(1-x)} = \sum_{k=0}^{\infty} \frac{(\log(1-x))^k}{k!} = \sum_{k=0}^{\infty} \frac{(-\sum_{j=1}^{\infty} x^j/j)^k}{k!}.$$

Comparing the coefficients of $1, x, \dots, x^{p-1}$ we find that $P(x) = 1 - x$. Therefore

$$\sum_{k=0}^{p-1} \frac{\pi^k}{k!} = P(1 - \zeta_p) + (1 - \zeta_p)^p Q(1 - \zeta_p) \equiv \zeta_p \pmod{\pi^p}.$$

If $j \in \mathbb{N}$ and $\zeta_p^j \equiv \sum_{k=0}^{p-1} (j\pi)^k/k! \pmod{\pi^p}$, then

$$\begin{aligned} \zeta_p^{j+1} &\equiv \sum_{k=0}^{p-1} \frac{j^k \pi^k}{k!} \sum_{l=0}^{p-1} \frac{\pi^l}{l!} \\ &\equiv \sum_{n=0}^{p-1} \left(\sum_{k=0}^n \binom{n}{k} j^k \right) \frac{\pi^n}{n!} = \sum_{n=0}^{p-1} \frac{(j+1)^n \pi^n}{n!} \pmod{\pi^p}. \end{aligned}$$

Thus, by induction, $\zeta_p^a \equiv \sum_{k=0}^{p-1} (a\pi)^k/k! \pmod{\pi^p}$ for any $a \in \mathbb{N}$.

A general integer a can be written in the form $pq + r$ with $q, r \in \mathbb{Z}$ and $0 \leq r < p$. In view of the above,

$$\begin{aligned} \zeta_p^a &= \zeta_p^r \equiv \sum_{k=0}^{p-1} \frac{(r\pi)^k}{k!} = 1 + \sum_{k=1}^{p-1} \frac{(a - pq)^k \pi^k}{k!} \\ &\equiv 1 + \sum_{k=1}^{p-1} \frac{a^k \pi^k}{k!} = \sum_{k=0}^{p-1} \frac{(a\pi)^k}{k!} \pmod{p\pi}. \end{aligned}$$

This concludes the proof. \square

Lemma 2.2. *Let $k \in \mathbb{Z}^+$ and $m \in \mathbb{N}$. Then*

$$\sum_{\substack{i_1, \dots, i_k \in \mathbb{N} \\ \sum_{j=1}^k i_j = k}} (-1)^{i_1 + \dots + i_k} \frac{(\sum_{j=1}^k i_j - 1)!}{i_1! \cdots i_k!} \prod_{j=1}^k \left(\frac{B_j^{(m)}}{j!} \right)^{i_j} = m \frac{(-1)^k B_k}{k!k}. \quad (2.5)$$

Proof. For $0 < x < 2\pi$ we have

$$\begin{aligned} & \frac{d}{dx} \left(\log \frac{e^x - 1}{x} - \sum_{n=1}^{\infty} \frac{B_n}{n} \cdot \frac{(-x)^n}{n!} \right) \\ &= \frac{e^x}{e^x - 1} - \frac{1}{x} - \sum_{n=1}^{\infty} B_n \frac{(-1)^n x^{n-1}}{n!} \\ &= \frac{1}{1 - e^{-x}} + \sum_{n=0}^{\infty} B_n \frac{(-x)^{n-1}}{n!} \\ &= \frac{1}{1 - e^{-x}} + \frac{1}{-x} \cdot \frac{-x}{e^{-x} - 1} = 0. \end{aligned}$$

So $f(x) = \log((e^x - 1)/x) - \sum_{n=1}^{\infty} (-x)^n B_n / (n!n)$ is a constant for $x \in (0, 2\pi)$. Letting $x \rightarrow 0$ we find that the constant is zero.

In light of the above,

$$\begin{aligned} m \frac{(-1)^{k-1} B_k}{k!k} &= -m [x^k] \log \left(\frac{e^x - 1}{x} \right) = [x^k] \log \left(\frac{x}{e^x - 1} \right)^m \\ &= [x^k] \log \left(1 + \sum_{j=1}^{\infty} B_j^{(m)} \frac{x^j}{j!} \right) \\ &= [x^k] \sum_{n=1}^k \frac{(-1)^{n-1}}{n} \left(\sum_{j=1}^k \frac{B_j^{(m)}}{j!} x^j \right)^n \\ &= [x^k] \sum_{n=1}^k \frac{(-1)^{n-1}}{n} \sum_{\substack{i_1, \dots, i_k \in \mathbb{N} \\ i_1 + \dots + i_k = n}} \frac{n!}{i_1! \cdots i_k!} \prod_{j=1}^k \left(\frac{B_j^{(m)}}{j!} x^j \right)^{i_j} \\ &= \sum_{\substack{i_1, \dots, i_k \in \mathbb{N} \\ \sum_{j=1}^k i_j = k}} (-1)^{i_1 + \dots + i_k - 1} \frac{(\sum_{j=1}^k i_j - 1)!}{i_1! \cdots i_k!} \prod_{j=1}^k \left(\frac{B_j^{(m)}}{j!} \right)^{i_j}. \end{aligned}$$

So we have the desired (2.5). \square

Proof of Theorem 2.1. (i) When $p = 2$, (2.1) is trivial since $\pi = \zeta_2 - 1 = -2$.

Now we consider the case $p > 2$. In view of (2.4),

$$\begin{aligned} \sum_{a=1}^{p-1} \frac{\zeta_p^a}{a^p} &\equiv \sum_{k=0}^{p-1} \frac{\pi^k}{k!} \sum_{a=1}^{p-1} \frac{1}{a^{p-k}} \\ &\equiv \sum_{a=1}^{(p-1)/2} \left(\frac{1}{a^p} + \frac{1}{(p-a)^p} \right) + \pi \sum_{a=1}^{p-1} \frac{1}{a^{p-1}} + \sum_{1 < k < p} \frac{\pi^k}{k!} \sum_{a=1}^{p-1} a^{k-1} \\ &\equiv \sum_{a=1}^{(p-1)/2} \left(\frac{1}{a^p} + \frac{1}{(-a)^p} \right) + \pi(p-1) \equiv -\pi \pmod{p\pi}. \end{aligned}$$

(It is well known that $\sum_{a=1}^{p-1} a^j \equiv 0 \pmod{p}$ for any $j \in \mathbb{N}$ with $p-1 \nmid j$, see, e.g., [IR, pp. 235–236].) For the norm

$$\alpha := N_{\mathbb{Q}_p(\zeta_p)/\mathbb{Q}} \left(\sum_{a=1}^{p-1} \frac{\zeta_p^a}{a^p} \right) \in \mathbb{Z}_p,$$

we have

$$\begin{aligned} \alpha &= \prod_{k=1}^{p-1} \sum_{a=1}^{p-1} \frac{(\zeta_p^k)^a}{a^p} = \prod_{k=1}^{p-1} \left(k^p \sum_{a=1}^{p-1} \frac{\zeta_p^{ka}}{(ka)^p} \right) \\ &\equiv \prod_{k=1}^{p-1} \left(k^p \sum_{a=1}^{p-1} \frac{\zeta_p^{ka}}{(\{ka\}_p)^p} \right) = ((p-1)!)^p \prod_{k=1}^{p-1} \sum_{a=1}^{p-1} \frac{\zeta_p^a}{a^p} \\ &\equiv (-1)^p \left(\sum_{a=1}^{p-1} \frac{\zeta_p^a}{a^p} \right)^{p-1} \equiv -\pi^{p-1} \left(\frac{1}{-\pi} \sum_{a=1}^{p-1} \frac{\zeta_p^a}{a^p} \right)^{p-1} \equiv -\pi^{p-1} \pmod{p^2}. \end{aligned}$$

(Note that $(\beta + p\gamma)^p \equiv \beta^p \pmod{p^2}$ for any $\beta, \gamma \in \mathbb{Z}_p[\zeta_p]$.) Thus $\alpha \equiv -\pi^{p-1} \equiv p \pmod{p\pi}$ and hence $\text{ord}_p(\alpha - p) \geq \text{ord}_p(p\pi) = 1 + 1/(p-1) > 1$. As $\alpha - p \in \mathbb{Z}_p$, we must have $\text{ord}_p(\alpha - p) \geq 2$ and so $-\pi^{p-1} \equiv \alpha \equiv p \pmod{p^2}$. This proves (2.1).

(ii) Let $b, m \in \mathbb{N}$ and $m \equiv -n \pmod{p}$. Observe that if $0 < |x| < 2\pi$ then

$$\begin{aligned} &\left(\sum_{k=0}^{p-2} B_k^{(m)} \frac{x^k}{k!} + \sum_{k=p-1}^{\infty} B_k^{(m)} \frac{x^k}{k!} \right) \left(\sum_{k=1}^{p-1} \frac{x^{k-1}}{k!} + \sum_{k=p}^{\infty} \frac{x^{k-1}}{k!} \right)^m \\ &= \left(\frac{x}{e^x - 1} \right)^m \left(\frac{e^x - 1}{x} \right)^m = 1. \end{aligned}$$

By comparing the coefficients of $1, x, \dots, x^{p-2}$ we find that

$$\left(\sum_{k=0}^{p-2} B_k^{(m)} \frac{x^k}{k!} \right) \left(\sum_{k=1}^{p-1} \frac{x^{k-1}}{k!} \right)^m = 1 + x^{p-1} Q(x)$$

for some $Q(x) \in \mathbb{Z}_p[x]$. It follows that

$$\left(\sum_{k=0}^{p-2} B_k^{(m)} \frac{(a\pi)^k}{k!} \right) \left(\sum_{k=1}^{p-1} \frac{(a\pi)^{k-1}}{k!} \right)^m \equiv 1 \pmod{p}$$

and

$$\left(\sum_{k=0}^{p-2} B_k^{(m)} \frac{(a\pi)^k}{k!} \right)^{p^b} \equiv \left(\sum_{k=1}^{p-1} \frac{(a\pi)^{k-1}}{k!} \right)^{-p^b m} \pmod{p^{b+1}}.$$

(Note that $(\beta + p^i \gamma)^p \equiv \beta^p \pmod{p^{i+1}}$ for any $i \in \mathbb{N}$ and $\beta, \gamma \in \mathbb{Z}_p[\zeta_p]$.)

Since $\sum_{k=1}^{p-1} (a\pi)^{k-1}/k! = 1 + \pi\beta$ for some $\beta \in \mathbb{Z}_p[\zeta_p]$, we have

$$\left(\sum_{k=1}^{p-1} \frac{(a\pi)^{k-1}}{k!} \right)^{p^{b+1}} = (1 + \pi\beta)^{p^{b+1}} \equiv 1 \pmod{p^{b+1}\pi}.$$

Note that $p^b n \equiv -p^b m \pmod{p^{b+1}}$. Therefore

$$\begin{aligned} \left(\sum_{k=1}^{p-1} \frac{(a\pi)^{k-1}}{k!} \right)^{p^b n} &\equiv \left(\sum_{k=1}^{p-1} \frac{(a\pi)^{k-1}}{k!} \right)^{-p^b m} \\ &\equiv \left(\sum_{j=0}^{p-2} B_j^{(m)} \frac{(a\pi)^j}{j!} \right)^{p^b} \pmod{p^{b+1}}. \end{aligned}$$

In view of Lemma 2.1,

$$\frac{\zeta_p^a - 1}{\pi} \equiv a \sum_{k=1}^{p-1} \frac{(a\pi)^{k-1}}{k!} \pmod{p}.$$

Thus

$$\begin{aligned} \left(\frac{\zeta_p^a - 1}{\pi} \right)^{p^b n} &\equiv \left(a \sum_{k=1}^{p-1} \frac{(a\pi)^{k-1}}{k!} \right)^{p^b n} \\ &\equiv a^{p^b n} \left(\sum_{j=0}^{p-2} B_j^{(m)} \frac{(a\pi)^j}{j!} \right)^{p^b} \pmod{p^{b+1}}. \end{aligned} \tag{2.6}$$

In the case $b = 0$, this yields (2.2).

Below we assume $b > 0$ and want to prove (2.3).

By the multi-nomial theorem and the fact that $\pi^{2p-2} \equiv 0 \pmod{p^2}$,

$$\begin{aligned}
& \left(\sum_{j=0}^{p-2} B_j^{(m)} \frac{(a\pi)^j}{j!} \right)^p \\
&= \sum_{\substack{i_0, \dots, i_{p-2} \in \mathbb{N} \\ i_0 + \dots + i_{p-2} = p}} \frac{p!}{i_0! \cdots i_{p-2}!} \prod_{j=0}^{p-2} \left(B_j^{(m)} \frac{(a\pi)^j}{j!} \right)^{i_j} \\
&\equiv \sum_{k=0}^{2p-3} (a\pi)^k \sum_{\substack{\sum_{j=0}^{p-2} i_j = p \\ \sum_{j=0}^{p-2} i_j j = k}} \frac{p!}{i_0! \cdots i_{p-2}!} \prod_{j=0}^{p-2} \left(\frac{B_j^{(m)}}{j!} \right)^{i_j} \pmod{p^2}.
\end{aligned}$$

If $i_0, \dots, i_{p-2} \in \mathbb{N}$, $\sum_{j=0}^{p-2} i_j = p$ and $\sum_{j=0}^{p-2} i_j j = k \not\equiv 0 \pmod{p}$ then i_0, \dots, i_{p-2} are all smaller than p and hence $p! / \prod_{j=0}^{p-2} i_j! \equiv 0 \pmod{p}$. Thus

$$\begin{aligned}
& \left(\sum_{j=0}^{p-2} B_j^{(m)} \frac{(a\pi)^j}{j!} \right)^p \\
&\equiv (a\pi)^0 \frac{p!}{p!0! \cdots 0!} \left(\frac{B_0^{(m)}}{0!} \right)^p \\
&+ \llbracket p \neq 2 \rrbracket (a\pi)^p \frac{p!}{0!p!0! \cdots 0!} \left(\frac{B_1^{(m)}}{1!} \right)^p \\
&+ \sum_{0 < k < p-1} (a\pi)^k \sum_{\substack{\sum_{j=0}^{p-2} i_j = p \\ \sum_{j=0}^{p-2} i_j j = k}} \frac{p!}{i_0! \cdots i_{p-2}!} \prod_{j=0}^{p-2} \left(\frac{B_j^{(m)}}{j!} \right)^{i_j} \pmod{p^2}.
\end{aligned}$$

Note that $B_0^{(m)} = 1$ and

$$B_1^{(m)} = [x] \left(\frac{x}{e^x - 1} \right)^m = [x] \left(1 - \frac{x}{2} + \sum_{k=2}^{\infty} B_k \frac{x^k}{k!} \right)^m = -\frac{m}{2}.$$

If $p \neq 2$ then $(-m/2)^p \equiv -m/2 \equiv n/2 \pmod{p}$. If $0 < k < p-1$, then

$$\begin{aligned}
& \sum_{\substack{\sum_{j=0}^{p-2} i_j = p \\ \sum_{j=0}^{p-2} i_j j = k}} \frac{p!}{i_0! \cdots i_{p-2}!} \prod_{j=0}^{p-2} \left(\frac{B_j^{(m)}}{j!} \right)^{i_j} \\
&= \sum_{\substack{i_1, \dots, i_k \in \mathbb{N} \\ \sum_{j=1}^k i_j j = k}} \frac{p!(B_0^{(m)}/0!)^{p-\sum_{j=1}^k i_j}}{(p-\sum_{j=1}^k i_j)!} \prod_{j=1}^k \frac{(B_j^{(m)}/j!)^{i_j}}{i_j!} \\
&= \sum_{\substack{i_1, \dots, i_k \in \mathbb{N} \\ \sum_{j=1}^k i_j j = k}} \prod_{0 \leq i < \sum_{j=1}^k i_j} (p-i) \times \prod_{j=1}^k \frac{(B_j^{(p-n)}/j!)^{i_j}}{i_j!} \\
&\equiv p \sum_{\substack{i_1, \dots, i_k \in \mathbb{N} \\ \sum_{j=1}^k i_j j = k}} (-1)^{\sum_{j=1}^k i_j - 1} \frac{(\sum_{j=1}^k i_j - 1)!}{i_1! \cdots i_k!} \prod_{j=1}^k \left(\frac{B_j^{(m)}}{j!} \right)^{i_j} \pmod{p^2}.
\end{aligned}$$

Therefore, with help of Lemma 2.2, we have

$$\left(\sum_{j=0}^{p-2} B_j^{(m)} \frac{(a\pi)^j}{j!} \right)^p \equiv 1 + p\pi S \pmod{p^2}$$

where

$$\begin{aligned}
S &= \llbracket p \neq 2 \rrbracket \frac{n}{2} \cdot \frac{a^p \pi^{p-1}}{p} - m \sum_{0 < k < p-1} a^k \pi^{k-1} \frac{(-1)^k B_k}{k!k} \\
&\equiv \llbracket p \neq 2 \rrbracket \frac{an}{2} \left(\frac{\pi^{p-1}}{p} + 1 \right) + n \sum_{1 < k < p-1} a^k \pi^{k-1} \frac{B_k}{k!k} \\
&\equiv n \sum_{1 < k < p-1} a^k \pi^{k-1} \frac{B_k}{k!k} \pmod{p} \quad (\text{by (2.1)}).
\end{aligned}$$

(Recall that $B_1 = -1/2$ and $B_{2j+1} = 0$ for all $j \in \mathbb{Z}^+$.)

Since $b > 0$, it follows from the above that

$$\begin{aligned}
\left(\sum_{j=0}^{p-2} B_j^{(m)} \frac{(a\pi)^j}{j!} \right)^{p^b} &\equiv (1 + p\pi S)^{p^{b-1}} \\
&\equiv 1 + p^{b-1} p\pi S + \sum_{1 < k \leq p^{b-1}} p^{b-1} \binom{p^{b-1} - 1}{k-1} \frac{p^k \pi^k}{k} S^k \\
&\equiv 1 + p^b \pi S \pmod{p^{b+1}},
\end{aligned}$$

where we have noted that $\pi^2/2 \in \mathbb{Z}_p[\zeta_p]$ and $p^{k-2}/k \in \mathbb{Z}_p$ for $k = 3, 4, \dots$ (cf. [S03, Lemma 2.1]). Combining this with (2.6), we find that

$$\begin{aligned} (\zeta_p^a - 1)^{p^b n} &\equiv (a\pi)^{p^b n} (1 + p^b \pi S) \\ &\equiv (a\pi)^{p^b n} + p^b n \sum_{1 < k < p-1} (a\pi)^{p^b n+k} \frac{B_k}{k!k} \pmod{p^{b+1} \pi^{p^b n}}. \end{aligned}$$

This proves (2.3) and we are done. \square

3. PROOFS OF THEOREM 1.1 AND LEMMA 1.1

Lemma 3.1. *Let p be a prime, and let $n \in \mathbb{N}$ and $r \in \mathbb{Z}$. Then*

$$pC_p(n, r) = \sum_{a=0}^{p-1} \zeta_p^{-ar} (1 - \zeta_p^a)^n. \quad (3.1)$$

Proof. This known result can be easily proved by using $\llbracket p \mid k - r \rrbracket = p^{-1} \sum_{a=0}^{p-1} \zeta_p^{a(k-r)}$. \square

Proof of Lemma 1.1. For $a = 1, \dots, p-1$, as $(\zeta_p^a - 1)/\pi \equiv a \pmod{\pi}$ by Lemma 2.1, we have

$$\left(\frac{\zeta_p^a - 1}{\pi} \right)^n \equiv a^n \pmod{p^{\text{ord}_p(n)} \pi}$$

since

$$\left(\frac{\zeta_p^a - 1}{\pi} \right)^p \equiv a^p \pmod{p\pi}, \quad \left(\frac{\zeta_p^a - 1}{\pi} \right)^{p^2} \equiv a^{p^2} \pmod{p^2\pi}, \quad \dots$$

Let g be a primitive root modulo p . Then $g^n \not\equiv 1 \pmod{p}$ (as $p-1 \nmid n$), and also

$$\begin{aligned} (g^n - 1) \sum_{a=1}^{p-1} a^n &= \sum_{a=1}^{p-1} (ag)^n - \sum_{a=1}^{p-1} a^n \\ &\equiv \sum_{a=1}^{p-1} (\{ag\}_p)^n - \sum_{a=1}^{p-1} a^n \equiv 0 \pmod{p^{\text{ord}_p(n)+1}}. \end{aligned}$$

Therefore $\sum_{a=1}^{p-1} a^n \equiv 0 \pmod{p^{\text{ord}_p(n)+1}}$ and hence

$$\sum_{a=1}^{p-1} \left(\frac{\zeta_p^a - 1}{\pi} \right)^n \equiv \sum_{a=1}^{p-1} a^n \equiv 0 \pmod{p^{\text{ord}_p(n)} \pi}.$$

On the other hand,

$$\sum_{a=0}^{p-1} \left(\frac{\zeta_p^a - 1}{\pi} \right)^n = \frac{pC_p(n, 0)}{(-\pi)^n}$$

by Lemma 3.1. So we have

$$\text{ord}_p(C_p(n, 0)) \geq \text{ord}_p \left(p^{\text{ord}_p(n)-1} \pi^{n+1} \right) = \text{ord}_p(n) - 1 + \frac{n+1}{p-1}$$

and hence

$$\text{ord}_p(F_p(n, 0)) = \text{ord}_p(C_p(n, 0)) - \left\lfloor \frac{n-1}{p-1} \right\rfloor > \text{ord}_p(n) - 1.$$

Since $F_p(n, 0) \in \mathbb{Z}$, this shows that $F_p(n, 0)/n \in \mathbb{Z}_p$. We are done. \square

Lemma 3.2. *Let p be a prime, and let $n \in \mathbb{N}$, $r \in \mathbb{Z}$ and $r \not\equiv 0 \pmod{p}$. Then*

$$\sum_{a=1}^{p-1} a^n (\zeta_p^{ar} - 1) \equiv -\frac{(r\pi)^{n^*}}{n^*!} p^{\llbracket p-1 \mid n \rrbracket} \equiv n_*! (-r\pi)^{n^*} p^{\llbracket p-1 \mid n \rrbracket} \pmod{p\pi}. \quad (3.2)$$

Proof. In view of Lemma 2.1,

$$\sum_{a=1}^{p-1} a^n (\zeta_p^{ar} - 1) \equiv \sum_{a=1}^{p-1} a^n \sum_{k=1}^{p-1} \frac{(ar\pi)^k}{k!} = \sum_{k=1}^{p-1} \frac{(r\pi)^k}{k!} \sum_{a=1}^{p-1} a^{n+k} \pmod{p\pi}.$$

Since $\sum_{a=1}^{p-1} a^{n+k} \equiv -\llbracket p-1 \mid n+k \rrbracket \pmod{p}$, we have

$$\begin{aligned} \sum_{a=1}^{p-1} a^n (\zeta_p^{ar} - 1) &\equiv - \sum_{\substack{1 \leq k \leq p-1 \\ p-1 \mid k-n^*}} \frac{(r\pi)^k}{k!} = -\frac{(r\pi)^{n^*}}{n^*!} \left(\frac{(r\pi)^{p-1}}{(p-1)!} \right)^{\llbracket n^*=0 \rrbracket} \\ &\equiv -\frac{(r\pi)^{n^*}}{n^*!} (-\pi^{p-1})^{\llbracket n^*=0 \rrbracket} \equiv -\frac{(r\pi)^{n^*}}{n^*!} p^{\llbracket p-1 \mid n \rrbracket} \\ &\equiv n_*! (-r\pi)^{n^*} p^{\llbracket p-1 \mid n \rrbracket} \pmod{p\pi}. \end{aligned}$$

This proves (3.2). \square

Proof of Theorem 1.1. Without loss of generality, we assume $m > n$ and write $m - n = p^b(p-1)d$ with $b, d \in \mathbb{Z}^+$ and $p \nmid d$. Clearly $2 \mid p^b(p-1)$. Set

$$D = \frac{1}{\pi^{n^*}} \sum_{a=1}^{p-1} \zeta_p^{-ar} \left(\frac{\zeta_p^a - 1}{\pi} \right)^n \left(\left(\frac{\zeta_p^a - 1}{\pi} \right)^{p^b(p-1)d} - 1 \right).$$

Then

$$\begin{aligned}
(-1)^n D &= \frac{(-1)^{n+p^b(p-1)d}}{\pi^{p^b(p-1)d+n+n^*}} \sum_{a=1}^{p-1} \zeta_p^{-ar} (\zeta_p^a - 1)^{n+p^b(p-1)d} \\
&\quad - \frac{(-1)^n}{\pi^{n+n^*}} \sum_{a=1}^{p-1} \zeta_p^{-ar} (\zeta_p^a - 1)^n \\
&= \frac{pC_p(n+p^b(p-1)d, r)}{\pi^{p^b(p-1)d+n+n^*}} - \frac{pC_p(n, r) - \llbracket n=0 \rrbracket}{\pi^{n+n^*}} \quad (\text{by Lemma 3.1}) \\
&= \frac{(-p)^{(n+n^*)/(p-1)}}{\pi^{n+n^*}} F_p(n, r) - \frac{(-p)^{p^b d+(n+n^*)/(p-1)}}{\pi^{p^b(p-1)d+n+n^*}} F_p(m, r)
\end{aligned}$$

and hence

$$\begin{aligned}
(-1)^n D \left(\frac{\pi^{p-1}}{-p} \right)^{(n+n^*)/(p-1)} &= F_p(n, r) - \left(\frac{-p}{\pi^{p-1}} \right)^{p^b d} F_p(m, r) \\
&\equiv F_p(n, r) - F_p(m, r) \pmod{p^b \pi}.
\end{aligned}$$

(Note that $(-p/\pi^{p-1})^{p^b} \equiv 1 \pmod{p^b \pi}$ since $-p/\pi^{p-1} \equiv 1 \pmod{\pi}$.)

Let a be an integer not divisible by p . In view of Theorem 2.1(ii),

$$\left(\frac{\zeta_p^a - 1}{\pi} \right)^n \equiv \sum_{j=0}^{p-2} a^{n+j} B_j^{\{\{-n\}_p\}} \frac{\pi^j}{j!} \pmod{p},$$

and

$$\begin{aligned}
&\left(\frac{\zeta_p^a - 1}{\pi} \right)^{p^b(p-1)d} - 1 \\
&\equiv a^{p^b(p-1)d} - 1 + p^b(p-1)d \sum_{1 < k < p-1} \frac{B_k}{k!k} a^{p^b(p-1)d+k} \pi^k \\
&\equiv -p^b d \sum_{1 < k < p-1} \frac{B_k}{k!k} a^k \pi^k \pmod{p^{b+1}}
\end{aligned}$$

with help of the congruence $a^{\varphi(p^{b+1})} \equiv 1 \pmod{p^{b+1}}$ (Euler's theorem).

In light of the above,

$$\pi^{n^*} D \equiv -p^b d \sum_{j=0}^{p-2} \sum_{1 < k < p-1} \frac{B_j^{\{\{-n\}_p\}} B_k}{j!k!k} \pi^{j+k} G_r(n+j+k) \pmod{p^{b+1}},$$

where $G_r(s) = \sum_{a=1}^{p-1} a^s \zeta_p^{-ar}$ for $s \in \mathbb{Z}$. By Lemma 3.2, for $k = 0, \dots, p-2$ we have

$$\begin{aligned} G_r(n+k) &= \sum_{a=1}^{p-1} a^{n+k} (\zeta_p^{-ar} - 1) + \sum_{a=1}^{p-1} a^{n+k} \\ &\equiv \llbracket k < n^* \rrbracket (n_* + k)! (r\pi)^{n^* - k} - \llbracket k = n^* \rrbracket \\ &\equiv \llbracket k \leq n^* \rrbracket (n_* + k)! (r\pi)^{n^* - k} \pmod{\pi^{n^* + 1}}. \end{aligned}$$

(Note that $(n_* + n^*)! = (p-1)! \equiv -1 \pmod{p}$ by Wilson's theorem.) So $\pi^{n^*} D$ is congruent to

$$-p^b d \sum_{1 < k \leq n^*} \sum_{j=0}^{n^* - k} \frac{B_j^{\{\{-n\}_p\}} B_k}{j! k! k} \pi^{j+k} (n_* + j + k)! (r\pi)^{n^* - j - k}$$

modulo $p^b \pi^{n^* + 1}$ and hence

$$\begin{aligned} D &\equiv -p^b d \sum_{1 < l \leq n^*} (n_* + l)! r^{n^* - l} \sum_{1 < k \leq l} \frac{B_k}{k! k} \cdot \frac{B_{l-k}^{\{\{-n\}_p\}}}{(l-k)!} \\ &\equiv -p^b d \times n_*! \Sigma \pmod{p^b \pi}, \end{aligned}$$

where

$$\Sigma = \sum_{1 < l \leq n^*} \binom{n_* + l}{n_*} r^{n^* - l} \sum_{1 < k \leq l} \binom{l}{k} \frac{B_k}{k} B_{l-k}^{\{\{-n\}_p\}}.$$

Therefore

$$F_p(m, r) - F_p(n, r) \equiv (-1)^{n-1} D \equiv p^b d (-1)^n n_*! \Sigma \pmod{p^b \pi},$$

i.e., the p -adic order of the rational number

$$R = \frac{F_p(m, r) - F_p(n, r)}{m - n} + (-1)^{n_*} n_*! \Sigma$$

is at least $\text{ord}_p(\pi) = 1/(p-1) > 0$. It follows that $\text{ord}_p(R) \geq 1$.

If $0 < l \leq n^*$, then

$$\begin{aligned} \binom{n_* + l}{n_*} &= \prod_{k=1}^l \frac{p-1-n^*+k}{k} \\ &\equiv (-1)^l \prod_{k=1}^l \frac{n^* - k + 1}{j} = (-1)^l \binom{n^*}{l} \pmod{p}. \end{aligned}$$

Note also that

$$\begin{aligned}
& \sum_{1 < l \leq n^*} (-1)^l \binom{n^*}{l} r^{n^* - l} \sum_{1 < k \leq l} \binom{l}{k} \frac{B_k}{k} B_{l-k}^{\{\{-n\}_p\}} \\
&= (-1)^{n^*} \sum_{1 < k \leq n^*} \binom{n^*}{k} \frac{B_k}{k} \sum_{l=k}^{n^*} \binom{n^* - k}{l - k} B_{l-k}^{\{\{-n\}_p\}} (-r)^{n^* - k - (l - k)} \\
&= (-1)^{n^*} \sum_{1 < k \leq n^*} \binom{n^*}{k} \frac{B_k}{k} B_{n^* - k}^{\{\{-n\}_p\}} (-r).
\end{aligned}$$

So we have

$$\begin{aligned}
\frac{F_p(m, r) - F_p(n, r)}{m - n} &\equiv (-1)^{n^* - 1} n^*! \Sigma \equiv \frac{\Sigma}{n^*!} \\
&\equiv \frac{(-1)^{n^*}}{n^*!} \sum_{1 < k \leq n^*} \binom{n^*}{k} \frac{B_k}{k} B_{n^* - k}^{\{\{-n\}_p\}} (-r) \pmod{p}.
\end{aligned}$$

This concludes the proof. \square

4. PROOF OF THEOREM 1.3

The following lemma in the case $b = 1$ is a known result due to Beeger in 1913 (cf. [Mu, p. 23]).

Lemma 4.1. *Let p be an odd prime, and let b be a positive integer. Then*

$$w_{p^b} \equiv \frac{pB_{\varphi(p^b)} - p + 1}{p^b} \pmod{p}, \quad (4.1)$$

where w_{p^b} denotes the generalized Wilson quotient $(1 + \prod_{0 < a < p^b, p \nmid a} a) / p^b$.

Proof. Let g be a primitive root modulo p^b . As Gauss discovered,

$$\prod_{\substack{a=1 \\ p \nmid a}}^{p^b - 1} a \equiv \prod_{k=0}^{\varphi(p^b) - 1} g^k = g^{\varphi(p^b)(\varphi(p^b) - 1)/2} \equiv g^{\varphi(p^b)/2} \equiv -1 \pmod{p^b}.$$

So w_{p^b} is an integer.

Clearly

$$\frac{3B_{\varphi(3)} - 3 + 1}{3} = \frac{3/6 - 3 + 1}{3} = -\frac{1}{2} \equiv w_3 = \frac{2! + 1}{3} \pmod{3}.$$

Below we assume $p^b > 3$.

Let $k > 1$ be an integer. Recall that

$$\begin{aligned} \sum_{a=1}^{p-1} a^k &= \frac{B_{k+1}(p) - B_{k+1}}{k+1} = \frac{1}{k+1} \sum_{j=0}^k \binom{k+1}{j+1} B_{k-j} p^{j+1} \\ &= pB_k + pk \sum_{j=1}^k \binom{k-1}{j-1} (pB_{k-j}) \frac{p^{j-1}}{j(j+1)}. \end{aligned}$$

Since $p \neq 2$, by [S03, Lemma 2.1] we have $p^{j-2}/(j(j+1)) \in \mathbb{Z}_p$ for $j = 3, 4, \dots$. Note also that $pB_{k-j} \in \mathbb{Z}_p$ ($0 \leq j \leq k$) by the von Staudt-Clausen theorem. So

$$\sum_{a=1}^{p-1} a^k \equiv pB_k + pk \left(\frac{p}{2} B_{k-1} + (k-1)pB_{k-2} \frac{p}{2 \times 3} \right) \pmod{p^{\text{ord}_p(k)+2}}.$$

When $p-1 \mid k$, we have $B_{k-1} \in \mathbb{Z}_p$ and $pB_{k-2} \equiv -\llbracket p=3 \ \& \ k \neq 2 \rrbracket \pmod{p}$ by the von Staudt-Clausen theorem, therefore

$$\sum_{a=1}^{p-1} a^k \equiv pB_k - \llbracket p=3 \ \& \ k \neq 2 \rrbracket p \frac{k(k-1)}{2} \pmod{p^{\text{ord}_p(k)+2}}.$$

Putting $k = \varphi(p^b) > 2$, we obtain

$$\begin{aligned} \sum_{a=1}^{p-1} a^{\varphi(p^b)} &\equiv pB_{\varphi(p^b)} - \llbracket p=3 \rrbracket p \varphi(p^b) \frac{\varphi(p^b) - 1}{2} \\ &\equiv pB_{\varphi(p^b)} + \llbracket p=3 \rrbracket p^b \pmod{p^{b+1}}. \end{aligned}$$

(Note that if $p=3$ then $b > 1$ and hence $p \mid \varphi(p^b)$.) Thus

$$\begin{aligned} \sum_{a=1}^{p-1} \frac{a^{\varphi(p^b)} - 1}{p^b} &\equiv \frac{pB_{\varphi(p^b)} + \llbracket p=3 \rrbracket p^b - p + 1}{p^b} \\ &\equiv \frac{pB_{\varphi(p^b)} - p + 1}{p^b} + \llbracket p=3 \rrbracket \pmod{p}. \end{aligned}$$

If a_1 and a_2 are two integers relatively prime to p^b , then

$$\begin{aligned} \frac{(a_1 a_2)^{\varphi(p^b)} - 1}{p^b} &= \frac{a_1^{\varphi(p^b)} - 1}{p^b} a_2^{\varphi(p^b)} + \frac{a_2^{\varphi(p^b)} - 1}{p^b} \\ &\equiv \frac{a_1^{\varphi(p^b)} - 1}{p^b} + \frac{a_2^{\varphi(p^b)} - 1}{p^b} \pmod{p^b} \end{aligned}$$

by Euler's theorem. Therefore

$$\begin{aligned} \sum_{\substack{a=1 \\ p \nmid a}}^{p^b-1} \frac{a^{\varphi(p^b)} - 1}{p^b} &\equiv \frac{(\prod_{0 < a < p^b, p \nmid a} a)^{\varphi(p^b)} - 1}{p^b} = \frac{(-1 + p^b w_{p^b})^{\varphi(p^b)} - 1}{p^b} \\ &\equiv \frac{(1 - \varphi(p^b) p^b w_{p^b}) - 1}{p^b} = -\varphi(p^b) w_{p^b} \equiv p^{b-1} w_{p^b} \pmod{p^b}. \end{aligned}$$

Suppose that n is an integer with $0 < n < b$. Then

$$\begin{aligned} \sum_{\substack{a=1 \\ p \nmid a}}^{p^{n+1}-1} \frac{a^{\varphi(p^b)} - 1}{p^b} &= \sum_{\substack{a=1 \\ p \nmid a}}^{p^n-1} \sum_{s=0}^{p-1} \frac{(p^n s + a)^{\varphi(p^b)} - 1}{p^b} \\ &= \frac{1}{p^b} \sum_{\substack{a=1 \\ p \nmid a}}^{p^n-1} \sum_{s=0}^{p-1} \left(a^{\varphi(p^b)} - 1 + \sum_{k=1}^{\varphi(p^b)} \frac{\varphi(p^b)}{k} \binom{\varphi(p^b) - 1}{k-1} (p^n s)^k a^{\varphi(p^b)-k} \right) \\ &= p \sum_{\substack{a=1 \\ p \nmid a}}^{p^n-1} \frac{a^{\varphi(p^b)} - 1}{p^b} + (p-1) \sum_{k=1}^{\varphi(p^b)} \binom{\varphi(p^b) - 1}{k-1} \frac{p^{kn-1}}{k} \sum_{s=1}^{p-1} s^k \sum_{\substack{a=1 \\ p \nmid a}}^{p^n-1} a^{\varphi(p^b)-k}. \end{aligned}$$

If $n > 1$ and $k \geq 2$, then

$$\frac{p^{kn-1}}{k} = p^{k(n-1)+1} \frac{p^{k-2}}{k} \equiv 0 \pmod{p^{n+1}}$$

because $k(n-1) \geq 2(n-1) \geq n$ and $p^{k-2}/k \in \mathbb{Z}_p$ by [S03, Lemma 2.1]. In the case $n = 1$, as $p^{k-3}/k \in \mathbb{Z}_p$ for $k = 4, 5, \dots$ (cf. [S03, Lemma 2.1]) and

$$\frac{p^2}{3} \sum_{s=1}^{p-1} s^3 = \frac{p^2}{3} \sum_{s=1}^{(p-1)/2} (s^3 + (p-s)^3) \equiv 0 \pmod{p^2},$$

we have

$$\frac{p^{k-1}}{k} \sum_{s=1}^{p-1} s^k \equiv 0 \pmod{p^2} \quad \text{for } k = 3, 4, 5, \dots$$

Thus

$$\begin{aligned}
& \sum_{\substack{a=1 \\ p \nmid a}}^{p^{n+1}-1} \frac{a^{\varphi(p^b)} - 1}{p^b} - p \sum_{\substack{a=1 \\ p \nmid a}}^{p^n-1} \frac{a^{\varphi(p^b)} - 1}{p^b} \\
& \equiv (p-1)p^{n-1} \sum_{s=1}^{p-1} s \sum_{\substack{a=1 \\ p \nmid a}}^{p^n-1} a^{\varphi(p^b)-1} \\
& \quad + \llbracket n=1 \rrbracket (p-1)(\varphi(p^b) - 1) \frac{p}{2} \sum_{s=1}^{p-1} s^2 \sum_{a=1}^{p-1} a^{\varphi(p^b)-2} \\
& \equiv (p-1)p^{n-1} \frac{p(p-1)}{2} \sum_{\substack{a=1 \\ p \nmid a}}^{(p^n-1)/2} \left(\frac{1}{a} + \frac{1}{p^n - a} \right) \\
& \quad + \llbracket n=1 \rrbracket (p-1)(\varphi(p^b) - 1) \frac{p}{2} \cdot \frac{p(p-1)(2p-1)}{6} \sum_{a=1}^{p-1} a^{\varphi(p^b)-2} \\
& \equiv \llbracket p=3 \ \& \ n=1 \rrbracket \frac{p}{2} \times \frac{2 \cdot 3 \cdot 5}{6} \sum_{a=1}^2 1 \equiv -p \llbracket p=3 \ \& \ n=1 \rrbracket \pmod{p^{n+1}}.
\end{aligned}$$

In view of the above,

$$\begin{aligned}
& p^{b-1} \sum_{a=1}^{p-1} \frac{a^{\varphi(p^b)} - 1}{p^b} - \sum_{\substack{a=1 \\ p \nmid a}}^{p^b-1} \frac{a^{\varphi(p^b)} - 1}{p^b} \\
& = \sum_{0 < n < b} p^{b-n-1} \left(p \sum_{\substack{a=1 \\ p \nmid a}}^{p^n-1} \frac{a^{\varphi(p^b)} - 1}{p^b} - \sum_{\substack{a=1 \\ p \nmid a}}^{p^{n+1}-1} \frac{a^{\varphi(p^b)} - 1}{p^b} \right) \\
& \equiv \sum_{0 < n < b} p^{b-n-1} p \llbracket p=3 \ \& \ n=1 \rrbracket = p^{b-1} \llbracket p=3 \rrbracket \pmod{p^b}
\end{aligned}$$

and hence

$$p^{b-1} w_{p^b} \equiv \sum_{\substack{a=1 \\ p \nmid a}}^{p^b-1} \frac{a^{\varphi(p^b)} - 1}{p^b} \equiv p^{b-1} \sum_{a=1}^{p-1} \frac{a^{\varphi(p^b)} - 1}{p^b} - p^{b-1} \llbracket p=3 \rrbracket \pmod{p^b}.$$

Therefore

$$w_{p^b} \equiv \sum_{a=1}^{p-1} \frac{a^{\varphi(p^b)} - 1}{p^b} - \llbracket p=3 \rrbracket \equiv \frac{pB_{\varphi(p^b)} - p + 1}{p^b} \pmod{p}.$$

This concludes the proof. \square

Lemma 4.2. *Let p be an odd prime and let $n \in \mathbb{Z}^+$, $r \in \mathbb{Z}$ and $r \not\equiv 0 \pmod{p}$. Then, for any $b = 1, \dots, \text{ord}_p(pn)$, we have*

$$\begin{aligned} & \frac{r^n}{(-p)^N} \sum_{a=1}^{p-1} (a\pi)^{pn} \zeta_p^{-ar} - (-1)^{(b-1)n} \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k \\ & \equiv n_*! \left(p^b n_* H_{n_*} - n_* (pB_{\varphi(p^b)} - p + 1) - pnq_p(r) \right) \pmod{p^b \pi}, \end{aligned} \quad (4.2)$$

where

$$b' = \frac{p^b - 1}{p - 1} \quad \text{and} \quad N = \frac{pn + n^*}{p - 1} = \left\lfloor \frac{pn - 1}{p - 1} \right\rfloor + 1.$$

Proof. Write $pn = p^b m$ with $m \in \mathbb{Z}^+$. Then

$$\begin{aligned} r^{(p-1)n} &= (1 + pq_p(r))^{p^{b-1}m} \\ &= 1 + p^b m q_p(r) + \sum_{1 < k \leq p^{b-1}m} p^{b-1} m \binom{p^{b-1}m - 1}{k - 1} \frac{p^k}{k} q_p(r)^k \\ &\equiv 1 + p^b m q_p(r) \pmod{p^{b+1}} \end{aligned}$$

since $p^{k-2}/k \in \mathbb{Z}_p$ for $k = 2, 3, \dots$. Thus

$$\begin{aligned} \sum_{a=1}^{p-1} a^{pn} \zeta_p^{-ar} &= \frac{(-1)^{pn} r^{-n}}{r^{(p-1)n}} \sum_{a=1}^{p-1} (-ar)^{p^b m} \zeta_p^{-ar} \equiv \frac{(-1)^n r^{-n}}{1 + p^b m q_p(r)} \sum_{s=1}^{p-1} s^{p^b m} \zeta_p^s \\ &\equiv \frac{(-1)^n}{r^n} (1 - p^b m q_p(r)) \sum_{a=1}^{p-1} a^{pn} \zeta_p^a \pmod{p^{b+1}}. \end{aligned}$$

Let ω be the Teichmüller character of the multiplicative group

$$(\mathbb{Z}/p\mathbb{Z})^* = \{\bar{a} = a + p\mathbb{Z} : a = 1, \dots, p-1\}.$$

Then for each $a = 1, \dots, p-1$ the value $\omega(\bar{a})$ is just the unique $(p-1)$ -th root of unity (in the algebraic closure of \mathbb{Q}_p) with $\omega(\bar{a}) \equiv a \pmod{p}$. (See, e.g., [Wa, p. 51].) Since $a^{p^b} \equiv \omega(\bar{a})^{p^b} \pmod{p^{b+1}}$, we have

$$\sum_{a=1}^{p-1} a^{pn} \zeta_p^a \equiv \sum_{a=1}^{p-1} \omega(\bar{a})^{pn} \zeta_p^a = \sum_{a=1}^{p-1} \omega(\bar{a})^{-n^*} \zeta_p^a \pmod{p^{b+1}}.$$

By the Gross-Koblitz formula for Gauss sums (cf. [BEW, p. 350] and [GK]),

$$G(n^*) := \sum_{a=1}^{p-1} \omega(\bar{a})^{-n^*} \zeta_p^a = -\pi_0^{n^*} \Gamma_p \left(\frac{n^*}{p-1} \right)$$

where Γ_p is Morita's p -adic Γ -function (see [BEW, p. 277] or [Mu, p. 59 and pp. 67–70] for the definition and basic properties), and π_0 is the unique element in $\mathbb{Z}_p[\zeta_p]$ satisfying

$$\pi_0^{p-1} = -p \quad \text{and} \quad \pi_0 \equiv \zeta_p - 1 \pmod{(\zeta_p - 1)^2}.$$

(See [Go, pp. 172–173] for the existence of π_0 .) Clearly $\pi_0 \equiv \zeta_p - 1 \equiv \pi \pmod{\pi^2}$ and hence $\pi_0/\pi \equiv 1 \pmod{\pi}$. (Furthermore, we have $\pi_0/\pi = (\pi_0/\pi)^p \pi^{p-1}/(-p) \equiv 1 \pmod{p}$.)

In view of the above,

$$\sum_{a=1}^{p-1} a^{pn} \zeta_p^{-ar} \equiv \frac{(-1)^{n-1}}{r^n} \pi_0^{n^*} (1 - pnq_p(r)) \Gamma_p \left(\frac{n^*}{p-1} \right) \pmod{p^{b+1}}$$

and hence

$$\begin{aligned} & (-1)^{n-1} r^n \sum_{a=1}^{p-1} (a\pi)^{pn} \zeta_p^{-ar} \\ & \equiv \pi_0^{pn+n^*} \left(\frac{\pi}{\pi_0} \right)^{pn} (1 - pnq_p(r)) \Gamma_p \left(\frac{n^*}{p-1} \right) \pmod{p^{b+1} \pi^{pn}} \\ & \equiv (-p)^N \left(\frac{\pi}{\pi_0} \right)^{p^b m} (1 - pnq_p(r)) \Gamma_p \left(\frac{n^*}{p-1} \right) \pmod{p^b \pi^{pn+n^*+1}} \\ & \equiv (-p)^N (1 - pnq_p(r)) \Gamma_p \left(\frac{n^*}{p-1} \right) \pmod{p^{b+N} \pi}. \end{aligned}$$

(Note that $(\pi/\pi_0)^{p^b} \equiv 1 \pmod{p^b \pi}$.)

Since

$$\frac{n^*}{p-1} = 1 - \frac{n_*}{p-1} \equiv 1 + n_* \frac{p^{b+1} - 1}{p-1} = 1 + n_* + pn_* + \cdots + p^b n_* \pmod{p^{b+1}},$$

we have

$$\Gamma_p \left(\frac{n^*}{p-1} \right) \equiv \Gamma_p((p^b + b')n_* + 1) = (-1)^{(p^b + b')n_* + 1} \prod_{\substack{k=1 \\ p \nmid k}}^{(p^b + b')n_*} k \pmod{p^{b+1}}.$$

Observe that

$$\begin{aligned} \prod_{\substack{k=1 \\ p \nmid k}}^{(p^b + b')n_*} k &= \prod_{s=0}^{n_*-1} \prod_{\substack{t=1 \\ p \nmid t}}^{p^b-1} (p^b s + t) \times \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} (p^b n_* + k) \\ &= \left(\prod_{\substack{t=1 \\ p \nmid t}}^{p^b-1} t \right)^{n_*} \prod_{s=0}^{n_*-1} \prod_{\substack{t=1 \\ p \nmid t}}^{p^b-1} \left(1 + p^b \frac{s}{t} \right) \times \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k \times \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} \left(1 + p^b \frac{n_*}{k} \right) \end{aligned}$$

and hence

$$\begin{aligned}
& \left(\prod_{\substack{k=1 \\ p \nmid k}}^{(p^b+b')n_*} k \right) / \left(\left(\prod_{\substack{t=1 \\ p \nmid t}}^{p^b-1} t \right)^{n_*} \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k \right) \\
& \equiv 1 + p^b \sum_{s=0}^{n_*-1} \sum_{\substack{t=1 \\ p \nmid t}}^{p^b-1} \frac{s}{t} + p^b n_* \left(\sum_{0 < k < (b'-1)n_*} \frac{1}{k} + \sum_{j=1}^{n_*} \frac{1}{n_*(b'-1)+j} \right) \\
& \equiv 1 + p^b n_* \sum_{j=1}^{n_*} \frac{1}{j} = 1 + p^b n_* H_{n_*} \pmod{p^{b+1}}
\end{aligned}$$

since $p \mid b' - 1$ and $\sum_{k=1}^{p-1} 1/k = \sum_{0 < k < p/2} (1/k + 1/(p-k)) \equiv 0 \pmod{p}$.

By Lemma 4.1,

$$\begin{aligned}
\left(- \prod_{\substack{t=1 \\ p \nmid t}}^{p^b-1} t \right)^{n_*} &= (1 - p^b w_{p^b})^{n_*} \equiv 1 - n_* p^b w_{p^b} \\
&\equiv 1 - n_*(pB_{\varphi(p^b)} - p + 1) \pmod{p^{b+1}}.
\end{aligned}$$

Therefore

$$\begin{aligned}
& \Gamma_p \left(\frac{n_*}{p-1} \right) / \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k \\
& \equiv (-1)^{(p^b+b')n_*+1} (1 + p^b n_* H_{n_*}) (-1)^{n_*} (1 - n_*(pB_{\varphi(p^b)} - p + 1)) \\
& \equiv (-1)^{bn+1} (1 + p^b n_* H_{n_*} - n_*(pB_{\varphi(p^b)} - p + 1)) \pmod{p^{b+1}}.
\end{aligned}$$

Note that $b' - 1 = p \sum_{0 \leq i < b-1} p^i$ and hence

$$\begin{aligned}
\prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k &= \prod_{0 \leq s < n_*(b'-1)/p} \prod_{t=1}^{p-1} (ps+t) \times \prod_{j=1}^{n_*} (n_*(b'-1)+j) \\
&\equiv ((p-1)!)^{n_*(b'-1)/p} n_*! \equiv (-1)^{n(b-1)} n_*! \pmod{p}.
\end{aligned}$$

So there is a $u \in \mathbb{Z}$ such that

$$U := (-1)^{(b-1)n} \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k = n_*! + pu.$$

Combining the above, we finally get

$$\begin{aligned}
& \frac{r^n}{(-p)^N} \sum_{a=1}^{p-1} (a\pi)^{pn} \zeta_p^{-ar} \\
& \equiv (-1)^{n-1} (1 - pnq_p(r)) \Gamma_p \left(\frac{n_*}{p-1} \right) \\
& \equiv (-1)^{n-1} (1 - p^b m q_p(r)) \times (-1)^{bn+1} \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k \\
& \quad \times \left(1 + p^b n_* \left(H_{n_*} - \frac{pB_{\varphi(p^b)} - p + 1}{p^b} \right) \right) \\
& \equiv (n_*! + pu) \left(1 - p^b m q_p(r) + p^b n_* \left(H_{n_*} - \frac{pB_{\varphi(p^b)} - p + 1}{p^b} \right) \right) \\
& \equiv U + n_*! (-pnq_p(r) + p^b n_* H_{n_*} - n_* (pB_{\varphi(p^b)} - p + 1)) \pmod{p^b \pi}.
\end{aligned}$$

This yields the desired (4.2). \square

Proof of Theorem 1.3. Let $\bar{n} = pn$. By Lemma 3.1 and Theorem 2.1(ii),

$$\begin{aligned}
(-1)^{p\bar{n}} p C_p(p\bar{n}, r) &= \sum_{a=0}^{p-1} \zeta_p^{-ar} (\zeta_p^a - 1)^{p\bar{n}} \\
&\equiv \sum_{a=1}^{p-1} \zeta_p^{-ar} (a\pi)^{p\bar{n}} \pmod{p^{\text{ord}_p(p\bar{n})} \pi^{p\bar{n}}}.
\end{aligned}$$

As $b \leq \text{ord}_p(\bar{n})$ and $p\bar{n} - pn = (p-1)\bar{n} \equiv 0 \pmod{\varphi(p^{b+1})}$, we have

$$(-1)^n p C_p(p\bar{n}, r) \equiv \pi^{(p-1)\bar{n}} \sum_{a=1}^{p-1} (a\pi)^{pn} \zeta_p^{-ar} \pmod{p^{b+1} \pi^{p\bar{n}}}.$$

Set $N = (\bar{n} + n^*)/(p-1)$. Then

$$N + \bar{n} = \frac{p\bar{n} + n^*}{p-1} = \left\lfloor \frac{p\bar{n} - 1}{p-1} \right\rfloor + 1.$$

Thus

$$\frac{(-1)^n}{(-p)^N} \sum_{a=1}^{p-1} (a\pi)^{pn} \zeta_p^{-ar} = \frac{p C_p(p\bar{n}, r)}{(-p)^{N+\bar{n}}} \left(\frac{-p}{\pi^{p-1}} \right)^{\bar{n}} \equiv -F_p(p\bar{n}, r) \pmod{p^b \pi}.$$

(Note that $(-p/\pi^{p-1})^{\bar{n}} \equiv 1 \pmod{p^b\pi}$ since $-p/\pi^{p-1} \equiv 1 \pmod{\pi}$ and $p^b \mid \bar{n}$.) Combining this with Lemma 4.2, we obtain

$$\begin{aligned} & -(-r)^n F_p(p\bar{n}, r) - (-1)^{(b-1)n} \prod_{\substack{k=1 \\ p \nmid k}}^{b'n_*} k \\ & \equiv n_*! (p^b n_* H_{n_*} - n_*(pB_{\varphi(p^b)} - p + 1) - pnq_p(r)) \pmod{p^b\pi} \end{aligned}$$

and hence

$$\begin{aligned} & \frac{(-r)^n F_p(p\bar{n}, r) + (-1)^{(b-1)n} \prod_{1 \leq k \leq b'n_*, p \nmid k} k}{n_*!} \\ & \equiv n_*(pB_{\varphi(p^b)} - p + 1) - p^b n_* H_{n_*} + pnq_p(r) \pmod{p^{b+1}}. \end{aligned} \quad (4.3)$$

By Theorem 1.1,

$$\frac{F_p(p\bar{n}, r) - F_p(\bar{n}, r)}{(p-1)\bar{n}} \equiv (-1)^{n_*-1} \frac{n_*!}{r^n} \sum_{1 < k \leq n_*} \binom{n_* + k}{n_*} \frac{B_k}{kr^k} \pmod{p}$$

and so

$$\frac{(-r)^n (F_p(p\bar{n}, r) - F_p(\bar{n}, r))}{n_*!} \equiv \bar{n} \sum_{1 < k \leq n_*} \binom{n_* + k}{n_*} \frac{B_k}{kr^k} \pmod{p^{b+1}}. \quad (4.4)$$

From (4.3) and (4.4) we immediately get the desired congruence (1.17). \square

5. CONGRUENCES FOR EXTENDED FLECK QUOTIENTS

Let p be a prime, and let $a \in \mathbb{Z}^+$, $n \in \mathbb{N}$ and $r \in \mathbb{Z}$. In 1977 C. S. Weisman [We] extended Fleck's inequality by showing that

$$\text{ord}_p \left(\sum_{k \equiv r \pmod{p^a}} \binom{n}{k} (-1)^k \right) \geq \left\lfloor \frac{n - p^{a-1}}{\varphi(p^a)} \right\rfloor.$$

An extension of this result was given by the author in [S06]. During his study of the ψ -operator in Fontaine's theory in 2005, D. Wan finally obtained the following extension of Fleck's inequality (cf. [W] and [SW1]): For any $l \in \mathbb{N}$ we have

$$\text{ord}_p \left(\sum_{k \equiv r \pmod{p^a}} \binom{n}{k} (-1)^k \binom{(k-r)/p^a}{l} \right) \geq \left\lfloor \frac{n - lp^a - p^{a-1}}{\varphi(p^a)} \right\rfloor,$$

i.e., the extended Fleck quotient

$$F_{p^a}^{(l)}(n, r) := (-p)^{\lfloor (n - lp^a - p^{a-1})/\varphi(p^a) \rfloor} \sum_{k \equiv r \pmod{p^a}} \binom{n}{k} (-1)^k \binom{(k-r)/p^a}{l}$$

is an integer. In this section we study $F_{p^a}^{(l)}(n, r) \pmod{p}$.

Theorem 5.1. *Let p be a prime, and let $a \in \mathbb{Z}^+$, $l, n \in \mathbb{N}$, $r \in \mathbb{Z}$ and $s, t \in \{0, \dots, p^{a-1} - 1\}$. Let $m \in \mathbb{N}$ with $m \equiv -n \pmod{p}$. Then*

$$\begin{aligned} & (-1)^{l+t-1} F_{p^a}^{(l)}(p^{a-1}n + s, p^{a-1}r + t) \\ & \equiv \llbracket n > l \rrbracket \binom{s}{t} \binom{\lfloor (n-l-1)/(p-1) \rfloor}{l} (n-l)_* B_{(n-l)_*}^{(m)}(-r) \pmod{p}, \end{aligned} \quad (5.1)$$

provided that we have one of the following (i)–(iii):

- (i) $a = 1$ or $p \mid n$ or $p-1 \nmid n-l-1$;
- (ii) $\lfloor s/p^{a-2} \rfloor = 2 \lfloor t/p^{a-2} \rfloor$ and $p \neq 2$;
- (iii) $\lfloor s/p^{a-2} \rfloor = \lfloor t/p^{a-2} \rfloor = p-1$.

Now we deduce Theorem 1.4 from Theorem 5.1.

Proof of Theorem 1.4. Let $d \in \mathbb{Z}^+$ with $d \leq \max\{p^{a-2}, 1\}$. Then

$$\left\lfloor \frac{(p^a n - p^{a-1} m - d) - lp^a - p^{a-1}}{\varphi(p^a)} \right\rfloor = \left\lfloor \frac{p(n-l) - m - 2}{p-1} \right\rfloor = n-l.$$

Also, $\lfloor (p^{a-1} - d)/p^{a-2} \rfloor = p-1$ if $a > 1$. Thus

$$\begin{aligned} & \frac{1}{(-p)^{n-l}} \sum_{l < k \leq n} \binom{p^a n - p^{a-1} m - d}{p^a k - p^{a-1} m - d} (-1)^{p^a k - p^{a-1} m - d} \binom{k-1}{l} \\ & = F_{p^a}^{(l)}(p^a n - p^{a-1} m - d, p^a - p^{a-1} m - d) \\ & = F_{p^a}^{(l)}(p^{a-1}(pn - m - 1) + p^{a-1} - d, p^{a-1}(p - m - 1) + p^{a-1} - d) \\ & \equiv (-1)^{l+(p^{a-1}-d)-1} \binom{p^{a-1}-d}{p^{a-1}-d} \binom{\lfloor (pn-m-1-l-1)/(p-1) \rfloor}{l} \\ & \quad \times (pn-m-1-l)_*! B_{(pn-m-1-l)_*}^{(m+1)}(m+1-p) \quad (\text{by Theorem 5.1}) \\ & \equiv (-1)^{l+d} \binom{n}{l} (n-m-1-l)! B_{p-1-(n-m-1-l)}^{(m+1)}(m+1-p) \pmod{p}. \end{aligned}$$

Since $B_0^{(m+1)}, \dots, B_{p-2}^{(m+1)} \in \mathbb{Z}_p$ and

$$\begin{aligned} & (-1)^{p-n+l+m} B_{p-n+l+m}^{(m+1)}(m+1-p) = B_{p-n+l+m}^{(m+1)}(p) \\ & = \sum_{j=0}^{p-n+l+m} \binom{p-n+l+m}{j} B_j^{(m+1)} p^{p-n+l+m-j} \equiv B_{p-n+l+m}^{(m+1)} \pmod{p}, \end{aligned}$$

by the above we have

$$\begin{aligned} & \frac{1}{(-p)^{n-l}} \sum_{l < k \leq n} \binom{p^a n - p^{a-1} m - d}{p^a k - p^{a-1} m - d} (-1)^{p^a k - p^{a-1} m - d} \binom{k-1}{l} \\ & \equiv (-1)^{l+d} \frac{n!/l!}{\prod_{k=0}^m (n-l-k)} (-1)^{p-n+l+m} B_{p-n+l+m}^{(m+1)} \pmod{p}, \end{aligned}$$

which is equivalent to (1.20). \square

To prove Theorem 5.1 we need some lemmas.

Lemma 5.1. *Let $f(x)$ be a function from \mathbb{Z} to a field, and let $m, n \in \mathbb{Z}^+$. Then, for any $r \in \mathbb{Z}$ we have*

$$\sum_{k=0}^n \binom{n}{k} (-1)^k f\left(\left\lfloor \frac{k-r}{m} \right\rfloor\right) = \sum_{k \equiv \bar{r} \pmod{m}} \binom{n-1}{k} (-1)^{k-1} \Delta f\left(\frac{k-\bar{r}}{m}\right),$$

where $\bar{r} = r + m - 1$ and $\Delta f(x) = f(x+1) - f(x)$.

Proof. This is Lemma 2.1 of Sun [S06]. \square

Lemma 5.2. *Let p be a prime, and let $l, n \in \mathbb{N}$ with $n > p$. Then*

$$\begin{aligned} & F_p^{(l)}(n, r) + \llbracket l > 0 \rrbracket F_p^{(l-1)}(n-p, r) \\ & \equiv - \sum_{k=1}^{p-1} \frac{1}{k} \sum_{j=0}^{k-1} F_p^{(l)}(n-p+1, r-j) \pmod{p}. \end{aligned} \quad (5.2)$$

Proof. Set $n' = n - (p-1) > 0$. With help of the Chu-Vandermonde convolution identity,

$$\begin{aligned} & F_p^{(l)}(n, r) \\ & = (-p)^{-\lfloor (n-lp-1)/(p-1) \rfloor} \sum_{k \equiv r \pmod{p}} \sum_{j=0}^{p-1} \binom{p-1}{j} \binom{n'}{k-j} (-1)^k \binom{(k-r)/p}{l} \\ & = -\frac{1}{p} \sum_{j=0}^{p-1} \binom{p-1}{j} (-p)^{-\lfloor (n'-lp-1)/(p-1) \rfloor} \sum_{p|k-r} \binom{n'}{k-j} (-1)^k \binom{(k-r)/p}{l} \\ & = -\frac{1}{p} \sum_{j=0}^{p-1} \binom{p-1}{j} (-1)^j F_p^{(l)}(n', r-j). \end{aligned}$$

For any $j = 0, \dots, p-1$, clearly

$$\begin{aligned} & \binom{p-1}{j} (-1)^j = \prod_{0 < i \leq j} \left(1 - \frac{p}{i}\right) \\ & \equiv 1 - \sum_{0 < i \leq j} \frac{p}{i} \equiv (-1)^{p-1} + p \sum_{j < k < p} \frac{1}{k} \pmod{p^2}. \end{aligned}$$

(Recall that $H_{p-1} = \sum_{k=1}^{p-1} 1/k \equiv 0 \pmod{p}$ if $p \neq 2$.) Also,

$$\begin{aligned}
& -\frac{1}{p} \sum_{j=0}^{p-1} F_p^{(l)}(n', r-j) \\
&= (-p)^{-1 - \lfloor (n' - lp - 1)/(p-1) \rfloor} \sum_{k=0}^{n'} \binom{n'}{k} (-1)^k \binom{\lfloor (k - r + p - 1)/p \rfloor}{l} \\
&= (-p)^{-\lfloor ((n' - 1) - (l-1)p - 1)/(p-1) \rfloor} \sum_{k \equiv r \pmod{p}} \binom{n' - 1}{k} (-1)^{k-1} \binom{(k - r)/p}{l-1} \\
&= -\llbracket l > 0 \rrbracket F_p^{(l-1)}(n' - 1, r),
\end{aligned}$$

where we have applied Lemma 5.1 with $f(x) = \binom{x}{l}$ for the second equality and view $\binom{x}{-1}$ as 0. Therefore

$$\begin{aligned}
F_p^{(l)}(n, r) &\equiv (-1)^p \llbracket l > 0 \rrbracket F_p^{(l-1)}(n' - 1, r) - \sum_{j=0}^{p-1} \sum_{j < k < p} \frac{F_p^{(l)}(n', r-j)}{k} \\
&\equiv -\llbracket l > 0 \rrbracket F_p^{(l-1)}(n' - 1, r) - \sum_{k=1}^{p-1} \frac{1}{k} \sum_{j=0}^{k-1} F_p^{(l)}(n', r-j) \pmod{p}.
\end{aligned}$$

This proves (5.2). \square

Lemma 5.3. *Let p be a prime, and let $l, n \in \mathbb{N}$ and $r \in \mathbb{Z}$. If $n > lp$, then*

$$F_p^{(l)}(n, r) \equiv (-1)^l \binom{\lfloor (n - l - 1)/(p-1) \rfloor}{l} F_p(n - lp, r) \pmod{p}. \quad (5.3)$$

Proof. We use induction on $l + n$.

Clearly $l = 0$ and $n = 1$ if $l + n = 1$. In the case $l = 0$, (5.3) holds trivially for $n > 0$.

Below we let $l > 0$ and assume the corresponding result for smaller values of $l + n$. As $n > lp$, we have $n' - 1 > (l-1)p$ where $n' = n - p + 1$. By the induction hypothesis, $(-1)^{l-1} F_p^{(l-1)}(n' - 1, r)$ is congruent to

$$\begin{aligned}
& \binom{\lfloor (n' - 1 - (l-1) - 1)/(p-1) \rfloor}{l-1} F_p(n' - 1 - (l-1)p, r) \\
&= \binom{\lfloor (n' - l - 1)/(p-1) \rfloor}{l-1} F_p(n - lp, r)
\end{aligned}$$

modulo p .

Clearly $n' > lp - p + 1 \geq l$. If $n' \leq lp$ then

$$\frac{n' - l - 1}{p - 1} - l = \frac{n' - lp - 1}{p - 1} < 0$$

and

$$\sum_{k=1}^{p-1} \frac{1}{k} \sum_{j=0}^{k-1} (-1)^l F_p^{(l)}(n', r - j) \equiv 0 \pmod{p}.$$

If $n' > lp$, then by the induction hypothesis,

$$\begin{aligned} & \sum_{k=1}^{p-1} \frac{1}{k} \sum_{j=0}^{k-1} (-1)^l F_p^{(l)}(n', r - j) \\ & \equiv \sum_{k=1}^{p-1} \frac{1}{k} \sum_{j=0}^{k-1} \binom{\lfloor (n' - l - 1)/(p - 1) \rfloor}{l} F_p(n' - lp, r - j) \\ & \equiv - \binom{\lfloor (n' - l - 1)/(p - 1) \rfloor}{l} F_p(n - lp, r) \pmod{p}, \end{aligned}$$

where we have applied Lemma 5.2 for Fleck quotients.

The above, together with Lemma 5.2, yields that

$$\begin{aligned} (-1)^l F_p^{(l)}(n, r) & \equiv (-1)^{l-1} \llbracket l > 0 \rrbracket F_p^{(l-1)}(n' - 1, r) \\ & \quad - \sum_{k=1}^{p-1} \frac{1}{k} \sum_{j=0}^{k-1} (-1)^l F_p^{(l)}(n', r - j) \\ & \equiv \binom{\lfloor (n' - l - 1)/(p - 1) \rfloor}{l - 1} F_p(n - lp, r) \\ & \quad + \binom{\lfloor (n' - l - 1)/(p - 1) \rfloor}{l} F_p(n - lp, r) \\ & \equiv \binom{\lfloor (n - l - 1)/(p - 1) \rfloor}{l} F_p(n - lp, r) \pmod{p}. \end{aligned}$$

The induction proof is now complete. \square

Lemma 5.4. *Let p be a prime, and let $a \in \mathbb{Z}^+$, $n \in \mathbb{N}$, $r \in \mathbb{Z}$ and $s, t \in \{0, \dots, p^{a-1} - 1\}$. If one of (i)-(iii) in Theorem 5.1 is satisfied, then*

$$F_{p^a}^{(l)}(p^{a-1}n + s, p^{a-1}r + t) \equiv (-1)^t \binom{s}{t} F_p^{(l)}(n, r) \pmod{p} \quad (5.4)$$

Proof. (5.4) holds trivially in the case $a = 1$. Below we assume $a \geq 2$.

Write $s = \sum_{k=0}^{a-2} s_k p^k$ and $t = \sum_{k=0}^{a-2} t_k p^k$ with $s_k, t_k \in \{0, \dots, p-1\}$. By [SW1, Theorem 1.1], if $a > 2$ then

$$\begin{aligned}
& F_{p^a}^{(l)}(p^{a-1}n + s, p^{a-1}r + t) \\
&= F_{p^a}^{(l)}\left(p\left(p^{a-2}n + \sum_{k=1}^{a-2} s_k p^{k-1}\right) + s_0, p\left(p^{a-2}r + \sum_{k=1}^{a-2} t_k p^{k-1}\right) + t_0\right) \\
&\equiv (-1)^{t_0} \binom{s_0}{t_0} F_{p^{a-1}}^{(l)}\left(p^{a-2}n + \sum_{k=1}^{a-2} s_k p^{k-1}, p^{a-2}r + \sum_{k=1}^{a-2} t_k p^{k-1}\right) \\
&\equiv \dots \equiv \left(\prod_{k=0}^{a-3} (-1)^{t_k} \binom{s_k}{t_k}\right) F_{p^2}^{(l)}(pn + s_{a-2}, pr + t_{a-2}) \pmod{p}.
\end{aligned}$$

Observe that $s_{a-2} = \lfloor s/p^{a-2} \rfloor$ and $t_{a-2} = \lfloor t/p^{a-2} \rfloor$. If (i) or (ii) holds, then

$$F_{p^2}^{(l)}(pn + s_{a-2}, pr + t_{a-2}) \equiv (-1)^{t_{a-2}} \binom{s_{a-2}}{t_{a-2}} F_p^{(l)}(n, r) \pmod{p} \quad (5.5)$$

by [SW1, Theorem 1.2]. Suppose that (iii) holds (i.e., $s_{a-2} = t_{a-2} = p-1$) but (i) fails. By [SW1, Lemma 3.3],

$$\begin{aligned}
& (-1)^{\lfloor (pn + s_{a-2} - (n-1)p - 1)/(p-1) \rfloor} F_p^{(n-1)}(pn + s_{a-2}, t_{a-2}) \\
&\equiv (-1)^{n+t_{a-2}} n \binom{s_{a-2}}{t_{a-2}} \frac{\sigma}{p} \pmod{p},
\end{aligned}$$

where

$$\begin{aligned}
\sigma &= 1 + (-1)^p \frac{\prod_{i=2}^p (p(n-1) + p - 1 + i)}{\prod_{i=1}^{p-1} i} = 1 + (-1)^p \prod_{k=1}^{p-1} \left(1 + \frac{pn}{k}\right) \\
&\equiv 1 + (-1)^p \left(1 + pn \sum_{k=1}^{p-1} \frac{1}{k}\right) \equiv 0 \pmod{p^2}.
\end{aligned}$$

(Note that if $p = 2$ then n is odd since (i) fails.) Thus $F_p^{(n-1)}(pn + s_{a-2}, t_{a-2}) \equiv 0 \pmod{p}$ and hence (5.5) holds by [SW1, Lemma 3.2].

Provided (i) or (ii) or (iii), by the above we have

$$\begin{aligned}
& F_{p^a}^{(l)}(p^{a-1}n + s, p^{a-1}r + t) \\
&\equiv \prod_{k=0}^{a-2} (-1)^{t_k} \binom{s_k}{t_k} \times F_p^{(l)}(n, r) \\
&\equiv (-1)^{\sum_{k=0}^{a-2} t_k p^k} \binom{\sum_{k=0}^{a-2} s_k p^k}{\sum_{k=0}^{a-2} t_k p^k} F_p^{(l)}(n, r) = (-1)^t \binom{s}{t} F_p^{(l)}(n, r) \pmod{p},
\end{aligned}$$

where we have applied Lucas' theorem (cf. [HS]). This completes the proof. \square

Proof of Theorem 5.1. In view of Lemma 5.4, it suffices to show that $(-1)^l F_p^{(l)}(n, r)$ is congruent to

$$-[[n > l]] \binom{\lfloor (n-l-1)/(p-1) \rfloor}{l} (n-l)_* B_{(n-l)_*}^{(m)}(-r)$$

modulo p . In the case $n \leq lp$, this is easy since the last expression vanishes.

Below we assume $n > lp$. By Lemma 5.3 and (1.4),

$$\begin{aligned} & (-1)^l F_p^{(l)}(n, r) \\ & \equiv \binom{\lfloor (n-l-1)/(p-1) \rfloor}{l} F_p(n-lp, r) \\ & \equiv - \binom{\lfloor (n-l-1)/(p-1) \rfloor}{l} (n-lp)_*! B_{(n-lp)_*}^{(m)}(-r) \pmod{p}. \end{aligned}$$

Since $(n-lp)^* = (n-l)^*$ and $(n-lp)_* = (n-l)_*$, the desired result follows and we are done. \square

Acknowledgment. This paper is based on the previous work [SW2] joint with D. Wan. The author is indebted to Prof. Wan for his work in [SW2], and Prof. K. Ono for his comments on the author's related talk given at the University of Wisconsin at Madison in April 2006. The author also thanks his two Ph.D. students H. Pan and H. Q. Cao for discussion on a particular case of (1.20).

REFERENCES

- [BEW] B. C. Berndt, R. J. Evans and K. S. Williams, *Gauss and Jacobi Sums*, John Wiley & Sons, New York, 1998.
- [C] L. Carlitz, *Some congruences for the Bernoulli numbers*, Amer. J. Math. **75** (1953), 163–172.
- [Ch] S. Chowla, *On the class number of real quadratic fields*, Proc. Nat. Acad. Sci. U.S.A. **47** (1961), 878.
- [Co] P. Colmez, *Une correspondance de Langlands locale p -adique pour les représentations semi-stables de dimension 2*, preprint, 2004.
- [DS] D. M. Davis and Z. W. Sun, *A number-theoretic approach to homotopy exponents of $SU(n)$* , J. Pure Appl. Algebra, in press. Available from the website <http://arxiv.org/abs/math.AT/0508083>.
- [D] L. E. Dickson, *History of the Theory of Numbers*, Vol. I, AMS Chelsea Publ., 1999.
- [GL] I. M. Gessel and T. Lengyel, *On the order of Stirling numbers and alternating binomial coefficient sums*, Fibonacci Quart. **39** (2001), 444–454.
- [G1] J. W. L. Glaisher, *Congruences relating to the sums of product of the first n numbers and to other sums of product*, Quart. J. Math. **31** (1900), 1–35.

- [G2] J. W. L. Glaisher, *On the residues of the sums of products of the first $p - 1$ numbers, and their powers, to modulus p^2 or p^3* , *Quart. J. Math.* **31** (1900), 321–353.
- [Go] F. Q. Gouvêa, *p -adic Numbers: An Introduction*, 2nd ed., Springer, New York, 1997.
- [GK] B. Gross and N. Koblitz, *Gauss sums and the p -adic Γ -function*, *Annals of Math.* **109** (1979), 569–581.
- [GKP] R. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*, Addison-Wesley, New York, 1989.
- [HS] H. Hu and Z. W. Sun, *An extension of Lucas' theorem*, *Proc. Amer. Math. Soc.* **129** (2001), 3471–3478.
- [IR] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory* (Graduate texts in math.; 84), 2nd ed., Springer, New York, 1990.
- [M] L. J. Mordell, *The congruence $((p-1)/2)! \equiv \pm 1 \pmod{p}$* , *Amer. Math. Monthly* **68** (1961), 145–146.
- [Mu] M. R. Murty, *Introduction to p -adic Analytic Number Theory* (AMS/IP studies in adv. math.; vol. 27), Amer. Math. Soc., Providence, RI; Internat. Press, Somerville, MA, 2002.
- [PS] H. Pan and Z. W. Sun, *New identities involving Bernoulli polynomials*, *J. Combin. Theory Ser. A* **113** (2006), 156–175.
- [S02] Z. W. Sun, *On the sum $\sum_{k \equiv r \pmod{m}} \binom{n}{k}$ and related congruences*, *Israel J. Math.* **128** (2002), 135–156.
- [S03] Z. W. Sun, *General congruences for Bernoulli polynomials*, *Discrete Math.* **262** (2003), 253–276.
- [S06] Z. W. Sun, *Polynomial extension of Fleck's congruence*, *Acta Arith.* **122** (2006), 91–100.
- [SD] Z. W. Sun and D. M. Davis, *Combinatorial congruences modulo prime powers*, *Trans. Amer. Math. Soc.*, in press, <http://arxiv.org/abs/math.NT/0508087>.
- [SW1] Z. W. Sun and D. Wan, *Lucas type congruences for cyclotomic ψ -coefficients*, preprint, 2005. On-line version: <http://arxiv.org/abs/math.NT/0512012>.
- [SW2] Z. W. Sun and D. Wan, *On Fleck quotients*, preprint, 2006. On-line version: <http://arxiv.org/abs/math.NT/0512012>.
- [W] D. Wan, *Combinatorial congruences and ψ -operators*, *Finite Fields Appl.*, in press, <http://arxiv.org/abs/math.NT/0603462>.
- [Wa] L. C. Washington, *Introduction to Cyclotomic Fields* (Graduate texts in math.; 83), 2nd ed., Springer, New York, 1997.
- [We] C. S. Weisman, *Some congruences for binomial coefficients*, *Michigan Math. J.* **24** (1977), 141–151.